## **PCT**

(51) International Patent Classification 7.

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification /:	1	(11) International Publication Number	: WO 00/66178
A61K 48/00, C07H 21/04, C07K 14/00, 16/00, C12N 15/63, 15/85, 15/86	A1	(43) International Publication Date:	9 November 2000 (09.11.00)

(21) International Application Number:

PCT/US00/11664

(22) International Filing Date:

28 April 2000 (28.04.00)

(30) Priority Data:

60/132,021

30 April 1999 (30.04.99)

US

(71) Applicant (for all designated States except US): EMORY UNIVERSITY [US/US]; 1380 South Oxford Road, Atlanta, GA 30322 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BODEN, Scott, D. [US/US]; 2842 Cravey Drive, N.E., Atlanta, GA 30345 (US). HAIR, Gregory, A. [US/US]; 1887 Chrysler Drive, N.E., Atlanta, GA 30345 (US).

(74) Agents: \_GARRETT, Arthur, S. et\_al.; Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P., 1300 I Street, N.W., Washington, DC 20005-3315 (US).

(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CII, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GII, GM, KE, LS, MW, SD, SI., SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.

(54) Title: LIM MINERALIZATION PROTEIN SPLICE VARIANTS

#### (57) Abstract

The present invention is directed to isolated nucleic acid molecules that encode LIM mineralization protein, or LMP. The invention further provides vectors comprising splice variants of nucleotide sequences that encode LMP, as well as host cells comprising those vectors. Moreover, the present invention relates to methods of inducing bone formation by transfecting osteogenic precursor cells with an isolated nucleic acid molecule comprising a nucleotide sequence encoding splice variants of LIM mineralization protein. The transfection may occur ex vivo or in vivo by direct injection of virus or naked plasmid DNA. In a particular embodiment, the invention provides a method of fusing a spine by transfecting osteogenic precursor cells with an isolated nucleic acid molecule having a nucleotide sequence encoding LIM mineralization protein, admixing the transfected osteogenic precursor cells with a matrix and contacting the matrix with the spine. Finally, the invention relates to methods for inducing systemic bone formation by stable transfection of host cells with the vectors of the invention.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	CA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Мопасо	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
ВG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Vobago
BJ	Benin	115	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of Americ
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
СМ	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	ŞE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

### LIM Mineralization Protein Splice Variants

#### BACKGROUND OF THE INVENTION

#### 5 1. Field of the Invention

10

15

20

25

30

The field of the invention relates generally to osteogenic cells and the formation of bone and boney tissue in mammalian species. Specifically, the invention concerns a novel family of proteins, and nucleic acids encoding those proteins, that enhances the efficacy of bone mineralization *in vitro* and *in vivo*. The invention provides methods for treating a variety of pathological conditions associated with bone and boney tissue, such as, for example, spine fusion, fracture repair and osteoporosis.

#### 2. Description of the Related Art

Osteoblasts are thought to differentiate from pluripotent mesenchymal stem cells. The maturation of an osteoblast results in the secretion of an extracellular matrix which can mineralize and form bone. The regulation of this complex process is not well understood but is thought to involve a group of signaling glycoproteins known as bone morphogenetic proteins (BMPs). These proteins have been shown to be involved with embryonic dorsal-ventral patterning, limb bud development, and fracture repair in adult animals. B. L. Hogan, Genes & Develop., 10:1580 (1996). This group of transforming growth factor-beta superfamily secreted proteins has a spectrum of activities in a variety of cell types at different stages of differentiation; differences in physiological activity between these closely related molecules have not been clarified. D. M. Kingsley, Trends Genet., 10:16 (1994).

To better discern the unique physiological role of different BMP signaling proteins, we recently compared the potency of BMP-6 with that of BMP-2 and BMP-4, for inducing rat calvarial osteoblast differentiation. Boden *et al.*, Endocrinology, 137:3401 (1996). We studied this process in first passage (secondary) cultures of fetal rat calvaria that require BMP or glucocorticoid for

initiation of differentiation. In this model of membranous bone formation, glucocorticoid (GC) or a BMP will initiate differentiation to mineralized bone nodules capable of secreting osteocalcin, the osteoblast-specific protein. This secondary culture system is distinct from primary rat osteoblast cultures which undergo spontaneous differentiation. In this secondary system, glucocorticoid resulted in a ten-fold induction of BMP-6 mRNA and protein expression which was responsible for the enhancement of osteoblast differentiation. Boden *et al.*, Endocrinology, 138:2920 (1997).

In addition to extracellular signals, such as the BMPs, intracellular signals or regulatory molecules may also play a role in the cascade of events leading to formation of new bone. One broad class of intracellular regulatory molecules are the LIM proteins, which are so named because they possess a characteristic structural motif known as the LIM domain. The LIM domain is a cysteine-rich structural motif composed of two special zinc fingers that are joined by a 2-amino acid spacer. Some proteins have only LIM domains, while others contain a variety of additional functional domains. LIM proteins form a diverse group, which includes transcription factors and cytoskeletal proteins. The primary role of LIM domains appears to be in mediating protein-protein interactions, through the formation of dimers with identical or different LIM domains, or by binding distinct proteins.

In LIM homeodomain proteins, that is, proteins having both LIM domains and a homeodomain sequence, the LIM domains function as negative regulatory elements. LIM homeodomain proteins are involved in the control of cell lineage determination and the regulation of differentiation, although LIM-only proteins may have similar roles. LIM-only proteins are also implicated in the control of cell proliferation since several genes encoding such proteins are associated with oncogenic chromosome translocations.

Humans and other mammalian species are prone to diseases or injuries that require the processes of bone repair and/or regeneration. For example, treatment of fractures would be improved by new treatment regimens that could stimulate the natural bone repair mechanisms, thereby reducing the time

5

10

15

20

25

required for the fractured bone to heal. In another example, individuals afflicted with systemic bone disorders, such as osteoporosis, would benefit from treatment regimens that would results in systemic formation of new bone. Such treatment regimens would reduce the incidence of fractures arising from the loss of bone mass that is a characteristic of this disease.

For at least these reasons, extracellular factors, such as the BMPs, have been investigated for the purpose of using them to stimulate formation of new bone *in vivo*. Despite the early successes achieved with BMPs and other extracellular signalling molecules, their use entails a number of disadvantages. For example, relatively large doses of purified BMPs are required to enhance the production of new bone, thereby increasing the expense of such treatment methods. Furthermore, extracellular proteins are susceptible to degradation following their introduction into a host animal. In addition, because they are typically immunogenic, the possibility of stimulating an immune response to the administered proteins is ever present.

Due to such concerns, it would be desirable to have available treatment regimens that use an intracellular signalling molecule to induce new bone formation. Advances in the field of gene therapy now make it possible to introduce into osteogenic precursor cells, that is, cells involved in bone formation, or peripheral blood leukocytes, nucleotide fragments encoding intracellular signals that form part of the bone formation process. Gene therapy for bone formation offers a number of potential advantages: (1) lower production costs; (2) greater efficacy, compared to extracellular treatment regiments, due to the ability to achieve prolonged expression of the intracellular signal; (3) it would by-pass the possibility that treatment with extracellular signals might be hampered due to the presence of limiting numbers of receptors for those signals; (4) it permits the delivery of transfected potential osteoprogenitor cells directly to the site where localized bone formation is required; and (5) it would permit systemic bone formation, thereby providing a treatment regimen for osteoporosis and other metabolic bone diseases.

5

10

15

20

25

#### SUMMARY OF THE INVENTION

The present invention seeks to overcome the drawbacks in the prior art by providing novels compositions and methods for inducing bone formation using an intracellular signalling molecule that participates early in the cascade of events that leads to bone formation. Applicants have discovered 10-4/RLMP (SEQ ID NO: 1, SEQ ID NO: 2), a novel LIM gene with a sequence originally isolated from stimulated rat calvarial osteoblast cultures. The gene has been cloned, sequenced and assayed for its ability to enhance the efficacy of bone mineralization in vitro. The protein RLMP affects mineralization of bone matrix as well as differentiation of cells into the osteoblast lineage. Unlike other known cytokines, for example, BMPs, RLMP is not a secreted protein, but is instead an intracellular signaling molecule. This feature has the advantage of providing intracellular signaling amplification as well as easier assessment of transfected cells. It is also suitable for more efficient and specific in vivo applications. Suitable clinical applications include enhancement of bone repair in fractures, bone defects, bone grafting, and normal homeostasis in patients presenting with osteoporosis.

Applicants have also cloned, sequenced and deduced the amino acid sequence of a corresponding human protein, named human LMP-1. The human protein demonstrates enhanced efficacy of bone mineralization *in vitro* and *in vivo*.

In addition, the applicants have characterized a truncated (short) version of LMP-1, termed HLMP-1s. This short version resulted from a point mutation in one source of a cDNA clone, providing a stop codon which truncated the protein. The short version (LMP-1s) is fully functional when expressed in cell culture and *in vivo*.

Using PCR analysis of human heart cDNA library, Applicants have identified two alternative splice variants (referred to as HLMP-2 and HLMP-3) that differ from HLMP-1 in a region between base pairs 325 and 444 in the

10

15

20

25

nucleotide sequence encoding HLMP-1. The HLMP-2 sequence has a 119 base pair deletion and an insertion of 17 base pairs in this region. Compared to HLMP-1, the nucleotide sequence encoding HLMP-3 has no deletions, but it does have the same 17 base pairs as HLMP-2, which are inserted at position 444 in the HLMP-1 sequence.

Additional features and advantages of the invention will be set forth in the description which follows, and in part will be apparent from the description, or may be learned by practice of the invention. The objectives and other advantages of the invention will be realized and attained by the methods and compositions of matter particularly pointed out in the written description and claims hereof.

In one broad aspect, the invention relates to an isolated nucleic acid molecule comprising a nucleic acid sequence encoding any LIM mineralization protein, wherein the nucleic acid molecule hybridizes under standard conditions to a nucleic acid molecule complementary to the full length of SEQ. ID NO: 25, and wherein the molecule hybridizes under highly stringent conditions to a nucleic acid molecule complementary to the full length of SEQ. ID NO: 26. In a specific aspect, the isolated nucleic acid molecule encodes HLMP-1, HLMP-1s, RLMP, HLMP-2, or HLMP-3. In addition, the invention is directed to vectors comprising these nucleic acid molecules, as well as host cells comprising the vectors. In another specific aspect, the invention relates to the proteins themselves.

In a second broad aspect, the invention relates to antibody that is specific for LIM mineralization protein, including HLMP-1, HLMP-1s, RLMP, HLMP-2, and HLMP-3. In one specific aspect, the antibody is a polyclonal antibody. In another specific aspect, the antibody is a monoclonal antibody.

In a third broad aspect, the invention relates to method of inducing bone formation wherein osteogenic precursor cells are transfected with an isolated nucleic acid molecule comprising a nucleotide sequence encoding LIM mineralization protein. In one specific aspect, the isolated nucleic acid molecule is in a vector, which may be a plasmid or a virus, such as adenovirus

5

10

15

20

25

or retrovirus. The transfection may occur *ex vivo* or *in vivo* by direct injection of the isolated nucleic acid molecule. The transfected isolated nucleic acid molecule may encode HLMP-1, HLMP-1s, RLMP, HLMP-2, or HLMP-3.

In a further aspect, the invention relates to methods of fusing a spine by transfecting osteogenic precursor cells with an isolated nucleic acid molecule having a nucleotide sequence encoding LIM mineralization protein, admixing the transfected osteogenic precursor cells with a matrix and contacting the matrix with the spine.

In yet another aspect, the invention relates to methods for inducing systemic bone formation by stable transfection of host cells with the vectors of the invention.

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory and are intended to provide further explanation of the invention as claimed.

15

10

5

#### ABBREVIATIONS AND DEFINITIONS

	ВМР	Bone Morphogenetic Protein
	HLMP-1	Human LMP-1, also
20		designated as Human LIM
		Protein or HLMP
	HLMP-1s	Human LMP-1 Short
		(truncated) protein
	HLMPU	Human LIM Protein Unique
25		Region
	LMP	LIM mineralization protein
	MEM	Minimal essential medium
	Trm	Triamcinolone
	-GlyP	Beta-glycerolphosphate
30	RACE	Rapid Amplification of cDNA
		Ends

		RLMP	Rat LIM mineralization protein,
			also designated as RLMP-1
		RLMPU	Rat LIM Protein Unique
			Region
5		RNAsin	RNase inhibitor
		ROB	Rat Osteoblast
		10-4	Clone containing cDNA
			sequence for RLMP (SEQ ID
			NO: 2)
10		UTR	Untranslated Region
			•
		HLMP-2	Human LMP Splice Variant 2
	A		
		HLMP-3	Human LMP Splice Variant 3
15			

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to novel mammalian LIM proteins, herein designated LIM mineralization proteins, or LMP. The invention relates more particularly to human LMP, known as HLMP or HLMP-1, or alternative splice variants of human LMP, which are known as HLMP-2 or HLMP-3. The applicants have discovered that these proteins enhance bone mineralization in mammalian cells grown *in vitro*. When produced in mammals, LMP also induces bone formation *in vivo*.

Ex vivo transfection of bone marrow cells, osteogenic precursor cells, peripheral blood leukocytes, or mesenchymal stem cells with nucleic acid that encodes LMP or HLMP, followed by reimplantation of the transfected cells in the donor, is suitable for treating a variety of bone-related disorders or injuries. For example, one can use this method to: augment long bone fracture repair; generate bone in segmental defects; provide a bone graft substitute for

20

25

fractures; facilitate tumor reconstruction or spine fusion; and provide a local treatment (by injection) for weak or osteoporotic bone, such as in osteoporosis of the hip, vertebrae, or wrist. Transfection with LMP or HLMP-encoding nucleic acid is also useful in: the percutaneous injection of transfected marrow cells to accelerate the repair of fractured long bones; treatment of delayed union or non-unions of long bone fractures or pseudoarthrosis of spine fusions; and for inducing new bone formation in avascular necrosis of the hip or knee.

In addition to *ex vivo*-based methods of gene therapy, transfection of a recombinant DNA vector comprising a nucleic acid sequence that encodes LMP or HLMP can be accomplished *in vivo*. When a DNA fragment that encodes LMP or HLMP is inserted into an appropriate viral vector, for example, an adenovirus vector, the viral construct can be injected directly into a body site were endochondral bone formation is desired. By using a direct, percutaneous injection to introduce the LMP or HLMP sequence stimulation of bone formation can be accomplished without the need for surgical intervention either to obtain bone marrow cells (to transfect *ex vivo*) or to reimplant them into the patient at the site where new bone is required. Alden *et al.*, Neurosurgical Focus (1998), have demonstrated the utility of a direct injection method of gene therapy using a cDNA that encodes BMP-2, which was cloned into an adenovirus vector.

It is also possible to carry out *in vivo* gene therapy by directly injecting into an appropriate body site, a naked, that is, unencapsulated, recombinant plasmid comprising a nucleic acid sequence that encodes HLMP. In this embodiment of the invention, transfection occurs when the naked plasmid DNA is taken up, or internalized, by the appropriate target cells, which have been described. As in the case of *in vivo* gene therapy using a viral construct, direct injection of naked plasmid DNA offers the advantage that little or no surgical intervention is required. Direct gene therapy, using naked plasmid DNA that encodes the endothelial cell mitogen VEGF (vascular endothelial growth factor), has been successfully demonstrated in human patients. Baumgartner *et al.*, Circulation, 97(12):1114-23 (1998).

10

15

20

25

By using an adenovirus vector to deliver LMP into osteogenic cells, transient expression of LMP is achieved. This occurs because adenovirus does not incorporate into the genome of target cells that are transfected. Transient expression of LMP, that is, expression that occurs during the lifetime of the transfected target cells, is sufficient to achieve the objects of the invention. Stable expression of LMP, however, can occur when a vector that incorporates into the genome of the target cell is used as a delivery vehicle. Retrovirus-based vectors, for example, are suitable for this purpose.

Stable expression of LMP is particularly useful for treating various systemic bone-related disorders, such as osteoporosis and osteogenesis imperfecta. For this embodiment of the invention, in addition to using a vector that integrates into the genome of the target cell to deliver an LMP-encoding nucleotide sequence into target cells, LMP expression is placed under the control of a regulatable promoter. For example, a promoter that is turned on by exposure to an exogenous inducing agent, such as tetracycline, is suitable. Using this approach, one can stimulate formation of new bone on a systemic basis by administering an effective amount of the exogenous inducing agent. Once a sufficient quantity of bone mass is achieved, administration of the exogenous inducing agent is discontinued. This process may be repeated as needed to replace bone mass lost, for example, as a consequence of osteoporosis.

Antibodies specific for HLMP are particularly suitable for use in methods for assaying the osteoinductive, that is, bone-forming, potential of patient cells. In this way one can identify patients at risk for slow or poor healing of bone repair. Also, HLMP-specific antibodies are suitable for use in marker assays to identify risk factors in bone degenerative diseases, such as, for example, osteoporosis.

Following well known and conventional methods, the genes of the present invention are prepared by ligation of nucleic acid segments that encode LMP to other nucleic acid sequences, such as cloning and/or expression vectors. Methods needed to construct and analyze these

5

10

15

20

25

recombinant vectors, for example, restriction endonuclease digests, cloning protocols, mutagenesis, organic synthesis of oligonucleotides and DNA sequencing, have been described. For DNA sequencing DNA, the dieoxyterminator method is the preferred.

Many treatises on recombinant DNA methods have been published, including Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, 2nd edition (1988), Davis et al., Basic Methods in Molecular Biology, Elsevier (1986), and Ausubel et al., Current Protocols in Molecular Biology, Wiley Interscience (1988). These reference manuals are specifically incorporated by reference herein.

Primer-directed amplification of DNA or cDNA is a common step in the expression of the genes of this invention. It is typically performed by the polymerase chain reaction (PCR). PCR is described in U.S. Patent No. 4,800,159 to Mullis *et al.* and other published sources. The basic principle of PCR is the exponential replication of a DNA sequence by successive cycles of primer extension. The extension products of one primer, when hybridized to another primer, becomes a template for the synthesis of another nucleic acid molecule. The primer-template complexes act as substrate for DNA polymerase, which in performing its replication function, extends the primers. The conventional enzyme for PCR applications is the thermostable DNA polymerase isolated from *Thermus aquaticus*, or Taq DNA polymerase.

Numerous variations of the basic PCR method exist, and a particular procedure of choice in any given step needed to construct the recombinant vectors of this invention is readily performed by a skilled artisan. For example, to measure cellular expression of 10-4/RLMP, RNA is extracted and reverse transcribed under standard and well known procedures. The resulting cDNA is then analyzed for the appropriate mRNA sequence by PCR.

The gene encoding the LIM mineralization protein is expressed in an expression vector in a recombinant expression system. Of course, the constructed sequence need not be the same as the original, or its complimentary sequence, but instead may be any sequence determined by the

5

10

15

20

25

degeneracy of the DNA code that nonetheless expresses an LMP having bone forming activity. Conservative amino acid substitutions, or other modifications, such as the occurrance of an amino-terminal methionine residue, may also be employed.

A ribosome binding site active in the host expression system of choice is ligated to the 5' end of the chimeric LMP coding sequence, forming a synthetic gene. The synthetic gene can be inserted into any one of a large variety of vectors for expression by ligating to an appropriately linearized plasmid. A regulatable promoter, for example, the *E. coli* lac promoter, is also suitable for the expression of the chimeric coding sequences. Other suitable regulatable promoters include trp, tac, recA, T7 and lambda promoters.

DNA encoding LMP is transfected into recipient cells by one of several standard published procedures, for example, calcium phosphate precipitation, DEAE-Dextran, electroporation or protoplast fusion, to form stable transformants. Calcium phosphate precipitation is preferred, particularly when performed as follows.

DNAs are coprecipitated with calcium phosphate according to the method of Graham and Van Der, Virology, 52:456 (1973), before transfer into cells. An aliquot of 40-50 g of DNA, with salmon sperm or calf thymus DNA as a carrier, is used for  $0.5 \times 10^8$  cells plated on a 100 mm dish. The DNA is mixed with 0.5 ml of 2X Hepes solution (280 mM NaCl, 50 mM Hepes and 1.5 mM Na<sub>2</sub>HPO<sub>4</sub>, pH 7.0), to which an equal volume of 2x CaCl<sub>2</sub> (250 mM CaCl<sub>2</sub> and 10 mM Hepes, pH 7.0) is added. A white granular precipitate, appearing after 30-40 minutes, is evenly distributed dropwise on the cells, which are allowed to incubate for 4-16 hours at 37°C. The medium is removed and the cells shocked with 15% glycerol in PBS for 3 minutes. After removing the glycerol, the cells are fed with Dulbecco's Minimal Essential Medium (DMEM) containing 10% fetal bovine serum.

DNA can also be transfected using: the DEAE-Dextran methods of Kimura *et al.*, <u>Virology</u>, 49:394 (1972) and Sompayrac *et al.*, <u>Proc. Natl. Acad.</u> <u>Sci. USA</u>, 78:7575 (1981); the electroporation method of Potter, <u>Proc. Natl.</u>

5

10

15

20

25

Acad. Sci. USA, 81:7161 (1984); and the protoplast fusion method of Sandri-Goddin et al., Molec. Cell. Biol., 1:743 (1981).

Phosphoramidite chemistry in solid phase is the preferred method for the organic synthesis of oligodeoxynucleotides and polydeoxynucleotides. In addition, many other organic synthesis methods are available. Those methods are readily adapted by those skilled in the art to the particular sequences of the invention.

The present invention also includes nucleic acid molecules that hybridize under standard conditions to any of the nucleic acid sequences encoding the LIM mineralization proteins of the invention. "Standard hybridization conditions" will vary with the size of the probe, the background and the concentration of the nucleic acid reagents, as well as the type of hybridization, for example, *in situ*, Southern blot, or hybrization of DNA-RNA hybrids (Northern blot). The determination of "standard hybridization conditions" is within the level of skill in the art. For example, see U.S. Patent 5,580,775 to Fremeau *et al.*, herein incorporated by reference for this purpose. See also, Southern, E. M., J. Mol. Biol., 98:503 (1975), Alwine *et al.*, Meth. Enzymol., 68:220 (1979), and Sambrook *et al.*, Molecular Cloning: A laboratory Manual, 2nd edition, pp. 7.19-7.50, Cold Spring Harbor Press (1989).

One preferred set of standard hybrization conditions involves a blot that is prehybridized at 42°C for 2 hours in 50% formamide, 5X SSPE (150 nM NaCl, 10 mM Na H<sub>2</sub>PO<sub>4</sub> [pH 7.4], 1 mM EDTA [pH 8.0]), 5X Denhardt's solution (20 mg Ficoll, 20 mg polyvinylpyrrolidone and 20 mg BSA per 100 ml water), 10% dextran sulphate, 1% SDS and 100 g/ml salmon sperm DNA. A <sup>32</sup>P-labelled cDNA probe is added, and hybridization is continued for 14 hours. Afterward, the blot is washed twice with 2X SSPE, 0.1% SDS for 20 minutes at 22°C, followed by a 1 hour wash at 65°C in 0.1X SSPE, 0.1 %SDS. The blot is then dried and exposed to x-ray film for 5 days in the presence of an intensifying screen.

5

15

20

25

Under "highly stringent conditions," a probe will hybridize to its target sequence if those two sequences are substantially identical. As in the case of standard hybridization conditions, one of skill in the art can, given the level of skill in the art and the nature of the particular experiment, determine the conditions under which only susbstantially identical sequences will hybridize.

Another aspect of the invention includes the proteins encoded by the nucleic acid sequences. In still another embodiment, the inventon relates to the identification of such proteins based on anti-LMP antibodies. In this embodiment, protein samples are prepared for Western blot analysis by lysing cells and separating the proteins by SDS-PAGE. The proteins are transferred to nitrocellulose by electroblotting as described by Ausubel *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons (1987). After blocking the filter with instant nonfat dry milk (1 gm in 100 ml PBS), anti-LMP antibody is added to the filter and incubated for 1 hour at room temperature. The filter is washed thoroughly with phosphate buffered saline (PBS) and incubated with horseradish peroxidase (HRPO)-antibody conjugate for 1 hour at room temperature. The filter is again washed thoroughly with PBS and the antigen bands are identified by adding diaminobenzidine (DAB).

Monospecific antibodies are the reagent of choice in the present invention, and are specifically used to analyze patient cells for specific characteristics associated with the expression of LMP. "Monospecific antibody" as used herein is defined as a single antibody species or multiple antibody species with homogenous binding characteristics for LMP. "Homogeneous binding" as used herein refers to the ability of the antibody species to bind to a specific antigen or epitope, such as those associated with LMP, as described above. Monospecific antibodies to LMP are purified from mammalian antisera containing antibodies reactive against LMP or are prepared as monoclonal antibodies reactive with LMP using the technique of Kohler and Milstein, Nature, 256:495-97 (1975). The LMP specific antibodies are raised by immunizing animals such as, for example, mice, rats, guinea

5

10

15

20

25

pigs, rabbits, goats or horses, with an appropriate concentration of LMP either with or without an immune adjuvant.

In this process, preimmune serum is collected prior to the first immunization. Each animal receives between about 0.1 mg and about 1000 mg of LMP associated with an acceptable immune adjuvant, if desired. Such acceptable adjuvants include, but are not limited to, Freund's complete, Freund's incomplete, alum-precipitate, water in oil emulsion containing Corynebacterium parvum and tRNA adjuvants. The initial immunization consists of LMP in, preferably, Freund's complete adjuvant injected at multiple sites either subcutaneously (SC), intraperitoneally (IP) or both. Each animal is bled at regular intervals, preferably weekly, to determine antibody titer. The animals may or may not receive booster injections following the initial immunization. Those animals receiving booster injections are generally given an equal amount of the antigen in Freund's incomplete adjuvant by the same route. Booster injections are given at about three week intervals until maximal titers are obtained. At about 7 days after each booster immunization or about weekly after a single immunization, the animals are bled, the serum collected, and aliquots are stored at about -20° C.

Monoclonal antibodies (mAb) reactive with LMP are prepared by immunizing inbred mice, preferably Balb/c mice, with LMP. The mice are immunized by the IP or SC route with about 0.1 mg to about 10 mg, preferably about 1 mg, of LMP in about 0.5 ml buffer or saline incorporated in an equal volume of an acceptable adjuvant, as discussed above. Freund's complete adjuvant is preferred. The mice receive an initial immunization on day 0 and are rested for about 3-30 weeks. Immunized mice are given one or more booster immunizations of about 0.1 to about 10 mg of LMP in a buffer solution such as phosphate buffered saline by the intravenous (IV) route. Lymphocytes from antibody-positive mice, preferably splenic lymphocytes, are obtained by removing the spleens from immunized mice by standard procedures known in the art. Hybridoma cells are produced by mixing the splenic lymphocytes with an appropriate fusion partner, preferably myeloma cells, under conditions

10

15

20

25

which will allow the formation of stable hybridomas. Fusion partners may include, but are not limited to: mouse myelomas P3/NS1/Ag 4-1; MPC-11; S-194 and Sp 2/0, with Sp 2/0 being preferred. The antibody producing cells and myeloma cells are fused in polyethylene glycol, about 1000 mol. wt., at concentrations from about 30% to about 50%. Fused hybridoma cells are selected by growth in hypoxanthine, thymidine and aminopterin in supplemented Dulbecco's Modified Eagles Medium (DMEM) by procedures known in the art. Supernatant fluids are collected from growth positive wells on about days 14, 18, and 21, and are screened for antibody production by an immunoassay such as solid phase immunoradioassay (SPIRA) using LMP as the antigen. The culture fluids are also tested in the Ouchterlony precipitation assay to determine the isotype of the mAb. Hybridoma cells from antibody positive wells are cloned by a technique such as the soft agar technique of MacPherson, "Soft Agar Techniques", in Tissue Culture Methods and Applications, Kruse and Paterson (eds.), Academic Press (1973). See, also, Harlow et al., Antibodies: A Laboratory Manual, Cold Spring Laboratory (1988).

Monoclonal antibodies may also be produced *in vivo* by injection of pristane- primed Balb/c mice, approximately 0.5 ml per mouse, with about 2x10<sup>6</sup> to about 6x10<sup>6</sup> hybridoma cells about 4 days after priming. Ascites fluid is collected at approximately 8-12 days after cell transfer and the monoclonal antibodies are purified by techniques known in the art.

In vitro production in anti-LMP mAb is carried out by growing the hydridoma cell line in DMEM containing about 2% fetal calf serum to obtain sufficient quantities of the specific mAb. The mAb are purified by techniques known in the art.

Antibody titers of ascites or hybridoma culture fluids are determined by various serological or immunological assays, which include, but are not limited to, precipitation, passive agglutination, enzyme-linked immunosorbent antibody (ELISA) technique and radioimmunoassay (RIA) techniques. Similar assays

5

10

15

20

25

are used to detect the presence of the LMP in body fluids or tissue and cell extracts.

It is readily apparent to those skilled in the art that the above described methods for producing monospecific antibodies may be utilized to produce antibodies specific for polypeptide fragments of LMP, full-length nascent LMP polypeptide, or variants or alleles thereof.

In another embodiment, the invention is directed to alternative splice variants of HLMP-1. PCR analysis of human heart cDNA revealed mRNA for two HLMP alternative splice variants, named HLMP-2 and HLMP-3, that differ from HLMP-1 in a region between base pairs 325 and 444 in the hLMP-1 sequence. The HLMP-2 sequence has a 119 base pair deletion and an insertion of 17 base pairs in this region. These changes preserve the reading frame, resulting in a 423 amino acid protein, which compared to HLMP-1, has a net loss of 34 amino acids (40 amino acids deleted plus 6 inserted amino acids). HLMP-2 contains the c-terminal LIM domains that are present in HLMP-1.

Compared to HLMP-1, HLMP-3 has no deletions, but it does have the same 17 base pair insertion at position 444. This insertion shifts the reading frame, causing a stop codon at base pairs 459-461. As a result, HLMP-3 encodes a protein of 153 amino acids. This protein lacks the c-terminal LIM domains that are present in HLMP-1 and HLMP-2. The predicted size of the proteins encoded by HLMP-2 and HLMP-3 was confirmed by western blot analysis.

PCR analysis of the tissue distribution of the three splice variants revealed that they are differentially expressed, with specific isoforms predominating in different tissues. HLMP-1 is apparently the predominant form expressed in leukocytes, spleen, lung, placenta, and fetal liver. HLMP-2 appears to be the predominant isoform in skeletal muscle, bone marrow, and heart tissue. HLMP-3, however, was not the predominant isoform in any tissue examined.

5

10

15

20

25

Overexpression of HLMP-3 in secondary rat osteoblast cultures induced bone nodule formation (287±56) similar to the effect seen for glucicorticoid (272±7) and HLMP-1 (232±200). Since HLMP-3 lacks the C-terminal LIM domains, there regions are not required for osteoinductive activity. Overexpression of HLMP-2, however, did not induce nodule formation (11±3). These data suggest that the amino acids encoded by the deleted 119 base pairs are necessary for osteoinduction. The data also suggest that the distribution of HLMP splice variants may be important for tissue-specific function. Surprisingly, we have shown that HLMP-2 inhibits steroid-induced osteoblast formation in secondary rat osteoblast cultures. Therefore, HLMP-2 will have therapeutic utility in clinical situations where bone formation is not desirable.

On July 22, 1997, a sample of 10-4/RLMP in a vector designated pCMV2/RLMP (which is vector pRc/CMV2 with insert 10-4 clone/RLMP) was deposited with the American Type Culture Collection (ATCC), 12301 Parklawn 15 Drive, Rockville, MD 20852. The culture accession number for that deposit is 209153. On March 19, 1998, a sample of the vector pHis-A with insert HLPM-1s was deposited at the American Type Culture Collection ("ATCC"). The culture accession number for that deposit is 209698. On April 14, 2000, samples of plasmids pHAhLMP-2 (vector pHisA with cDNA insert derived from 20 human heart muscle cDNA with HLMP-2) and pHAhLMP-3 (vector pHisA with cDNA insert derived from human heart muscle cDNA with HLMP-3) were deposited with the ATCC, 10801 University Blvd., Manassas, VA, 20110-2209, USA, under the conditions of the Budapest treaty. The accession numbers for these deposits are and , respectively. These deposits, as 25 required by the Budapest Treaty, will be maintained in the ATCC for at least 30 years and will be made available to the public upon the grant of a patent disclosing them. It should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by government action. 30

5

In assessing the nucleic acids, proteins, or antibodies of the invention, enzyme assays, protein purification, and other conventional biochemical methods are employed. DNA and RNA are analyzed by Southern blotting and Northern blotting techniques, respectively. Typically, the samples analyzed are size fractionated by gel electrophoresis. The DNA or RNA in the gels are then transferred to nitrocellulose or nylon membranes. The blots, which are replicas of sample patterns in the gels, were then hybridized with probes. Typically, the probes are radiolabelled, preferably with <sup>32</sup>P, although one could label the probes with other signal-generating molecules known to those in the art. Specific bands of interest can then be visualized by detection systems, such as autoradiography.

For purposes of illustrating preferred embodiments of the present invention, the following, non-limiting examples are included. These results demonstrate the feasibility of inducing or enhancing the formation of bone using the LIM mineralization proteins of the invention, and the isolated nucleic acid molecules encoding those proteins.

## Example 1: Calvarial Cell Culture

Rat calvarial cells, also known as rat osteoblasts ("ROB"), were obtained from 20-day pre-parturition rats as previously described. Boden *et al.*, Endocrinology, 137(8):3401-07 (1996). Primary cultures were grown to confluence (7 days), trypsinized, and passed into 6-well plates (1 x 10<sup>5</sup> cells/35 mm well) as first subculture cells. The subculture cells, which were confluent at day 0, were grown for an additional 7 days. Beginning on day 0, media were changed and treatments (Trm and/or BMPs) were applied, under a laminar flow hood, every 3 or 4 days. The standard culture protocol was as follows: days 1-7, MEM, 10% FBS, 50 g/ml ascorbic acid, ± stimulus; days 8-14, BGJb medium, 10% FBS, 5mM -GlyP (as a source of inorganic phosphate to permit mineralization). Endpoint analysis of bone nodule formation and osteocalcin secretion was performed at day 14. The dose of BMP was chosen as 50 ng/ml

5

10

15

20

25

based on pilot experiments in this system that demonstrated a mid-range effect on the dose-response curve for all BMPs studied.

#### **EXAMPLE 2: Antisense Treatment and Cell Culture**

5

10

15

20

25

To explore the potential functional role of LMP-1 during membranous bone formation, we synthesized an antisense oligonucleotide to block LMP-1 mRNA translation and treated secondary osteoblast cultures that were undergoing differentiation initiated by glucocorticoid. Inhibition of RLMP expression was accomplished with a highly specific antisense oligonucleotide (having no significant homologies to known rat sequences) corresponding to a 25 bp sequence spanning the putative translational start site (SEQ ID NO: 42). Control cultures either did not receive oligonucleotide or they received sense oligonucleotide. Experiments were performed in the presence (preincubation) and absence of lipofectamine. Briefly, 22 g of sense or antisense RLMP oligonucleotide was incubated in MEM for 45 minutes at room temperature. Following that incubation, either more MEM or pre-incubated lipofectamine/MEM (7% v/v; incubated 45 minutes at room temperature) was added to achieve an oligonucleotide concentration of 0.2 M. The resulting mixture was incubated for 15 minutes at room temperature. Oligonucleotide mixtures were then mixed with the appropriate medium, that is, MEM/Ascorbate/±Trm, to achieve a final oligonucleotide concentration of 0.1 M.

Cells were incubated with the appropriate medium (±stimulus) in the presence or absence of the appropriate oligonucleotides. Cultures originally incubated with lipofectamine were re-fed after 4 hours of incubation (37°C; 5% CO<sub>2</sub>) with media containing neither lipofectamine nor oligonucleotide. All cultures, especially cultures receiving oligonucleotide, were re-fed every 24 hours to maintain oligonucleotide levels.

30

LMP-1 antisense oligonucleotide inhibited mineralized nodule formation and osteocalcin secretion in a dose-dependent manner, similar to the effect of

BMP-6 oligonucleotide. The LMP-1 antisense block in osteoblast differentiation could not be rescued by addition of exogenous BMP-6, while the BMP-6 antisense oligonucleotide inhibition was reversed with addition of BMP-6. This experiment further confirmed the upstream position of LMP-1 relative to BMP-6 in the osteoblast differentiation pathway. LMP-1 antisense oligonucleotide also inhibited spontaneous osteoblast differentiation in primary rat osteoblast cultures.

#### **EXAMPLE 3: Quantitation of Mineralized Bone Nodule Formation**

10

15

20

25

30

5

Cultures of ROBs prepared according to Examples 1 and 2 were fixed overnight in 70% ethanol and stained with von Kossa silver stain. A semi-automated computerized video image analysis system was used to quantitate nodule count and nodule area in each well. Boden *et al.*, Endocrinology, 137(8):3401-07 (1996). These values were then divided to calculate the area per nodule values. This automated process was validated against a manual counting technique and demonstrated a correlation coefficient of 0.92 (p < 0.000001). All data are expressed as the mean ± standard error of the mean (S.E.M.) calculated from 5 or 6 wells at each condition. Each experiment was confirmed at least twice using cells from different calvarial preparations.

### **EXAMPLE 4: Quantitation of Osteocalcin Secretion**

Osteocalcin levels in the culture media were measured using a competitive radioimmunoassay with a monospecific polyclonal antibody (Pab) raised in our laboratory against the C-terminal nonapeptide of rat osteocalcin as described in Nanes *et al.*, Endocrinology, 127:588 (1990). Briefly, 1 g of nonapeptide was iodinated with 1 mCi <sup>125</sup>I-Na by the lactoperoxidase method. Tubes containing 200 I of assay buffer (0.02 M sodium phosphate, 1 mM EDTA, 0.001% thimerosal, 0.025% BSA) received media taken from cell cultures or osteocalcin standards (0 - 12,000 fmole) at 100 I/tube in assay

buffer. The Pab (1:40,000; 100 I) was then added, followed by the iodinated peptide (12,000 cpm; 100 I). Samples tested for non-specific binding were prepared similarly but contained no antibody.

Bound and free PAbs were separated by the addition of 700 I goat antirabbit IgG, followed by incubation for 18 hours at 4°C. After samples were centrifuged at 1200 rpm for 45 minutes, the supernatants were decanted and the precipitates counted in a gamma counter. Osteocalcin values were reported in fmole/100 I, which was then converted to pmole/ml medium (3-day production) by dividing those values by 100. Values were expressed as the mean ± S.E.M. of triplicate determinations for 5-6 wells for each condition. Each experiment was confirmed at least two times using cells from different calvarial preparations.

#### EXAMPLE 5: Effect of Trm and RLMP on Mineralization In Vitro

15

20

25

30

10

5

There was little apparent effect of either the sense or antisense oligonucleotides on the overall production of bone nodules in the non-stimulated cell culture system. When ROBs were stimulated with Trm, however, the antisense oligonucleotide to RLMP inhibited mineralization of nodules by > 95%. The addition of exogenous BMP-6 to the oligonucleotide-treated cultures did not rescue the mineralization of RLMP-antisense-treated nodules.

Osteocalcin has long been synonymous with bone mineralization, and osteocalcin levels have been correlated with nodule production and mineralization. The RLMP-antisense oligonucleotide significantly decreases osteocalcin production, but the nodule count in antisense-treated cultures does not change significantly. In this case, the addition of exogenous BMP-6 only rescued the production of osteocalcin in RLMP-antisense-treated cultures by 10-15%. This suggests that the action of RLMP is downstream of, and more specific than, BMP-6.

#### **EXAMPLE 6: Harvest and Purification of RNA**

Cellular RNA from duplicate wells of ROBs (prepared according to Examples 1 and 2 in 6-well culture dishes) was harvested using 4M guanidine isothiocyanate (GIT) solution to yield statistical triplicates. Briefly, culture supernatant was aspirated from the wells, which were then overlayed with 0.6 ml of GIT solution per duplicate well harvest. After adding the GIT solution, the plates were swirled for 5-10 seconds (being as consistent as possible). Samples were saved at -70°C for up to 7 days before further processing.

RNA was purified by a slight modification of standard methods according to Sambrook *et al.*, Molecular Cloning: a Laboratory Manual, 2nd Ed., chapter 7.19, Cold Spring Harbor Press (1989). Briefly, thawed samples received 60 12.0 M sodium acetate (pH 4.0), 550 I phenol (water saturated) and 150 I chloroform:isoamyl alcohol (49:1). After vortexing, the samples were centrifuged (10000 x g; 20 minutes; 4°C), the aqueous phase transferred to a fresh tube, 600 I isopropanol was added and the RNA precipitated overnight at -20°C.

Following the overnight incubation, the samples were centrifuged (10000 x g; 20 minutes) and the supernatant was aspirated gently. The pellets were resuspended in 400 TDEPC-treated water, extracted once with phenol:chloroform (1:1), extracted with chloroform:isoamyl alcohol (24:1) and precipitated overnight at -20°C after addition of 40 T sodium acetate (3.0 M; pH 5.2) and 1.0 ml absolute ethanol. To recover the cellular RNA, the samples were centrifuged (10000 x g; 20 min), washed once with 70% ethanol, air dried for 5-10 minutes and resuspended in 20 T of DEPC-treated water. RNA concentrations were calculated from optical densities that were determined with a spectrophotometer.

30

5

10

15

20

### **EXAMPLE 7: Reverse Transcription-Polymerase Chain Reaction**

Heated total RNA (5 g in 10.5 I total volume DEPC-H<sub>2</sub>O at 65°C for 5 minutes) was added to tubes containing 4 I 5X MMLV-RT buffer, 2 I dNTPs, 2 I dT17 primer (10 pmol/ml), 0.5 I RNAsin (40U/ml) and 1 I MMLV-RT (200 units/I). The samples were incubated at 37°C for 1 hour, then at 95°C for 5 minutes to inactivate the MMLV-RT. The samples were diluted by addition of 80 I of water.

Reverse-transcribed samples (5 I) were subjected to polymerase-chain reaction using standard methodologies (50 I total volume). Briefly, samples were added to tubes containing water and appropriate amounts of PCR buffer, 25 mM MgCl<sub>2</sub>, dNTPs, forward and reverse primers for glyceraldehyde 3-phosphate dehydrogenase (GAP, a housekeeping gene) and/or BMP-6), <sup>32</sup>P-dCTP, and Taq polymerase. Unless otherwise noted, primers were standardized to run consistently at 22 cycles (94°C, 30"; 58°C, 30"; 72°C, 20").

# EXAMPLE 8: Quantitation of RT-PCR Products by Polyacrylamide Gel Electrophoresis (PAGE) and PhosphorImager Analysis

20

25

5

10

15

RT-PCR products received 5 I/tube loading dye, were mixed, heated at 65°C for 10 min and centrifuged. Ten I of each reaction was subjected to PAGE (12% polyacrylamide:bis; 15 V/well; constant current) under standard conditions. Gels were then incubated in gel preserving buffer (10% v/v glycerol, 7% v/v acetic acid, 40% v/v methanol, 43% deionized water) for 30 minutes, dried (80°C) *in vacuo* for 1-2 hours and developed with an electronically-enhanced phosphoresence imaging system for 6-24 hours. Visualized bands were analyzed. Counts per band were plotted graphically.

### **EXAMPLE 9: Differential Display PCR**

RNA was extracted from cells stimulated with glucocorticoid (Trm, 1 nM). Heated, DNase-treated total RNA (5 g in 10.5 I total volume in DEPC-H<sub>2</sub>O at 65°C for 5 minutes) was reverse transcribed as described in Example 7, but H-T<sub>11</sub>M (SEQ ID. NO: 4) was used as the MMLV-RT primer. The resulting cDNAs were PCR-amplified as described above, but with various commercial primer sets (for example, H-T<sub>11</sub>G (SEQ ID NO: 4) and H-AP-10 (SEQ ID. NO: 5); GenHunter Corp, Nashville, TN). Radiolabelled PCR products were fractionated by gel electrophoresis on a DNA sequencing gel. After electrophoresis, the resulting gels were dried *in vacuo* and autoradiographs were exposed overnight. Bands representing differentially-expressed cDNAs were excised from the gel and reamplified by PCR using the method of Conner *et al.*, Proc. Natl. Acad. Sci. USA, 88:278 (1983). The products of PCR reamplification were cloned into

#### EXAMPLE 10: Screening of a UMR 106 Rat Osteosarcoma Cell cDNA Library

the vector PCR-II (TA cloning kit; InVitrogen, Carlsbad, CA).

A UMR 106 library (2.5 x 10<sup>10</sup> pfu/ml) was plated at 5 x 10<sup>4</sup> pfu/ml onto agar plates (LB bottom agar) and the plates were incubated overnight at 37°C. Filter membranes were overlaid onto plates for two minutes. Once removed, the filters were denatured, rinsed, dried and UV cross-linked. The filters were then incubated in pre-hyridization buffer (2X PIPES [pH 6.5], 5% formamide, 1% SDS and 100 g/ml denatured salmon sperm DNA) for 2 h at 42°C. A 260 base-pair radiolabelled probe (SEQ ID NO: 3; <sup>32</sup>P labelled by random priming) was added to the entire hybridization mix/filters, followed by hybridization for 18 hours at 42°C. The membranes were washed once at room temperature (10 min, 1 x SSC, 0.1% SDS) and three times at 55°C (15 min, 0.1 x SSC, 0.1% SDS).

10

15

20

25

After they were washed, the membranes were analyzed by autoradiography as described above. Positive clones were plaque purified. The procedure was repeated with a second filter for four minutes to minimize spurious positives. Plaque-purified clones were rescued as lambda SK(-) phagemids. Cloned cDNAs were sequenced as described below.

#### **EXAMPLE 11: Sequencing of Clones**

5

10

15

20

25

30

Cloned cDNA inserts were sequenced by standard methods. Ausubel *et al.*, <u>Current Protocols in Molecular Biology</u>, Wiley Interscience (1988). Briefly, appropriate concentrations of termination mixture, template and reaction mixture were subjected to an appropriate cycling protocol (95°C,30s; 68°C,30s; 72°C,60s; x 25). Stop mixture was added to terminate the sequencing reactions. After heating at 92°C for 3 minutes, the samples were loaded onto a denaturing 6% polyacrylamide sequencing gel (29:1 acrylamide:bisacrylamide). Samples were electrophoresed for about 4 hours at 60 volts, constant current. After electrophoresis, the gels were dried *in vacuo* and autoradiographed.

The autoradiographs were analyzed manually. The resulting sequences were screened against the databases maintained by the National Center for Biotechnology Information (NIH, Bethesda, MD; http://www.ncbi.nlm.nih.gov/) using the BLASTn program set with default parameters. Based on the sequence data, new sequencing primers were prepared and the process was repeated until the entire gene had been sequenced. All sequences were confirmed a minimum of three times in both orientations.

Nucleotide and amino acid sequences were also analyzed using the PCGENE software package (version 16.0). Per cent homology values for nucleotide sequences were calculated by the program NALIGN, using the following parameters: weight of non-matching nucleotides, 10; weight of non-matching gaps, 10; maximum number of nucleotides considered, 50; and minimum number of nucleotides considered, 50.

For amino acid sequences, per cent homology values were calculated using PALIGN. A value of 10 was selected for both the open gap cost and the unit gap cost.

#### 5 EXAMPLE 12: Cloning of RLMP cDNA

The differential display PCR amplification products described in Example 9 contained a major band of approximately 260 base pairs. This sequence was used to screen a rat osteosarcoma (UMR 106) cDNA library. Positive clones were subjected to nested primer analysis to obtain the primer sequences necessary for amplifying the full length cDNA. (SEQ. ID NOs: 11, 12, 29, 30 and 31) One of those positive clones selected for further study was designated clone 10-4.

Sequence analysis of the full-length cDNA in clone 10-4, determined by nested primer analysis, showed that clone 10-4 contained the original 260 base-pair fragment identified by differential display PCR. Clone 10-4 (1696 base pairs; SEQ ID NO: 2) contains an open reading frame of 1371 base pairs encoding a protein having 457 amino acids (SEQ ID NO: 1). The termination codon, TGA, occurs at nucleotides 1444-1446. The polyadenylation signal at nucleotides 1675-1680, and adjacent poly(A)\* tail, was present in the 3' noncoding region. There were two potential N-glycosylation sites, Asn-Lys-Thr and Asn-Arg-Thr, at amino acid positions 113-116 and 257-259 in SEQ ID NO: 1, respectively. Two potential cAMP- and cGMP-dependent protein kinase phosphorylation sites, Ser and Thr, were found at amino acid positions 191 and 349, respectively. There were five potential protein kinase C phosphorylation sites, Ser or Thr, at amino acid positions 3, 115, 166, 219, 442. One potential ATP/GTP binding site motif A (P-loop), Gly-Gly-Ser-Asn-Asn-Gly-Lys-Thr, was determined at amino acid positions 272-279.

In addition, two highly conserved putative LIM domains were found at amino acid positions 341-391 and 400-451. The putative LIM domains in this newly identified rat cDNA clone showed considerable homology with the LIM

10

15

20

25

domains of other known LIM proteins. However, the overall homology with other rat LIM proteins was less than 25%. RLMP (also designated 10-4) has 78.5% amino acid homology to the human enigma protein (see U.S. Patent No. 5,504,192), but only 24.5% and 22.7% amino acid homology to its closest rat homologs, CLP-36 and RIT-18, respectively.

#### **EXAMPLE 13: Northern Blot Analysis of RLMP Expression**

Thirty g of total RNA from ROBs, prepared according to Examples 1 and 2, was size fractionated by formaldehyde gel electrophoresis in 1% agarose flatbed gels and osmotically transblotted to nylon membranes. The blot was probed with a 600 base pair EcoR1 fragment of full-length 10-4 cDNA labeled with <sup>32</sup>P-dCTP by random priming.

Northern blot analysis showed a 1.7 kb mRNA species that hybridized with the RLMP probe. RLMP mRNA was up-regulated approximately 3.7-fold in ROBs after 24 hours exposure to BMP-6. No up-regulation of RMLP expression was seen in BMP-2 or BMP-4-stimulated ROBs at 24 hours.

## **EXAMPLE 14: Statistical Methods**

20

25

30

5

10

15

For each reported nodule/osteocalcin result, data from 5-6 wells from a representative experiment were used to calculate the mean ± S.E.M. Graphs may be shown with data normalized to the maximum value for each parameter to allow simultaneous graphing of nodule counts, mineralized areas and osteocalcin.

For each reported RT-PCR, RNase protection assay or Western blot analysis, data from triplicate samples of representative experiments, were used to determine the mean ± S.E.M. Graphs may be shown normalized to either day 0 or negative controls and expressed as fold-increase above control values.

Statistical significance was evaluated using a one-way analysis of variance with post-hoc multiple comparison corrections of Bonferroni as appropriate. D. V. Huntsberger, "The Analysis of Variance," in Elements of Statistical Variance, P. Billingsley (ed.), pp. 298-330, Allyn & Bacon Inc., Boston, MA (1977) and Sigmastat, Jandel Scientific, Corte Madera, CA. Alpha levels for significance were defined as p < 0.05.

EXAMPLE 15: Detection of Rat LIM Mineralization Protein by Western Blot Analysis

10

15

25

30

5

Polyclonal antibodies were prepared according to the methods of England *et al.*, <u>Biochim.Biophys. Acta</u>, 623:171 (1980) and Timmer *et al.*, <u>J. Biol. Chem.</u>, 268:24863 (1993).

HeLa cells were transfected with pCMV2/RLMP. Protein was harvested from the transfected cells according to the method of Hair *et al.*, <u>Leukemia</u> Research, 20:1 (1996). Western Blot Analysis of native RLMP was performed as described by Towbin *et al.*, <u>Proc. Natl. Acad. Sci. USA</u>, 76:4350 (1979).

EXAMPLE 16: Synthesis of the Rat LMP-Unique (RLMPU) derived Human

20 PCR product

Based on the sequence of the rat LMP-1 cDNA, forward and reverse PCR primers (SEQ ID NOs: 15 and 16) were synthesized and a unique 223 base-pair sequence was PCR amplified from the rat LMP-1 cDNA. A similar PCR product was isolated from human MG63 osteosarcoma cell cDNA with the same PCR primers.

RNA was harvested from MG63 osteosarcoma cells grown in T-75 flasks. Culture supernatant was removed by aspiration and the flasks were overlayed with 3.0 ml of GIT solution per duplicate, swirled for 5-10 seconds, and the resulting solution was transferred to 1.5 ml eppendorf tubes (5 tubes with 0.6 ml/tube). RNA was purified by a slight modification of standard

methods, for example, see Sambrook et al., Molecular Cloning: A Laboratory Manual, chapter 7, page 19, Cold Spring Harbor Laboratory Press (1989) and Boden et al., Endocrinology, 138:2820-28 (1997). Briefly, the 0.6 ml samples received 60 12.0 M sodium acetate (pH 4.0), 550 I water saturated phenol and 150 I chloroform:isoamyl alcohol (49:1). After addiiton of those reagents, the samples were vortexed, centrifuged (10000 x. g; 20 min; 4C) and the aqueous phase transferred to a fresh tube. Isopropanol (600 I) was added and the RNA was precipitated overnight at -20°C. The samples were centrifuged (10000 x g; 20 minutes) and the supernatant was aspirated gently. The pellets were resuspended in 400 I of DEPC-treated water, extracted once with phenol:chloroform (1:1), extracted with chloroform; isoamyl alcohol (24:1) and precipitated overnight at -20°C in 40 I sodium acetate (3.0 M; pH 5.2) and 1.0 ml absolute ethanol. After precipitation, the samples were centrifuged (10000 x g: 20 min), washed once with 70% ethanol, air dried for 5-10 minutes and resuspended in 20 I of DEPC-treated water. RNA concentrations were derived from optical densities.

Total RNA (5 g in 10.5 L total volume in DEPC-H<sub>2</sub>O) was heated at 65°C for 5 minutes, and then added to tubes containing 4 I 5X MMLV-RT buffer, 2 I dNTPs, 2 I dT17 primer (10 pmol/ml), 0.5 I RNAsin (40 U/ml) and 1 I MMLV-RT (200 units/I). The reactions were incubated at 37°C for 1 hour. Afterward, the MMLV-RT was inactivated by heating at 95°C for 5 minutes. The samples were diluted by addition of 80 L water.

Transcribed samples (5 I) were subjected to polymerase-chain reaction using standard methodologies (50 I total volume). Boden *et al.*, Endocrinology, 138:2820-28 (1997); Ausubel *et al.*, "Quantitation of rare DNAs by the polymerase chain reaction", *in* Current Protocols in Molecular Biology, chapter 15.31-1, Wiley & Sons, Trenton, NJ (1990). Briefly, samples were added to tubes containing water and appropriate amounts of PCR buffer (25 mM MgC1<sub>2</sub>, dNTPs, forward and reverse primers (for RLMPU; SEQ ID NOs: 15 and 16), <sup>32</sup>P-dCTP, and DNA polymerase. Primers were designed to run consistently at

10

15

20

25

22 cycles for radioactive band detection and 33 cycles for amplification of PCR product for use as a screening probe (94°C, 30 sec, 58°C, 30 sec; 72°C, 20 sec).

Sequencing of the agarose gel-purified MG63 osteosarcoma-derived PCR product gave a sequence more than 95% homologous to the RLMPU PCR product. That sequence is designated HLMP unique region (HLMPU; SEQ ID NO: 6).

## EXAMPLE 17: Screening of reverse-transcriptase-derived MG63 cDNA

10

Screening was performed with PCR using specific primers (SEQ ID NOs: 16 and 17) as described in Example 7. A 717 base-pair MG63 PCR product was agarose gel purified and sequenced with the given primers (SEQ. ID NOs: 12, 15, 16, 17, 18, 27 and 28). Sequences were confirmed a minimum of two times in both directions. The MG63 sequences were aligned against each other and then against the full-length rat LMP cDNA sequence to obtain a partial human LMP cDNA sequence (SEQ ID NO: 7).

### EXAMPLE 18: Screening of a Human Heart cDNA Library

20

25

30

15

Based on Northern blot experiments, it was determined that LMP-1 is expressed at different levels by several different tissues, including human heart muscle. A human heart cDNA library was therefore examined. The library was plated at 5 x 10<sup>4</sup> pfu/ml onto agar plates (LB bottom agar) and plates were grown overnight at 37° C. Filter membranes were overlaid onto the plates for two minutes. Afterward, the filters denatured, rinsed, dried, UV cross-linked and incubated in pre-hyridization buffer (2X PIPES [pH 6.5]; 5% formamide, 1% SDS, 100 g/ml denatured salmon sperm DNA) for 2 h at 42°C. A radiolabelled, LMP-unique, 223 base-pair probe (<sup>32</sup>P, random primer labelling; SEQ ID NO: 6) was added and hybridized for 18 h at 42°C. Following hybridization, the membranes were washed once at room temperature (10 min,

1 x SSC, 0.1% SDS) and three times at 55°C (15 min, 0.1 x SSC, 0.1% SDS). Double-positive plaque-purified heart library clones, identified by autoradiography, were rescued as lambda phagemids according to the manufacturers' protocols (Stratagene, La Jolla, CA).

Restriction digests of positive clones yielded cDNA inserts of varying sizes. Inserts greater than 600 base-pairs in length were selected for initial screening by sequencing. Those inserts were sequenced by standard methods as described in Example 11.

One clone, number 7, was also subjected to automated sequence analysis using primers corresponding to SEQ ID NOs: 11-14, 16 and 27. The sequences obtained by these methods were routinely 97-100% homologous. Clone 7 (Partial Human LMP-1 cDNA from a heart library; SEQ. ID NO: 8) contained sequence that was more than 87% homologous to the rat LMP cDNA sequence in the translated region.

15

20

25

30

10

5

### EXAMPLE 19: Determination of Full-Length Human LMP-1 cDNA

Overlapping regions of the MG63 human osteosarcoma cell cDNA sequence and the human heart cDNA clone 7 sequence were used to align those two sequences and derive a complete human cDNA sequence of 1644 base-pairs. NALIGN, a program in the PCGENE software package, was used to align the two sequences. The overlapping regions of the two sequences constituted approximately 360 base-pairs having complete homology except for a single nucleotide substitution at nucleotide 672 in the MG63 cDNA (SEQ ID NO: 7) with clone 7 having an "A" instead of a "G" at the corresponding nucleotide 516 (SEQ ID NO: 8).

The two aligned sequences were joined using SEQIN, another subprogram of PCGENE, using the "G" substitution of the MG63 osteosarcoma cDNA clone. The resulting sequence is shown in SEQ ID NO: 9. Alignment of the novel human-derived sequence with the rat LMP-1 cDNA was accomplished with NALIGN. The full-length human LMP-1 cDNA sequence

(SEQ. ID NO: 9) is 87.3% homologous to the translated portion of rat LMP-1 cDNA sequence.

#### EXAMPLE 20: Determination of Amino Acid Sequence of Human LMP-1

5

10

15

25

30

The putative amino acid sequence of human LMP-1 was determined with the PCGENE subprogram TRANSL. The open reading frame in SEQ ID NO: 9 encodes a protein comprising 457 amino acids (SEQ. ID NO: 10). Using the PCGENE subprogram Palign, the human LMP-1 amino acid sequence was found to be 94.1% homologous to the rat LMP-1 amino acid sequence.

# EXAMPLE 21: Determination of the 5 Prime Untranslated Region of the Human LMP cDNA

MG63 5' cDNA was amplified by nested RT-PCR of MG63 total RNA using a 5' rapid amplification of cDNA ends (5' RACE) protocol. This method included first strand cDNA synthesis using a lock-docking oligo (dT) primer with two degenerate nucleotide positions at the 3' end (Chenchik *et al.*, CLONTECHniques, X:5 (1995); Borson *et al.*, PC Methods Applic., 2:144 (1993)). Second-strand synthesis is performed according to the method of Gubler *et al.*, Gene, 25:263 (1983), with a cocktail of *Escherichia coli* DNA polymerase I, RNase H, and *E. coli* DNA ligase. After creation of blunt ends with T4 DNA polymerase, double-stranded cDNA was ligated to the fragment (5' -CTAATACGACTCACTATAGGGCTCGAGCGGCCGCCCGGGCAGGT- 3') (SEQ.ID NO: 19). Prior to RACE, the adaptor-ligated cDNA was diluted to a concentration suitable for Marathon RACE reactions (1:50). Adaptor-ligated double-stranded cDNA was then ready to be specifically cloned.

First-round PCR was performed with the adaptor-specific oligonucleotide, 5'-CCATCCTAATACGACTCACTATAGGGC- 3' (AP1) (SEQ.ID NO: 20) as sense primer and a Gene Specific Primer (GSP) from the unique region described in Example 16 (HLMPU). The second round of PCR was

performed using a nested primers GSP1-HLMPU (antisense/reverse primer) (SEQ. ID NO: 23) and GSP2-HLMPUF (SEQ. ID NO: 24) (see Example 16; sense/forward primer). PCR was performed using a commercial kit (Advantage cDNA PCR core kit; CloneTech Laboratories Inc., Palo Alto, CA) that utilizes an antibody-mediated, but otherwise standard, hot-start protocol. PCR conditions for MG63 cDNA included an initial hot-start denaturation (94°C, 60 sec) followed by: 94°C, 30 sec; 60°C, 30 sec; 68°C, 4 min; 30 cycles. The first-round PCR product was approximately 750 base-pairs in length whereas the nested PCR product was approximately 230 base-pairs. The first-round PCR product was cloned into linearized pCR 2.1 vector (3.9 Kb). The inserts were sequenced in both directions using M13 Forward and Reverse primers (SEQ. ID NO: 11; SEQ. ID NO: 12)

# EXAMPLE 22: Determination of Full-length Human LMP-1 cDNA with 5 Prime UTR

Overlapping MG63 human osteosarcoma cell cDNA 5'-UTR sequence (SEQ ID NO: 21), MG63 717 base-pair sequence (Example 17; SEQ ID NO: 8) and human heart cDNA clone 7 sequence (Example 18) were aligned to derive a novel human cDNA sequence of 1704 base-pairs (SEQ.ID NO: 22). The alignment was accomplished with NALIGN, (both PCGENE and Omiga 1.0; Intelligenetics). Over-lapping sequences constituted nearly the entire 717 base-pair region (Example 17) with 100% homology. Joining of the aligned sequences was accomplished with SEQIN.

25

30

10

15

20

#### **EXAMPLE 23: Construction of LIM Protein Expression Vector**

The construction of pHIS-5ATG LMP-1s expression vector was carried out with the sequences described in Examples 17 and 18. The 717 base-pair clone (Example 17; SEQ ID NO: 7) was digested with <u>Clal</u> and <u>EcoRV</u>. A small fragment (~250 base-pairs) was gel purified. Clone 7 (Example 18; SEQ

ID NO: 8) was digested with <u>Clal</u> and <u>Xbal</u> and a 1400 base-pair fragment was gel purified. The isolated 250 base-pair and 1400 base-pair restriction fragments were ligated to form a fragment of ~1650 base-pairs.

Due to the single nucleotide substitution in Clone 7 (relative to the 717 base-pair PCR sequence and the original rat sequence) a stop codon at translated base-pair 672 resulted. Because of this stop codon, a truncated (short) protein was encoded, hence the name LMP-1s. This was the construct used in the expression vector (SEQ ID NO: 32). The full length cDNA sequence with 5' UTR (SEQ ID NO: 33) was created by alignment of SEQ ID NO: 32 with the 5' RACE sequence (SEQ ID NO: 21). The amino acid sequence of LMP-1s (SEQ ID NO: 34) was then deduced as a 223 amino acid protein and confirmed by Western blot (as in Example 15) to run at the predicted molecular weight of ~-23.7 kD.

The pHis-ATG vector (InVitrogen, Carlsbad, CA) was digested with EcoRV and Xbal. The vector was recovered and the 1650 base-pair restriction fragment was then ligated into the linearized pHis-ATG. The ligated product was cloned and amplified. The pHis-ATG-LMP-1s Expression vector, also designated pHIS-A with insert HLMP-1s, was purified by standard methods.

EXAMPLE 24: Induction of Bone Nodule Formation and Mineralization *In vitro*with LMP Expression Vector

Rat Calvarial cells were isolated and grown in secondary culture according to Example 1. Cultures were either unstimulated or stimulated with glucocorticoid (GC) as described in Example 1. A modification of the Superfect Reagent (Qiagen, Valencia, CA) transfection protocol was used to transfect 3 g/well of each vector into secondary rat calvarial osteoblast cultures according to Example 25.

Mineralized nodules were visualized by Von Kossa staining, as described in Example 3. Human LMP-1s gene product overexpression alone induced bone nodule formation (~203 nodules/well) *in vitro*. Levels of nodules were

5

10

15

20

25

approximately 50% of those induced by the GC positive control (~412 nodules/well). Other positive controls included the pHisA-LMP-Rat expression vector (~152 nodules/well) and the pCMV2/LMP-Rat-Fwd Expression vector (~206 nodules/well), whereas the negative controls included the pCMV2/LMP-Rat-Rev. Expression vector (~2 nodules/well) and untreated (NT) plates (~4 nodules/well). These data demonstrate that the human cDNA was at least as osteoinductive as the rat cDNA. The effect was less than that observed with GC stimulation, most likely due to suboptimal

10

15

20

25

30

doses of Expression vector.

# EXAMPLE 25: LMP-Induced Cell Differentiation In Vitro and In Vivo

The rat LMP cDNA in clone 10-4 (see Example 12) was excised from the vector by double-digesting the clone with Notl and Apal overnight at 37°C. Vector pCMV2 MCS (InVitrogen, Carlsbad, CA) was digested with the same restriction enzymes. Both the linear cDNA fragment from clone 10-4 and pCMV2 were gel purified, extracted and ligated with T4 ligase. The ligated DNA was gel purified, extracted and used to transform *E. coli* JM109 cells for amplification. Positive agar colonies were picked, digested with Notl and Apal and the restriction digests were examined by gel electrophoresis. Stock cultures were prepared of positive clones.

A reverse vector was prepared in analogous fashion except that the restriction enzymes used were Xbal and HindIII. Because these restriction enzymes were used, the LMP cDNA fragment from clone 10-4 was inserted into pRc/CMV2 in the reverse (that is, non-translatable) orientation. The recombinant vector produced is designated pCMV2/RLMP.

An appropriate volume of pCMV10-4 (60 nM final concentration is optimal [3 g]; for this experiment a range of 0-600 nM/well [0-30 g/well] final concentration is preferred) was resuspended in Minimal Eagle Media (MEM) to 450 I final volume and vortexed for 10 seconds. Superfect was added (7.5 l/ml final solution), the solution was vortexed for 10 seconds and then incubated at

room termperature for 10 minutes. Following this incubation, MEM supplemented with 10% FBS (1 ml/well; 6 ml/plate) was added and mixed by pipetting.

The resulting solution was then promptly pipetted (1 ml/well) onto washed ROB cultures. The cultures were incubated for 2 hours at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub>. Afterward, the cells were gently washed once with sterile PBS and the appropriate normal incubation medium was added.

Results demonstrated significant bone nodule formation in all rat cell cultures which were induced with pCMV10-4. For example, pCMV10-4 transfected cells produced 429 nodules/well. Positive control cultures, which were exposed to Trm, produced 460 nodules/well. In contrast, negative controls, which received no treatment, produced 1 nodule/well. Similarly, when cultures were transfected with pCMV10-4 (reverse), no nodules were observed.

For demonstrating *de novo* bone formation *in vivo*, marrow was aspirated from the hindlimbs of 4-5 week old normal rats (rnu/+; heterozygous for recessive athymic condition). The aspirated marrow cells were washed in alpha MEM, centrifuged, and RBCs were lysed by resuspending the pellet in 0.83% NH<sub>4</sub>Cl in 10 mM Tris (pH 7.4). The remaining marrow cells were washed 3x with MEM and transfected for 2 hours with 9 g of pCMV-LMP-1s (forward or reverse orientation) per 3 x 10<sup>6</sup> cells. The transfected cells were then washed 2X with MEM and resuspended at a concentration of 3 x 10<sup>7</sup> cells/ml.

The cell suspension (100 I) was applied via sterile pipette to a sterile 2 x 5 mm type I bovine collagen disc (Sulzer Orthopaedics, Wheat Ridge, CO). The discs were surgically implanted subcutaneously on the skull, chest, abdomen or dorsal spine of 4-5 week old athymic rats (rnu/rnu). The animals were scarified at 3-4 weeks, at which time the discs or surgical areas were excised and fixed in 70% ethanol. The fixed specimens were analyzed by radiography and undecalcified histologic examination was performed on 5 m

5

10

15

20

25

thick sections stained with Goldner Trichrome. Experiments were also performed using devitalized (guanidine extracted) demineralized bone matrix (Osteotech, Shrewsbury, NJ) in place of collagen discs.

Radiography revealed a high level of mineralized bone formation that conformed to the form of the original collagen disc containing LMP-1s transfected marrow cells. No mineralized bone formation was observed in the negative control (cells transfected with a reverse-oriented version of the LMP-1s cDNA that did not code for a translated protein), and absorption of the carrier appeared to be well underway.

Histology revealed new bone trabeculae lined with osteroblasts in the LMP-1s transfected implants. No bone was seen along with partial resorption of the carrier in the negative controls.

Radiography of a further experiment in which 18 sets (9 negative control pCMV-LMP-REV & 9 experimental pCMV-LMP-1s) of implants were added to sites alternating between lumbar and thoracic spine in athymic rats demonstrated 0/9 negative control implants exhibiting bone formation (spine fusion) between vertebrae. All nine of the pCMV-LMP-1s treated implants exhibited solid bone fusions between vertebrae.

20 EXAMPLE 26: The Synthesis of pHIS-5' ATG LMP-1s Expression Vector from the sequences Demonstrated in Examples 2 and 3.

The 717 base-pair clone (Example 17) was digested with <u>Clal</u> and <u>Eco</u>RV (New England Biologicals, city, MA). A small fragment (~250 base-pairs) was gel purified. Clone No. 7 (Example 18) was digested with <u>Clal</u> and <u>Xbal</u>. A 1400 base-pair fragment was gel purified from that digest. The isolated 250 base-pair and 1400 base-pair cDNA fragments were ligated by standard methods to form a fragment of ~1650 bp. The pHis-A vector (InVitrogen) was digested with <u>Eco</u>RV and *Xbal*. The linearized vector was recovered and ligated to the chimeric 1650 base-pair cDNA fragment. The ligated product was cloned and amplified by standard methods, and the pHis-

5

10

15

25

A-5' ATG LMP-1s expression vector, also denominated as the vector pHis-A with insert HLMP-1s, was deposited at the ATCC as previously described.

EXAMPLE 27: The Induction of Bone Nodule Formation and Mineralization In

Vitro With pHis-5' ATG LMP-1s Expression Vector

Rat calvarial cells were isolated and grown in secondary culture according to Example 1. Cultures were either unstimulated or stimulated with glucocorticoid (GC) according to Example 1. The cultures were transfected with 3 g of recombinant pHis-A vector DNA/well as described in Example 25. Mineralized nodules were visualized by Von Kossa staining according to Example 3.

Human LMP-1s gene product overexpression alone (*i.e.*, without GC stimulation) induced significant bone nodule formation (~203 nodules/well) *in vitro*. This is approximately 50% of the amount of nodules produced by cells exposed to the GC positive control (~412 nodules/well). Similar results were obtained with cultures transfected with pHisA-LMP-Rat Expression vector (~152 nodules/well) and pCMV2/LMP-Rat-Fwd (~206 nodules/well). In contrast, the negative control pCMV2/LMP-Rat-Rev yielded (~2 nodules/well), while approximately 4 nodules/well were seen in the untreated plates. These data demonstrate that the human LMP-1 cDNA was at least as osteoinductive as the rat LMP-1 cDNA in this model system. The effect in this experiment was less than that observed with GC stimulation; but in some the effect was comparable.

25

30

5

10

15

20

# EXAMPLE 28: LMP Induces Secretion of a Soluble Osteoinductive Factor

Overexpression of RLMP-1 or HLMP-1s in rat calvarial osteoblast cultures as described in Example 24 resulted in significantly greater nodule formation than was observed in the negative control. To study the mechanism of action of LIM mineralization protein conditioned medium was harvested at

different time points, concentrated to 10 X, sterile filtered, diluted to its original concentration in medium containing fresh serum, and applied for four days to untransfected cells.

Conditioned media harvested from cells transfected with RLMP-1 or HLMP-1s at day 4 was approximately as effective in inducing nodule formation as direct overexpression of RLMP-1 in transfected cells. Conditioned media from cells transfected with RLMP-1 or HLMP-1 in the reverse orientation had no apparent effect on nodule formation. Nor did conditioned media harvested from LMP-1 transfected cultures before day 4 induce nodule formation. These data suggest that expression of LMP-1 caused the synthesis and/or secretion of a soluble factor, which did not appear in culture medium in effectie amounts until 4 days post transfection.

Since overexpression of rLMP-1 resulted in the secretion of an osteoinductive factor into the medium, Western blot analysis was used to determine if LMP-1 protein was present in the medium. The presence of rLMP-1 protein was assessed using antibody specific for LMP-1 (QDPDEE) and detected by conventional means. LMP-1 protein was found only in the cell layer of the culture and not detected in the medium.

Partial purification of the osteoinductive soluble factor was accomplished by standard 25% and 100% ammonium sulfate cuts followed by DE-52 anion exchange batch chromatography (100 mM or 500 mM NaCl). All activity was observed in the high ammonium sulfate, high NaCl fractions. Such localization is consistent with the possibility of a single factor being responsible for conditioning the medium.

25

30

5

10

15

20

EXAMPLE 29: Gene Therapy In Lumbar Spine Fusion Mediated by Low Dose Adenovirus

This study determined the optimal dose of adenoviral delivery of the LMP-1 cDNA (SEQ ID NO: 2) to promote spine fusion in normal, that is, immune competent, rabbits.

A replication-deficient human recombinant adenovirus was constructed with the LMP-1 cDNA (SEQ ID NO: 2) driven by a CMV promoter using the Adeno-Quest™ Kit (Quantum Biotechnologies, Inc., Montreal). A commercially available (Quantum Biotechnologies, Inc., Montreal) recombinant adenovirus containing the beta-galactosidase gene was used as a control.

Initially, an *in vitro* dose response experiment was performed to determine the optimal concentration of adenovirus-delivered LMP-1 ("AdV-LMP-1") to induce bone differentiation in rat calvarial osteoblast cultures using a 60-minute transduction with a multiplicity of infection ("MOI") of 0.025, 0.25, 2.5, or 25 plaque-forming units (pfu) of virus per cell. Positive control cultures were differentiated by a 7-day exposure to 10° M glucocorticoid ("GC"). Negative control cultures were left untreated. On day 14, the number of mineralized bone nodules was counted after von Kossa staining of the cultures, and the level of osteocalcin secreted into the medium (pmol/mL) was measured by radioimmunoassay (mean ± SEM).

The results of this experiment are shown in Table I. Essentially no spontaneous nodules formed in the untreated negative control cultures. The data show that a MOI equal to 0.25 pfu/cell is most effective for osteoinducing bone nodules, achieving a level comparable to the positive control (GC). Lower and higher doses of adenovirus were less effective.

TABLE I

Outcome			AdV-LMP-1 Dose (MOI)			
	Neg. Ctrl.	GC	0.025	0.25	2.5	25
Bone Nodules	0.5±0.2	188±35	79.8±13	145.1±1 3	26.4±15	87.6±2
Osteocalcin	1.0±0.1	57.8±9	28.6±11	22.8±1	18.3±3	26.0±2

5

10

15

In vivo experiments were then performed to determine if the optimal in vitro dose was capable of promoting intertransverse process spine fusions in skeletally mature New Zealand white rabbits. Nine rabbits were anesthetized and 3 cc of bone marrow was aspirated from the distal femur through the intercondylar notch using an 18 gauge needle. The buffy coat was then isolated, a 10-minute transduction with AdV-LMP-1 was performed, and the cells were returned to the operating room for implantation. Single level posterolateral lumbar spine arthrodesis was performed with decortication of transverse processes and insertion of carrier (either rabbit devitalized bone matrix or a collagen sponge) containing 8-15 million autologous nucleated buffy coat cells transduced with either AdV-LMP-1 (MOI = 0.4) or AdV-BGal (MOI = 0.4). Rabbits were euthanized after 5 weeks and spine fusions were assessed by manual palpation, plain x-rays, CT scans, and undecalcified histology.

The spine fusion sites that received AdV-LMP-1 induced solid, continuous spine fusion masses in all nine rabbits. In contrast, the sites receiving AdV-BGal, or a lower dose of AdV-LMP-1 (MOI = 0.04) made little or no bone and resulted in spine fusion at a rate comparable to the carrier alone (< 40%). These results were consistent as evaluated by manual palpation, CT scan, and histology. Plain radiographs, however, sometimes overestimated the amount of bone that was present, especially in the control sites. LMP-1 cDNA delivery and bone induction was successful with both of the carrier materials tested. There was no evidence of systemic or local immune response to the adenovirus vector.

These data demonstrate consistent bone induction in a previously validated rabbit spine fusion model which is quite challenging. Furthermore, the protocol of using autogenous bone marrow cells with intraoperative *ex vivo* gene transduction (10 minutes) is a more clinically feasible procedure than other methods that call for overnight transduction or cell expansion for weeks in culture. In addition, the most effective dose of recombinant adenovirus (MOI=0.25) was substantially lower than doses reported in other gene therapy

5

10

15

20

25

applications (MOI-40-500). We believe this is due to the fact that LMP-1 is an intracellular signaling molecule and may have powerful signal amplification cascades. Moreover, the observation that the same concentration of AdV-LMP-1 that induced bone in cell culture was effective *in vivo* was also surprising given the usual required increase in dose of other growth factors when translating from cell culture to animal experiments. Taken together, these observations indicate that local gene therapy using adenovirus to deliver the LMP-1 cDNA is possible and the low dose required will likely minimize the negative effects of immune response to the adenovirus vector.

10

15

20

25

30

5

# EXAMPLE 30: Use of Peripheral Venous Blood Nucleated Cells (Buffy Coat) for Gene Therapy With LMP-1 cDNA To Make Bone

In four rabbits we performed spine fusion surgery as above (Example 29) except the transduced cells were the buffy coat from venous blood rather than bone marrow. These cells were transfected with Adeno-LMP or pHIS-LMP plasmid and had equivalent successful results as when bone marrow cells were used. This discovery of using ordinary venous blood cells for gene delivery makes gene therapy more feasible clinically since it avoids painful marrow harvest under general anesthesia and yields two times more cells per mL of starting material.

### **EXAMPLE 31: Isolation of Human LMP-1 Splice Variants**

Intron/Exon mRNA transcript splice variants are a relatively common regulatory mechanism in signal transduction and cellular/tissue development. Splice variants of various genes have been shown to alter protein-protein, protein-DNA, protein-RNA, and protein-substrate interactions. Splice variants may also control tissue specificity for gene expression allowing different forms (and therefore functions) to be expressed in various tissues. Splice variants are a common regulatory phenomenon in cells. It is possible that the LMP

splice variants may result in effects in other tissues such as nerve regeneration, muscle regeneration, or development of other tissues.

To screen a human heart cDNA library for splice variants of the HLMP-1 sequence, a pair of PCR primer corresponding to sections of SEQ ID NO: 22 was prepared. The forward PCR primer, which was synthesized using standard techniques, corresponds to nucleotides 35-54 of SEQ ID NO: 22. It has the following sequence:

5' GAGCCGGCATCATGGATTCC 3' (SEQ ID NO: 35)

10

5

The reverse PCR primer, which is the reverse complement of nucleotides 820-839 in SEQ ID NO: 22, has the following sequence:

5' GCTGCCTGCACAATGGAGGT 3' (SEQ ID NO: 36)

15

20

25

The forward and reverse PCR primers were used to screen human heart cDNA (ClonTech, Cat No. 7404-1) for sequences similar to HLMP-1 by standard techniques, using a cycling protocol of 94°C for 30 seconds, 64°C for 30 seconds, and 72°C for 1 minute, repeated 30 times and followed by a 10 minute incubation at 72°C. The amplification cDNA sequences were gelpurified and submitted to the Emory DNA Sequence Core Facility for sequencing. The clones were sequenced using standard techniques and the sequences were examined with PCGENE (Intelligenetics; Programs SEQUIN and NALIGN) to determine homology to SEQ ID NO: 22. Two homologous nucleotide sequences with putative alternative splice sites compared to SEQ ID NO: 22 were then translated to their respective protein products with Intelligenetic's program TRANSL.

One of these two novel human cDNA sequences (SEQ ID NO: 37) comprises 1456 bp:

30

10 20 30 40 50 60

- 43 -

	CGACGCAGAG	CAGCGCCCTG	GCCGGGCCAA	GCAGGAGCCG	GCATCATGGA	TTCCTTCAAG
	70	80	90	100	110	120
	GTAGTGCTGG	AGGGGCCAGC	ACCTTGGGGC	TTCCGGCTGC	AAGGGGGCAA	GGACTTCAAT
	130	140	150	160	170	180
5	GTGCCCCTCT	CCATTTCCCG	GCTCACTCCT	GGGGCAAAG	CGGCGCAGGC	CGGAGTGGCC
	190	200	210	220	230	240
	GTGGGTGACT	GGGTGCTGAG	CATCGATGGC	GAGAATGCGG	GTAGCCTCAC	ACACATCGAA
	250	260	270	280	290	300
	GCTCAGAACA	AGATCCGGGC	CTGCGGGGAG	CGCCTCAGCC	TGGGCCTCAG	CAGGGCCCAG
10	310	320	* 330	340	* 350	360
	CCGGTTCAGA	GCAAACCGCA	GAAG <u>GTGCAG</u>	ACCCCTGACA	<u>A</u> ACAGCCGCT	CCGACCGCTG
	370	380	390	400	410	420
	GTCCCAGATG	CCAGCAAGCA	GCGGCTGATG	GAGAACACAG	AGGACTGGCG	GCCGCGGCCG
	430	440	450	460	470	480
15	GGGACAGGCC	AGTCGCGTTC	CTTCCGCATC	CTTGCCCACC	TCACAGGCAC	CGAGTTCATG
	490	500	510	520	530	540
	CAAGACCCGG	ATGAGGAGCA	CCTGAAGAAA	TCAAGCCAGG	TGCCCAGGAC	AGAAGCCCCA
	550	560	570	580	590	600
	GCCCCAGCCT	CATCTACACC	CCAGGAGCCC	TGGCCTGGCC	CTACCGCCCC	CAGCCCTACC
20	610	620	630	640	650	660
	AGCCGCCCGC	CCTGGGCTGT	GGACCCTGCG	TTTGCCGAGC	GCTATGCCCC	GGACAAAACG
	670	680	690	700	710	720
	AGCACAGTGC	TGACCCGGCA	CAGCCAGCCG	GCCACGCCCA	CGCCGCTGCA	GAGCCGCACC
	730	740	750	760	770	780
25	TCCATTGTGC	AGGCAGCTGC	CGGAGGGGTG	CCAGGAGGGG	GCAGCAACAA	CGGCAAGACT
	790	800	810	820	830	840
	CCCGTGTGTC	ACCAGTGCCA	CAAGGTCATC	CGGGGCCGCT	ACCTGGTGGC	GTTGGGCCAC
	850	860	870	880	890	900

	GCGTACCACC	CGGAGGAGTT	TGTGTGTAGC	CAGTGTGGGA	AGGTCCTGGA	AGAGGGTGGC
	910	920	930	940	950	960
	TTCTTTGAGG	AGAAGGGCGC	CATCTTCTGC	CCACCATGCT	ATGACGTGCG	CTATGCACCC
	970	980	990	1000	1010	1020
5	AGCTGTGCCA	AGTGCAAGAA	GAAGATTACA	GGCGAGATCA	TGCACGCCCT	GAAGATGACC
	1030	1040	1050	1060	1070	1080
	TGGCACGTGC	ACTGCTTTAC	CTGTGCTGCC	TGCAAGACGC	CCATCCGGAA	CAGGGCCTTC
	1090	1100	1110	1120	1130	1140
	TACATGGAGG	AGGGCGTGCC	CTATTGCGAG	CGAGACTATG	AGAAGATGTT	TGGCACGAAA
10	1150	1160	1170	1180	1190	1200
	TGCCATGGCT	GTGACTTCAA	GATCGACGCT	GGGGACCGCT	TCCTGGAGGC	CCTGGGCTTC
1946 <del>- 2</del> 1 1 1	1210	1220	1230	1240	1250	1260
	AGCTGGCATG	ACACCTGCTT	CGTCTGTGCG	ATATGTCAGA	TCAACCTGGA	AGGAAAGACC
	1270	1280	1290	1300	1310	1320
15	TTCTACTCCA	AGAAGGACAG	GCCTCTCTGC	AAGAGCCATG	CCTTCTCTCA	TGTGTGAGCC
	1330	1340	1350	1360	1370	1380
	CCTTCTGCCC	ACAGCTGCCG	CGGTGGCCCC	TAGCCTGAGG	GGCCTGGAGT	CGTGGCCCTG
	1390	1400	1410	1420	1430	1440
	CATTTCTGGG	TAGGGCTGGC	AATGGTTGCC	TTAACCCTGG	CTCCTGGCCC	GAGCCTGGGC
20	1450					
	TCCCGGGCCC	TGCCCA				

Reading frame shifts caused by the deletion of a 119 bp fragment (between \*) and the addition of a 17 bp fragment (underlined) results in a truncated gene product having the following derived amino acid sequence (SEQ ID NO: 38):

10 20 30 40 50 60

	MDSFKVVLEG	PAPWGFRLQG	GKDFNVPLSI	SRLTPGGKAA	QAGVAVGDWV	LSIDGENAGS
	70	80	90	100	110	120
	LTHIEAQNKI	RACGERLSLG	LSRAQPVQNK	PQK <u>VOTPDK</u> Q	PLRPLVPDAS	KQRLMENTED
	130	140	150	160	170	180
5	WRPRPGTGQS	RSFRILAHLT	GTEFMQDPDE	EHLKKSSQVP	RTEAPAPASS	TPQEPWPGPT
	190	200	210	220	230	240
	APSPTSRPPW	AVDPAFAERY	APDKTSTVLT	RHSQPATPTP	LQSRTSIVQA	AAGGVPGGGS
,	250	260	270	280	290	300
	NNGKTPVCHQ	CHQVIRARYL	VALGHAYHPE	EFVCSQCGKV	LEEGGFFEEK	GAIFCPPCYD
10	310	320	330	340	350	360
	VRYAPSCAKC	KKKITGEIMH	ALKMTWHVLC	FTCAACKTPI	RNRAFYMEEG	VPYCERDYEK
	370	380	390	400	410	420
	MFGTKCQWCD	FKIDAGDRFL	EALGFSWHDT	CFVCAICQIN	LEGKTFYSKK	DRPLCKSHAF
	SHV					

This 423 amino acid protein demonstrates 100% homology to the protein shown in Sequence ID No. 10, except for the sequence in the highlighted area (amino acids 94-99), which are due to the nucleotide changes depicted above.

The second novel human heart cDNA sequence (SEQ ID NO: 39) comprises 1575 bp:

	10	20	30	40	50	60
	CGACGCAGAG	CAGCGCCCTG	GCCGGGCCAA	GCAGGAGCCG	GCATCATGGA	TTCCTTCAAG
	70	80	90	100	110	120
25	GTAGTGCTGG	AGGGGCCAGC	ACCTTGGGGC	TTCCGGCTGC	AAGGGGGCAA	GGACTTCAAT
	130	140	150	160	170	180
	GTGCCCCTCT	CCATTTCCCG	GCTCACTCCT	GGGGGCAAAG	CGGCGCAGGC	CGGAGTGGCC
	190	200	210	220	230	240

15

	GTGGGTGACT	GGGTGCTGAG	CATCGATGGC	GAGAATGCGG	GTAGCCTCAC	ACACATCGAA
	250	260	270	280	290	300
	GCTCAGAACA	AGATCCGGGC	CTGCGGGGAG	CGCCTCAGCC	TGGGCCTCAG	CAGGGCCCAG
	310	320	330	340	350	360
5	CCGGTTCAGA	GCAAACCGCA	GAAGGCCTCC	GCCCCCGCCG	CGGACCCTCC	GCGGTACACC
	370	380	390	400	410	420
	TTTGCACCCA	GCGTCTCCCT	CAACAAGACG	GCCCGGCCCT	TTGGGGCGCC	CCCGCCCGCT
	430	440	450	460	470	480
	GACAGCGCCC	CGCAACAGAA	TGG <u>GTGCAGA</u>	CCCCTGACAA	ACAGCCGCTC	CGACCGCTGG
10	490	500	510	520	530	540
	TCCCAGATGC	CAGCAAGCAG	CGGC <u>TGA</u> TGG	AGAACACAGA	GGACTGGCGG	CCGCGGCCGG
	550	560	570	580	590	600
	GGACAGGCCA	GTCGCGTTCC	TTCCGCATCC	TTGCCCACCT	CACAGGCACC	GAGTTCATGC
	610	620	630	640	650	660
15	AAGACCCGGA	TGAGGAGCAC	CTGAAGAAAT	CAAGCCAGGT	GCCCAGGACA	GAAGCCCCAG
	670	680	690	700	710	720
	CCCCAGCCTC	ATCTACACCC	CAGGAGCCCT	GGCCTGGCCC	TACCGCCCCC	AGCCCTACCA
	730	740	750	. 760	770	780
	ecceccecc	CTGGGCTGTG	GACCCTGCGT	TTGCCGAGCG	CTATGCCCCG	GACAAAACGA
20	790	800	810	820	830	840
	GCACAGTGCT	GACCCGGCAC	AGCCAGCCGG	CCACGCCCAC	GCCGCTGCAG	AGCCGCACCT
	850	860	870	880	. 890	900
	CCATTGTGCA	GGCAGCTGCC	GGAGGGGTGC	CAGGAGGGG	CAGCAACAAC	GGCAAGACTC
	910	920	930	940	950	960
25	CCGTGTGTCA	CCAGTGCCAC	AAGGTCATCC	GGGGCCGCTA	CCTGGTGGCG	TTGGGCCACG
	970	980	990	1000	1010	1020
	CGTACCACCC	GGAGGAGTTT	GTGTGTAGCC	AGTGTGGGAA	GGTCCTGGAA	GAGGGTGGCT
	1030	1040	1050	1060	1070	1080

	TCTTTGAGGA	GAAGGGCGCC	ATCTTCTGCC	CACCATGCTA	TGACGTGCGC	TATGCACCCA
	1090	1100	1110	. 1120	1130	1140
	GCTGTGCCAA	GTGCAAGAAG	AAGATTACAG	GCGAGATCAT	GCACGCCCTG	AAGATGACCT
	1150	1160	1170	1180	1190	1200
5	GGCACGTGCA	CTGCTTTACC	TGTGCTGCCT	GCAAGACGCC	CATCCGGAAC	AGGGCCTTCT
	1210	1220	1230	1240	1250	1260
	ACATGGAGGA	GGGCGTGCCC	TATTGCGAGC	GAGACTATGA	GAAGATGTTT	GGCACGAAAT
	1270	1280	1290	1300	1310	1320
	GCCATGGCTG	TGACTTCAAG	ATCGACGCTG	GGGACCGCTT	CCTGGAGGCC	CTGGGCTTCA
10	1330	1340	1350	1360	1370	1380
	GCTGGCATGA	CACCTGCTTC	GTCTGTGCGA	TATGTCAGAT	CAACCTGGAA	GGAAAGACCT
=	1390	1400	1410	1420	1430	1440
	TCTACTCCAA	GAAGGACAGG	CCTCTCTGCA	AGAGCCATGC	CTTCTCTCAT	GTGTGAGCCC
	1450	1460	1470	1480	1490	1500
15	CTTCTGCCCA	CAGCTGCCGC	GGTGGCCCCT	AGCCTGAGGG	GCCTGGAGTC	GTGGCCCTGC
	1510	1520	1530	1540	1550	1560
	ATTTCTGGGT	AGGGCTGGCA	ATGGTTGCCT	TAACCCTGGC	TCCTGGCCCG	AGCCTGGGCT
	1570					
	CCCGGGCCCT	GCCCA				

Reading frame shifts caused by the addition of a 17 bp fragment (bolded, italicized and underlined) results in an early translation stop codon at position 565-567 (underlined).

The derived amino acid sequence (SEQ ID NO: 40) consists of 153 amino acids:

**25**<sub>.</sub>

60	50	40	30	20	10
LSIDGENAGS	QAGVAVGDWV	SRLTPGGKAA	GKDFNVPLSI	PAPWGFRLQG	MDSFKVVLEG
120	110	100	90	80	70

LTHIEAQNKI RACGERLSLG LSRAQPVQSK PQKASAPAAD PPRYTFAPSV SLNKTARPFG

130 140 150

#### APPPADSAPO ONGCRPLTNS RSDRWSOMPA SSG

This protein demonstrates 100% homology to SEQ ID NO: 10 until amino acid 94, where the addition of the 17 bp fragment depicted in the nucleotide sequence results in a frame shift. Over amino acids 94-153, the protein is not homologous to SEQ ID NO: 10. Amino acids 154-457 in SEQ ID NO: 10 are not present due to the early stop codon depicted in nucleotide sequence.

## EXAMPLE 32: Genomic HLMP-1 Nucleotide Sequence

Applicants have identified the genomic DNA sequence encoding HLMP-1, including putative regulatory elements associated with HLMP-1 expression. The entire genomic sequence is shown in SEQ ID. NO: 41. This sequence

was derived from AC023788 (clone RP11-564G9), Genome Sequencing Center, Washington University School of Medicine, St. Louis, MO.

The putative promoter region for HLMP-1 spans nucleotides 2,660-8,733 in SEQ ID NO: 41. This region comprises, among other things, at least ten potential glucocorticoid response elements ("GREs") (nucleotides 6148-6153, 6226-6231, 6247-6252, 6336-6341, 6510-6515, 6552-6557, 6727-6732, 6752-6757, 7738-7743, and 8255-8260), twelve potential Sma-2 homologues to Mothers against Drosophilla decapentaplegic ("SMAD") binding element sites (nucleotides 3569-3575, 4552-4558, 4582-4588, 5226-5232, 6228-6234, 6649-6655, 6725-6731, 6930-6936, 7379-7384, 7738-7742, 8073-8079, and 8378-8384), and three TATA boxes (nucleotides 5910-5913, 6932-6935, and 7380-7383). The three TATA boxes, all of the GREs, and eight of the SMAD binding elements ("SBEs") are grouped in the region spanning nucleotides 5,841-8,733 in SEQ ID NO: 41. These regulatory elements can be used, for example, to regulate expression of exogenous nucleotide sequences encoding proteins involved in the process of bone formation. This would permit systemic administration of therapeutic factors or genes relating to bone formation and repair, as well as factors or genes associated with tissue differentiation and development.

In addition to the putative regulatory elements, 13 exons corresponding to the nucleotide sequence encoding HLMP-1 have been identified. These exons span the following nucleotides in SEQ ID NO: 41:

	Exon 1	8733-8767
30	Exon 2	9790-9895
	Exon 3	13635-13787

5

10

15

20

	Exon 4	13877-13907
	Exon 5	14387-14502
	Exon 6	15161-15297
	Exon 7	15401-15437
5	Exon 8	16483-16545
	Exon 9	16689-16923
	Exon 10	18068-18248
	Exon 11	22117-22240
	Exon 12	22323-22440
10	Exon 13	22575-22911

In HLMP-2 there is another exon (Exon 5A), which spans nucleotides 14887-14904.

All cited publications and patents are hereby incorporated by reference in their entirety.

15 While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be appreciated by one skilled in the art from reading this disclosure that various changes in form and detail can be made without departing from the true scope of the invention.

#### We claim:

10

15

20

25

30

1. An isolated nucleic acid molecule comprising a SEQ ID NO: 37 or SEQ ID NO: 39.

- An isolated human LMP protein encoded by SEQ ID NO: 37 or SEQ
   ID NO: 39.
  - 3. A vector comprising the isolated nucleic acid molecule of claims 1.
  - 4. A host cell comprising the vector of claim 3.
  - The host cell of claim 4, wherein the host cell is selected from the group consisting of prokaryotic cells, yeast cells and mammalian cells
  - 6. The isolated nucleic acid molecule of claim 1, further comprising a label.
  - 7. A human LIM mineralization protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 38 and SEQ ID NO: 40.
  - 8. A monoclonal antibody specific for a HLMP-2 (SEQ ID NO: 38) or HLMP-3 (SEQ ID NO: 40).
  - 9. A method of inducing bone formation comprising transfecting osteogenic precursor cells or peripheral blood leukocytes with an isolated nucleic acid molecule comprising SEQ ID NO: 39.
  - 10. The method of claim 9, wherein the isolated nucleic acid molecule is in a vector.
    - 11. The method of claim 10, wherein the vector is an expression vector.
    - 12. The method of claim 11, wherein the vector is a plasmid.
    - 13. The method of claim 11, wherein the vector is a virus.
    - 14. The method of claim 13, wherein the virus is an adenovirus
    - 15. The method of claim 13, wherein the virus is a retrovirus.
  - 16. The method of claim 9, wherein the osteogenic precursor cells or peripheral blood leukocytes are transfected *ex vivo*.
  - 17. The method of claim 9, wherein the osteogenic precursor cells are transfected *in vivo* by direct injection of the isolated nucleic acid molecule.

18. The method of claim 9, wherein the LIM mineralization protein is HLMP-3 (SEQ ID NO: 40).

- 19. A method of fusing a spine, comprising:
- (a) transfecting osteogenic precursor cells or peripheral blood leukocytes with an isolated nucleic acid molecule comprising SEQ ID NO: 39;
  - (b) admixing the transfected osteogenic precursor cells or peripheral blood leukocytes with a matrix; and
- (c) contacting the matrix with the spine;
   wherein expression of the nucleotide sequence causes mineralized
   bone formation in the matrix.
  - 20. The method of claim 19, wherein the osteogenic precursor cells or peripheral blood cells are transfected *ex vivo*.
  - 21. A method of inducing systemic bone formation in a mammalian host in need thereof, comprising:
  - a) transfecting osteogenic precursor cells or peripheral blood leukocytes with a vector that is stablely maintained in the osteogenic precursor cells or peripheral blood leukocytes, the vector comprising SEQ ID NO: 39 operably linked to a regulatable promoter, wherein the regulatable promoter responds to an exogenous control compound; and
  - (b) administering to the host, as needed, an amount of the exogenous control substance effective to cause expression of SEQ ID NO: 39.

5

15

22. A method of stimulating production of an osteogenic soluble factor by an osteogenic cell, comprising:

- (a) transfecting the osteogenic cell or peripheral blood leukocyte with an isolated nucleic acid molecule comprising SEQ ID NO: 39; and
  - (b) overexpressing the isolated nucleic acid molecule.
  - 23. An osteogenic soluble factor produced by the method of claim 22.
- 24. The osteogenic soluble factor of claim 23, wherein the osteogenic factor is a protein.
- 25. A method of inhibiting the expression of HLMP-2 or HLMP-3
   10 comprising transfecting a cell wherein HLMP-2 or HLMP-3 is expressed with an antisense oligonucleotide.
  - 26. The method of claim of claim 17, wherein the isolated nucleic acid molecule is in a vector selected from the group consisting of a plasmid and a virus.
    - 27. The method of claim 26, wherein the vector is a plasmid, which plasmid is directly injected into muscle tissue.
  - 28. A method of inhibiting bone formation comprising transfecting osteogenic precursor cells or peripheral blood leukocytes with an isolated nucleic acid molecule comprising SEQ ID NO: 37.
  - 29. The method of claim 28, wherein the isolated nucleic acid molecule is in a vector.
    - 30. The method of claim 29, wherein the vector is an expression vector.
    - 31. The method of claim 30, wherein the vector is a plasmid.
    - 32. The method of claim 30, wherein the vector is a virus.
    - 33. The method of claim 32, wherein the virus is an adenovirus
    - 34. The method of claim 32, wherein the virus is a retrovirus.
    - 35. The method of claim 28, wherein the osteogenic precursor cells or peripheral blood leukocytes are transfected *ex vivo*.
- 36. The method of claim 28, wherein the osteogenic precursor cells aretransfected *in vivo* by direct injection of the isolated nucleic acid molecule.

5

15

20

37. The method of claim 28, wherein the LIM mineralization protein is HLMP-2 (SEQ ID NO: 38).

#### SEQUENCE LISTING

- <110> Boden M.D., Scott D Hair Ph.D., Gregory A
- <120> LIM Mineralization Protein Splice Variants
- <130> splice variant LMP
- <140> 6148.0222-00304
- <141> 2000-04-28
- <150> 60/132,021
- <151> 1999-04-30
- <160> 42
- <170> PatentIn Ver. 2.1
- <210> 1
- <211> 457
- <212> PRT
- <213> Rattus norvegicus
- <400> 1
- Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe
  1 5 10 15
- Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg 20 25 30
- Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp
  35 40 45
- Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
  50 55 60
- Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly
  65 70 75 80
- Leu Ser Arg Ala Gln Pro Ala Gln Ser Lys Pro Gln Lys Ala Leu Thr 85 90 95
- Pro Pro Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Ala Ser Leu 100 105 110
- Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Thr Asp Ser Ala 115 120 125
- Leu Ser Gln Asn Gly Gln Leu Leu Arg Gln Leu Val Pro Asp Ala Ser .130 135 140
- Lys Gln Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly 145 150 155 160

Thr Gly Gln Ser Arg Ser Phe Arg Ile Leu Ala His Leu Thr Gly Thr 175

Glu Phe Met Gln Asp Pro Asp Glu Glu Phe Met Lys Lys Ser Ser Gln 180

Val Pro Arg Thr Glu Ala Pro Ala Pro Ala Ser Thr Ile Pro Gln Glu 205

Ser Trp Pro Gly Pro Thr Thr Pro Ser Pro Thr Ser Arg Pro Pro Trp 210

Ala Val Asp Pro Ala Phe Ala Glu Arg Tyr Ala Pro Asp Lys Thr Ser 240

Thr Val Leu Thr Arg His Ser Gln Pro Ala Thr Pro Thr Pro Leu Gln 255

Asn Arg Thr Ser Ile Val Gln Ala Ala Ala Gly Gly Gly Thr Gly Gly Gly Ser Asn Asn Gly Lys Thr Pro Val Cys His Gln Cys His Lys Ile

- 275 280 285
- Ile Arg Gly Arg Tyr Leu Val Ala Leu Gly His Ala Tyr His Pro Glu 290 295 300
- Glu Phe Val Cys Ser Gln Cys Gly Lys Val Leu Glu Glu Gly Gly Phe 305 310 315 320
- Phe Glu Glu Lys Gly Ala Ile Phe Cys Pro Ser Cys Tyr Asp Val Arg 325 330 335
- Tyr Ala Pro Ser Cys Ala Lys Cys Lys Lys Lys Ile Thr Gly Glu Ile 340 345 350
- Met His Ala Leu Lys Met Thr Trp His Val Pro Cys Phe Thr Cys Ala 355 360 365
- Ala Cys Lys Thr Pro Ile Arg Asn Arg Ala Phe Tyr Met Glu Glu Gly 370 375 380
- Ala Pro Tyr Cys Glu Arg Asp Tyr Glu Lys Met Phe Gly Thr Lys Cys 385 390 395 400
- Arg Gly Cys Asp Phe Lys Ile Asp Ala Gly Asp Arg Phe Leu Glu Ala 405 410 415
- Leu Gly Phe Ser Trp His Asp Thr Cys Phe Val Cys Ala Ile Cys Gln
  420 425 430
- Ile Asn Leu Glu Gly Lys Thr Phe Tyr Ser Lys Lys Asp Lys Pro Leu
  435 440 445
- Cys Lys Ser His Ala Phe Ser His Val 450 455

```
<210> 2
<211> 1696
<212> DNA
<213> Rattus norvegicus
<400> 2
gcacgaggat cccagcgcgg ctcctggagg ccgccaggca gccgcccagc cgggcattca 60
ggagcaggta ccatggattc cttcaaggta gtgctggagg gacctgcccc ttggggcttc 120
cgtctgcaag ggggcaagga cttcaacgtg cccctctcca tctctcggct cactcctgga 180
ggcaaggccg cacaggccgg tgtggccgtg ggagactggg tactgagtat cgacggtgag 240
aacgccggaa gcctcacaca cattgaagcc cagaacaaga tccgtgcctg tggggagcgc 300
ctcagcctgg gtcttagcag agcccagcct gctcagagca aaccacagaa ggccctgacc 360
cetecegecg acceeegag gtacaetttt geaccaageg ceteceteaa caagaeggee 420
cggcccttcg gggcaccccc acctactgac agcgccctgt cgcagaatgg acagctgctc 480
agacagetgg tecetgatge cagcaageag eggetgatgg agaataetga agactggege 540
cegeggecag ggacaggeca gtecegttee tteegeatee ttgeteacet caegggeaca 600
gagttcatgc aagacccgga tgaggaattc atgaagaagt caagccaggt gcccaggaca 660
gaagccccag ccccagcctc aaccataccc caggaatcct ggcctggccc caccaccccc 720
agececacca geogeocace etgggeogta gateetgeat ttgetgageg etatgeecca 780
gacaaaacca gcacagtgct gacccgacac agccagccag ccacacctac gcctctgcag 840
aaccgcacct ccatagttca ggctgcagct ggagggggca caggaggagg cagcaacaat 900
ggcaagacgc ctgtatgcca ccagtgccac aagatcatcc-geggccgata cctggtagca-960
ctgggccacg cgtaccatcc tgaggaattt gtgtgcagcc agtgtgggaa ggtcctggaa 1020
gagggtggct tcttcgagga gaagggagct atcttttgcc cctcctgcta tgatgtgcgc 1080
tatgcaccca gctgtgccaa atgcaagaag aagatcactg gagagatcat gcatgcgctg 1140
aagatgacct ggcatgttcc ctgcttcacc tgtgcagcct gcaaaacccc tatccgcaac 1200
agggetttet acatggagga gggggetece tactgegage gagattaega gaagatgttt 1260
ggcacaaagt gtcgcggctg tgacttcaag atcgatgccg gggaccgttt cctggaagcc 1320
ctgggtttca gctggcatga tacgtgtttt gtttgcgcaa tatgtcaaat caacttggaa 1380
ggaaagacct tctactccaa gaaggacaag cccctgtgca agagccatgc cttttcccac 1440
gtatgagcac ctcctcacac tactgccacc ctactctgcc agaagggtga taaaatgaga 1500
gagetetete tecetegace tttetgggtg gggetggeag ceattgteet ageettgget 1560
cctggccaga tcctggggct ccctcctcac agtccccttt cccacacttc ctccaccacc 1620
accaccgtca ctcacaggtg ctagcctcct agccccagtt cactctggtg tcacaataaa 1680
                                                                  1696
cctgtatgta gctgtg
<210> 3
<211> 260
<212> DNA
<213> Rattus norvegicus
<400> 3
ttctacatgg aggaggggc tccctactgc gagcgagatt acgagaagat gtttggcaca 60
aagtgtcgcg gctgtgactt caagatcgat gccggggacc gtttcctgga agccctgggt 120
ttcagctggc atgatacgtg ttttgtttgc gcaatatgtc aaatcaactt ggaaggaaag 180
accttctact ccaagaagga caagccctg tgcaagagcc atgccttttc ccacgtatga 240
                                                                  260
gcacctcctc acactactgc
<210> 4
<211> 16
<212> DNA
```

<213> MMLV

```
<400> 4
                                                                  16
aagctttttt tttttg
<210> 5
<211> 13
<212> DNA
<213> MMLV
<400> 5
                                                                  13
aagcttggct atg
<210> 6
<211> 223
<212> DNA
<213> Homo sapiens
<400> 6
atcettgete aceteaeggg cacegagtte atgeaagace eggatgagga geacetgaag 60
aaatcaagcc aggtgcccag gacagaagcc ccagccccag cctcatctac accccaggag 120
ccctggcctg gccctaccgc ccccagccct accagccgc cgccctgggc tgtggaccct 180
gegtttgeeg agegetatge eecagacaaa accageacag tge
<210> 7
<211> 717
<212> DNA
<213> Homo sapiens
<400> 7
atggattcct tcaaggtagt gctggagggg ccagcacctt ggggcttccg gctgcaaggg 60
ggcaaggact tcaatgtgcc cctctccatt tcccggctca ctcctggggg caaagcggcg 120
caggccggag tggccgtggg tgactgggtg ctgagcatcg atggcgagaa tgcgggtagc 180
ctcacacaca tegaagetea gaacaagate egggeetgeg gggagegeet eageetggge 240
ctcagcaggg cccagccggt tcagagcaaa ccgcagaagg cctccgcccc cgccgcggac 300
cctccgcggt acacctttgc acccagcgtc tccctcaaca agacggcccg gccctttggg 360
gegececege egetgaeag egeceegeaa eagaatggae ageegeteeg acegetggte 420
ccagatgcca gcaagcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 480
acaggecagt egegtteett eegeateett geecaeetea eaggeacega gtteatgeaa 540
qaccoggatg aggagcacct gaagaaatca agccaggtgc ccaggacaga agccccagcc 600
ccagcctcat ctacacccca ggagccctgg cctggcccta ccgcccccag ccctaccagc 660
egecegeeet gggetgtgga eeetgegttt geegageget atgeeeegga caaaaeg
<210> 8
<211> 1488
<212> DNA
<213> Homo sapiens
atcgatggcg agaatgcggg tagcctcaca cacatcgaag ctcagaacaa gatccgggcc 60
tgcggggagc gcctcagcct gggcctcagc agggcccagc cggttcagag caaaccgcag 120
aaggeeteeg eeceegeege ggaceeteeg eggtacacet ttgcacecag egteteeete 180
aacaagacgg cccggccctt tggggcgccc ccgcccgctg acagcgcccc gcaacagaat 240
qqacaqccqc tccqaccqct ggtcccaqat gccaqcaaqc agcggctgat ggagaacaca 300
```

```
gaggactggc ggccgcggcc ggggacaggc cagtcgcgtt ccttccgcat ccttgcccac 360
ctcacaggca ccgagttcat gcaagacccg gatgaggagc acctgaagaa atcaagccag 420
cctaccgccc ccagccctac cagccgcccg ccctgagctg tggaccctgc gtttgccgag 540
egetatgece eggacaaaac gagcacagtg etgaceegge acagecagee ggecacgee 600
acgccgctgc agagccgcac ctccattgtg caggcagctg ccggaggggt gccaggaggg 660
ggcagcaaca acggcaagac tcccgtgtgt caccagtgcc acaaggtcat ccggggccgc 720
tacctggtgg cgttgggcca cgcgtaccac ccggaggagt ttgtgtgtag ccagtgtggg 780
aaggtcctgg aagagggtgg cttctttgag gagaagggcg ccatcttctg cccaccatgc 840
tatgacgtgc gctatgcacc cagctgtgcc aagtgcaaga agaagattac aggcgagatc 900
atgcacgccc tgaagatgac ctggcacgtg cactgcttta cctgtgctgc ctgcaagacg 960
cccatccgga acagggcctt ctacatggag gagggcgtgc cctattgcga gcgagactat 1020
gagaagatgt ttggcacgaa atgccatggc tgtgacttca agatcgacgc tggggaccgc 1080
ttcctggagg ccctgggctt cagctggcat gacacctgct tcgtctgtgc gatatgtcag 1140
atcaacctgg aaggaaagac cttctactcc aagaaggaca ggcctctctg caagagccat 1200
geettetete atgtgtgage eeettetgee cacagetgee geggtggeee etageetgag 1260
gggcctggag tcgtggccct gcatttctgg gtagggctgg caatggttgc cttaaccctg 1320
getectggee egageetggg etecegggee eetgeecace cacettatee teceacecca 1380
ctccctccac caccacagca caccggtgct ggccacacca gccccctttc acctccagtg 1440
ccacaataaa cctgtaccca gctgaattcc aaaaaaatcca aaaaaaaa
```

<400> 9 atggatteet teaaggtagt getggagggg ceageacett ggggetteeg getgeaaggg 60 ggcaaggact tcaatgtgcc cctctccatt tcccggctca ctcctggggg caaagcggcg 120 caggccggag tggccgtggg tgactgggtg ctgagcatcg atggcgagaa tgcgggtagc 180 ctcacacaca tcgaagctca gaacaagatc cgggcctgcg gggagcgcct cagcctgggc 240 ctcagcaggg cccagccggt tcagagcaaa ccgcagaagg cctccgcccc cgccgcggac 300 cctccgcggt acacctttgc acccagcgtc tccctcaaca agacggcccg gccctttggg 360 gegeeeege eegetgacag egeeegeaa cagaatggae ageegeteeg acegetggte 420 ccagatgcca gcaagcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 480 acaggccagt cgcgttcctt ccgcatcctt gcccacctca caggcaccga gttcatgcaa 540 gacccggatg aggagcacct gaagaaatca agccaggtgc ccaggacaga agccccagcc 600 ccagcctcat ctacacccca ggagccctgg cctggcccta ccgccccag ccctaccagc 660 cgcccgccct gggctgtgga ccctgcgttt gccgagcgct atgccccgga caaaacgagc 720 acagtgetga eeeggeacag eeageeggee aegeeeaege egetgeagag eegeacetee 780 attgtgcagg cagctgccgg aggggtgcca ggagggggca gcaacaacgg caagactccc 840 gtgtgtcacc agtgccacaa ggtcatccgg ggccgctacc tggtggcgtt gggccacgcg 900 taccaccegg aggagtttgt gtgtageeag tgtgggaagg teetggaaga gggtggette 960 tttgaggaga agggcgccat cttctgccca ccatgctatg acgtgcgcta tgcacccagc 1020 tgtgccaagt gcaagaagaa gattacaggc gagatcatgc acgccctgaa gatgacctgg 1080 cacgtgcact gctttacctg tgctgcctgc aagacgccca tccggaacag ggccttctac 1140 atggaggagg gcgtgcccta ttgcgagcga gactatgaga agatgtttgg cacgaaatgc 1200 catggctgtg acttcaagat cgacgctggg gaccgcttcc tggaggccct gggcttcagc 1260 tggcatgaca cctgcttcgt ctgtgcgata tgtcagatca acctggaagg aaagaccttc 1320 tactccaaga aggacaggcc tctctgcaag agccatgcct tctctcatgt gtgagcccct 1380 tetgeccaca getgeegegg tggeceetag cetgagggge etggagtegt ggecetgeat 1440 ttctgggtag ggctggcaat ggttgcctta accetggctc ctggcccgag cctgggctcc 1500 egggeeetg eccaeceace thateetece accecaete etceaecace acageaeace 1560 ggtgctggcc acaccagccc cetttcacct ccagtgccac aataaacctg tacccagctg 1620 aattccaaaa aatccaaaaa aaaa

<210> 10 <211> 457

<212> PRT

<213> Homo sapiens

<400> 10

Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe
1 5 10 15

Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg 20 25 30

Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp 35 40 45

Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
50 55 60

Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly 65 70 75 80

Leu Ser Arg Ala-Ghn-Pro Val Gln Ser Lys-Pro Gln Lys-Ala-Ser Ala 85 90 95

Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Val Ser Leu 100 105 110

Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala 115 120 125

Pro Gln Gln Asn Gly Gln Pro Leu Arg Pro Leu Val Pro Asp Ala Ser 130 135 140

Lys Gln Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly 145 150 155 160

Thr Gly Gln Ser Arg Ser Phe Arg Ile Leu Ala His Leu Thr Gly Thr
165 170 175

Glu Phe Met Gln Asp Pro Asp Glu Glu His Leu Lys Lys Ser Ser Gln 180 185 190

Val Pro Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu
195 200 205

Pro Trp Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro Trp 210 215 220

Ala Val Asp Pro Ala Phe Ala Glu Arg Tyr Ala Pro Asp Lys Thr Ser 225 230 235 240

Thr Val Leu Thr Arg His Ser Gln Pro Ala Thr Pro Thr Pro Leu Gln 245 250 255

Ser Arg Thr Ser Ile Val Gln Ala Ala Ala Gly Gly Val Pro Gly Gly

265 260 270 Gly Ser Asn Asn Gly Lys Thr Pro Val Cys His Gln Cys His Lys Val 280 Ile Arg Gly Arg Tyr Leu Val Ala Leu Gly His Ala Tyr His Pro Glu 295 Glu Phe Val Cys Ser Gln Cys Gly Lys Val Leu Glu Glu Gly Gly Phe 305 Phe Glu Glu Lys Gly Ala Ile Phe Cys Pro Pro Cys Tyr Asp Val Arg 325 330 Tyr Ala Pro Ser Cys Ala Lys Cys Lys Lys Lys Ile Thr Gly Glu Ile 345 Met His Ala Leu Lys Met Thr Trp His Val His Cys Phe Thr Cys Ala 355 . 360 Ala Cys Lys Thr Pro Ile Arg Asn Arg Ala Phe Tyr Met Glu Glu Gly 375 380 Val Pro Tyr Cys Glu Arg Asp Tyr Glu Lys Met Phe Gly Thr Lys Cys 390 His Gly Cys Asp Phe Lys Ile Asp Ala Gly Asp Arg Phe Leu Glu Ala Leu Gly Phe Ser Trp His Asp Thr Cys Phe Val Cys Ala Ile Cys Gln 425 Ile Asn Leu Glu Gly Lys Thr Phe Tyr Ser Lys Lys Asp Arg Pro Leu Cys Lys Ser His Ala Phe Ser His Val <210> 11 <211> 22 <212> DNA <213> Rattus norvegicus <400> 11 22 gccagggttt tcccagtcac ga

<210> 12 <211> 22 <212> DNA <213> Rattus norvegicus

gecagggttt teccagteae ga 22

WO 00/66178	PCT/US00/11664
<210> 13	
<211> 22	
<212> DNA	
<213> Homo sapiens	·
<400> 13	
tcttagcaga gcccagcctg ct	22
<210> 14	
<211> 22	
<212> DNA	
<213> Homo sapiens	
<400> 14	
gcatgaacte tgtgeeegtg ag	22
<210> 15	
<211> 20	
<212> DNA	
<213> Rattus norvegicus	
<400> 15	gan iz a maga a sa a sa a sa a sa a sa a sa a
atecttgete aceteaeggg	20
<210> 16	
<211> 22	
<212> DNA	
<213> Rattus norvegicus	
<400> 16	
gcactgtgct ggttttgtct gg	22
•	
<210> 17	•
<211> 23	
<212> DNA <213> Homo sapiens	
<400> 17	
catggattcc ttcaaggtag tgc	23
<210> 18	
<211> 20	
<212> DNA	
<213> Homo sapiens	
<400> 18 gttttgtctg gggcagagcg	20
greengery gggeagageg	20
<210> 19	
<211> 44	

```
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: adaptor for
     .Marathon RACE reactions
<400> 19
ctaatacgac tcactatagg gctcgagcgg ccgcccgggc aggt
                                                                  44
<210> 20
<211> 27
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: PCR primer
      specific for Marathon RACE adaptor
<400> 20
                                                                  27
ccatcctaat acgactcact atagggc
 <210> 21
<211> 765
<212> DNA
<213> Homo sapiens
<400> 21
ccgttgtttg taaaacgacg cagagcagcg ccctggccgg gccaagcagg agccggcatc 60
atggatteet teaaggtagt getggagggg ceageacett ggggetteeg getgeaaggg 120
qqcaaqqact tcaatqtqcc ctcctccatt tcccggctca cctctggggg caaggccgtg 180
caggceggag tggcegtaag tgactgggtg etgagcateg atggegagaa tgegggtage 240
ctcacacaca tegaagetea gaacaagate egggeetgeg gggagegeet eageetggge 300
ctcaacaggg cccagccggt tcagaacaaa ccgcaaaagg cctccgcccc cgccgcggac 360
ceteegeggt acacetttge accaagegte teceteaaca agaeggeeeg geeettgggg 420
gegeeeege eegetgacag egeeeegeag cagaatggac ageegeteeg acegetggte 480
ccagatgcca gcaagcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 540
acaggecagt geegtteett tegeateett geteacetta caggeacega gtteatgeaa 600
gaccoggatg aggagcacct gaagaaatca agccaggtgc ccaggacaga agccccagcc 660
ccagcctcat ctacaccca ggagecctgg cctggcccta ccgcccccag ccctaccagc 720
cgcccgccct gggctgtgga ccctgcgttt gccgagcgct atgcc
<210> 22
<211> 1689
<212> DNA
<213> Homo sapiens
cgacgcagag cagegccetg geegggccaa geaggagccg geatcatgga ttcettcaag 60
gtagtgctgg aggggccagc accttggggc ttccggctgc aagggggcaa ggacttcaat 120
gtgcccctct ccatttcccg gctcactcct gggggcaaag cggcgcaggc cggagtggcc 180
gtgggtgact gggtgctgaq catcgatggc gagaatgcgg gtagcctcac acacatcgaa 240
```

getcagaaca agateeggge etgeggggag egectcagee tgggeetcag eagggeecag 300

```
ccqqttcaqa qcaaaccgca gaaggcctcc gccccgccg cggaccctcc gcggtacacc 360
tttqcaccca gcgtctccct caacaagacg gcccggccct ttggggcgcc cccgcccgct 420
gacagegece egeaacagaa tggacageeg eteegacege tggteecaga tgecageaag 480
cagcggctga tggagaacac agaggactgg cggccgcggc cggggacagg ccagtcgcgt 540
tectteegea teettgeeea eeteacagge accgagttea tgeaagaeee ggatgaggag 600
cacctgaaga aatcaagcca ggtgcccagg acagaagccc cagccccagc ctcatctaca 660
ccccaggagc cctggcctgg ccctaccgcc cccagcccta ccagccgccc gccctgggct 720
gtggaccetg cgtttgecga gcgctatgcc ccggacaaaa cgagcacagt gctgaccegg 780
cacagecage eggecaegee caegecgetg cagageegea cetecattgt geaggeaget 840
qeeqqaqqqq tgecaggagg gggcagcaac aacggcaaga ctcccgtgtg tcaccagtgc 900
cacaaggtca teeggggeeg ctacetggtg gegttgggee aegegtaeea eeeggaggag 960
tttgtgtgta gccagtgtgg gaaggtcctg gaagagggtg gcttctttga ggagaagggc 1020
gccatcttct gcccaccatg ctatgacgtg cgctatgcac ccagctgtgc caagtgcaag 1080
aaqaaqatta caqqcqaqat catqcacqcc ctqaaqatqa cctqqcacqt qcactqcttt 1140
acctgtgctg cctgcaagac gcccatccgg aacagggcct tctacatgga ggagggcgtg 1200
ccctattgcg agcgagacta tgagaagatg tttggcacga aatgccatgg ctgtgacttc 1260
aagategaeg etggggaeeg etteetggag geeetggget teagetggea tgacaeetge 1320
ttcqtctqtq cqatatgtca gatcaacctg gaaggaaaga ccttctactc caagaaggac 1380
aggeetetet qeaagageea tgeettetet catgtgtgag eeeettetge ceacagetge 1440
cgcggtggcc cctagcctga ggggcctgga gtcgtggccc tgcatttctg ggtagggctg 1500
qeaatqqttq ccttaaccct ggctcctggc ccgagcctgg gctcccgggc ccctgcccac 1560
ccaccttate eteccacee acteceteca ccaccacage acaceggtge tggccacace 1620
-aqcccccttt cacctccagt/gccacaataa/acctgtaccc agctgaattc--caaaaaaatcc/1680---
aaaaaaaa
<210> 23
<211> 22
<212> DNA
<213> Homo sapiens
<400> 23
                                                                   22
gcactgtgct cgttttgtcc gg
<210> 24
<211> 21
<212> DNA
<213> Homo sapiens
<400> 24
teettgetea eeteaeggge a
                                                                   21
<210> 25
<211> 30
<212> DNA
<213> Homo sapiens
<400> 25
tecteateeg ggtettgeat gaacteggtg
                                                                   30
<210> 26
<211> 28
<212> DNA
```

WO 00/66178	PCT/US00/11664
<213> Homo sapiens	
<400> 26 gcccccgccc gctgacagcg ccccgcaa	28
<210> 27 <211> 24 <212> DNA <213> Homo sapiens	
<400> 27 teettgetea ceteaeggge accg	24
<210> 28 <211> 22 <212> DNA <213> Homo sapiens	
<400> 28 gtaatacgac tcactatagg gc	22
<210> 29 <211> 23 <212> DNA <213> Rattus norvegicus	
<400> 29 gcggctgatg gagaatactg aag	23
<210> 30 <211> 23 <212> DNA <213> Rattus norvegicus	
<400> 30 atcttgtggc actggtggca tac	23
<210> 31 <211> 22 <212> DNA <213> Rattus norvegicus	
<400> 31 tgtgtcgggt cagcactgtg ct	22
<210> 32 <211> 1620 <212> DNA <213> Homo sapiens	

```
<400> 32
atggatteet teaaggtagt getggagggg ceageacett ggggetteeg getgeaaggg 60
ggcaaggact tcaatgtgcc cctctccatt tcccggctca ctcctggggg caaagcggcg 120
caggooggag tggccgtggg tgactgggtg ctgagcatcg atggcgagaa tgcgggtagc 180
ctcacacaca tegaagetea gaacaagate egggeetgeg gggagegeet eageetggge 240
ctcagcaggg cccagccggt tcagagcaaa ccgcagaagg cctccgccc cgccgcggac 300
cctccgcggt acacctttgc acccagcgtc tccctcaaca agacggcccg gccctttggg 360
gegeeeeege eegetgaeag egeeeegeaa cagaatggae ageegeteeg acegetggte 420
ccagatgcca gcaagcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 480
acaggecagt egegtteett eegeateett geecacetea eaggeacega gtteatgeaa 540
gacceggatg aggagcacct gaagaaatca agccaggtge ccaggacaga agccccagee 600
ccagceteat etacacecca ggagecetgg cetggeeeta eegeeeecag ecetaceage 660
egecegeeet gagetgtgga eeetgegttt geegageget atgeeeegga caaaacgage 720
acagtgetga cceggcacag ccagceggcc acgcccacgc egetgeagag cegcacetec 780
attgtgcagg cagctgccgg aggggtgcca ggagggggca gcaacaacgg caagactccc 840
gtgtgtcacc agtgccacaa ggtcatccgg ggccgctacc tggtggcgtt gggccacgcg 900
taccaccegg aggagtttgt gtgtagccag tgtgggaagg tcctggaaga gggtggcttc 960
tttgaggaga agggcgccat cttctgccca ccatgctatg acgtgcgcta tgcacccage 1020
tgtgccaagt gcaagaagaa gattacaggc gagatcatgc acgccctgaa gatgacctgg 1080
cacgtgcact gctttacctg tgctgcctgc aagacgccca tccggaacag ggccttctac 1140
atggaggagg gcgtgcccta ttgcgagcga gactatgaga agatgtttgg cacgaaatgc 1200
catggctgtg acttcaagat cgacgctggg gaccgcttcc tggaggccct gggcttcagc 1260
tggcatgaca cctgcttcgt ctgtgcgata tgtcagatca acctggaagg-aaagaccttc-1320
tactccaaga aggacaggec tetetgcaag agccatgeet teteteatgt gtgageeeet 1380
tetgeccaca getgeegegg tggeccctag cetgagggge etggagtegt ggecetgeat 1440
ttctgggtag ggctggcaat ggttgcctta accctggctc ctggcccgag cctgggctcc 1500
egggeeectg eccaeceace ttatectece accecaetee etceaecace acageacace 1560
ggtgctggcc acaccagccc cctttcacct ccagtgccac aataaacctg tacccagctg 1620
<210> 33
<211> 1665
<212> DNA
<213> Homo sapiens
<400> 33
cgacgcagag cagcgccctg gccgggccaa gcaggagccg gcatcatgga ttccttcaag 60
gtagtgctgg aggggccagc accttggggc ttccggctgc aagggggcaa ggacttcaat 120
gtgeceetet ccattteeeg geteacteet gggggeaaag eggegeagge eggagtggee 180
gtgggtgact gggtgctgag catcgatggc gagaatgcgg gtagcctcac acacatcgaa 240
geteagaaca agateeggge etgeggggag egeeteagee tgggeeteag eagggeecag 300
ceggttcaga gcaaacegca gaaggcetce gcececgeeg eggaceetce geggtacaec 360
tttgcaccca gcgtctccct caacaagacg gcccggccct ttggggcgcc cccgcccgct 420
```

ccctattgcg agcgagacta tgagaagatg tttggcacga aatgccatgg ctgtgacttc 1260 aagatcgacg ctggggaccg cttcctggag gccctgggct tcagctggca tgacacctgc 1320 ttcgtctgtg cgatatgtca gatcaacctg gaaggaaaga ccttctactc caagaaggac 1380 aggcctctct gcaagagcca tgccttctct catgtgtgag ccccttctgc ccacagctgc 1440 cgcggtggcc cctagcctga ggggcctgga gtcgtggcc tgcatttctg ggtagggctg 1500 gcaatggttg ccttaaccct ggctcctgca ccgagcctgg gctcccgggc ccctgccac 1560 ccaccttatc ctccacccc actcctcca ccaccacagc acaccggtgc tggccacacc 1620 agccccttt cacctccagt gccacaataa acctgtaccc agctg

<210> 34

<211> 223

<212> PRT

<213> Homo sapiens

<400> 34

Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe
1 5 10 15

Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg 20 25 30

Leu Thr Pro Gly Gly-Lys Ala Ala Gln Ala-Gly Val-Ala Val Gly Asp--- 35 40 45

Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
50 55 60

Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly
65 70 75 80

Leu Ser Arg Ala Gln Pro Val Gln Ser Lys Pro Gln Lys Ala Ser Ala 85 90 95

Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Val Ser Leu 100 105 110

Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala 115 120 125

Pro Gln Gln Asn Gly Gln Pro Leu Arg Pro Leu Val Pro Asp Ala Ser 130 135 140

Lys Gln Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly
145 150 155 160

Thr Gly Gln Ser Arg Ser Phe Arg Ile Leu Ala His Leu Thr Gly Thr
165 170 175

Glu Phe Met Gln Asp Pro Asp Glu Glu His Leu Lys Lys Ser Ser Gln 180 185 190

Val Pro Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu 195 200 205

Pro Trp Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro

210 215 220 <210> 35 <211> 20 <212> DNA <213> Homo sapiens <400> 35 20 gagccggcat catggattcc <210> 36 <211> 20 <212> DNA <213> Homo sapiens <400> 36 20 gctgcctgca caatggaggt <210> 37 <211> 1456 <212> DNA <213> Homo sapiens <400> 37 cgacgcagag cagcgccctg gccgggccaa gcaggagccg gcatcatgga ttccttcaag 60 gtagtgctgg aggggccagc accttggggc ttccggctgc aagggggcaa ggacttcaat 120 gtgcccctct ccatttcccg gctcactcct gggggcaaag cggcgcaggc cggagtggcc 180 gtgggtgact gggtgctgag catcgatggc gagaatgcgg gtagcctcac acacatcgaa 240 getcagaaca agateeggge etgeggggag egeetcagee tgggeetcag eagggeecag 300 ccggttcaga gcaaaccgca gaaggtgcag acccctgaca aacagccgct ccgaccgctg 360 gtcccagatg ccagcaagca gcggctgatg gagaacacag aggactggcg gccgcggccg 420 gggacaggcc agtcgcgttc cttccgcatc cttgcccacc tcacaggcac cgagttcatg 480 caagacccgg atgaggagca cctgaagaaa tcaagccagg tgcccaggac agaagcccca 540 geoccageet catetacace ecaggageee tggeetggee etacegeeee cagecetace 600 ageegeeege cetgggetgt ggaceetgeg tttgeegage getatgeece ggacaaaaeg 660 agcacagtge tgaceeggea cageeageeg geeaegeeca egeegetgea gageegeaee 720 tccattgtgc aggcagctgc cggaggggtg ccaggagggg gcagcaacaa cggcaagact 780 cccgtgtgtc accagtgcca caaggtcatc cggggccgct acctggtggc gttgggccac 840 gcgtaccacc cggaggagtt tgtgtgtagc cagtgtggga aggtcctgga agagggtggc 900 ttetttgagg agaagggege catettetge ceaceatget atgacgtgeg etatgeacee 960 aqctqtqcca agtgcaagaa gaagattaca ggcgagatca tgcacgccct gaagatgacc 1020 tggcacgtgc actgctttac ctgtgctgcc tgcaagacgc ccatccggaa cagggccttc 1080 tacatggagg agggcgtgcc ctattgcgag cgagactatg agaagatgtt tggcacgaaa 1140 tgccatggct gtgacttcaa gatcgacgct ggggaccgct tcctggaggc cctgggcttc 1200 agctqqcatq acacctgctt cgtctgtgcg atatgtcaga tcaacctgga aggaaagacc 1260 ttctactcca agaaggacag gcctctctgc aagagccatg ccttctctca tgtgtgagcc 1320 cettetgece acagetgeeg eggtggeece tageetgagg ggeetggagt egtggeeetg 1380 cattletggg tagggetgge aatggttgee ttaaccetgg eteetggeee gageetggge 1440 1456 tecegggeee tgeeca

<210> 38 <211> 423

<212> PRT

<213> Homo sapiens

<400> 38

Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe
1 5 10 15

Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg 20 25 30

Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp 35 40 45

Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
50 55 60

Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly
65 70 75 80

Leu Ser Arg Ala Gln Pro Val Gln Asn Lys Pro Gln Lys Val Gln Thr 85 90 95

Pro Asp-Lys-Gln-Pro Leu Arg-Pro Leu Val Pro Asp Ala Ser Lys Gln
100 105 110

Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly Thr Gly
115 120 125

Gln Ser Arg Ser Phe Arg Ile Leu Ala His Leu Thr Gly Thr Glu Phe 130 135 140

Met Gln Asp Pro Asp Glu Glu His Leu Lys Lys Ser Ser Gln Val Pro 145 150 155 160

Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu Pro Trp
165 170 175

Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro Trp Ala Val 180 185 190

Asp Pro Ala Phe Ala Glu Arg Tyr Ala Pro Asp Lys Thr Ser Thr Val 195 200 205

Leu Thr Arg His Ser Gln Pro Ala Thr Pro Thr Pro Leu Gln Ser Arg 210 215 220

Thr Ser Ile Val Gln Ala Ala Ala Gly Gly Val Pro Gly Gly Gly Ser 225 230 235 240

Asn Asn Gly Lys Thr Pro Val Cys His Gln Cys His Gln Val Ile Arg
245 250 255

Ala Arg Tyr Leu Val Ala Leu Gly His Ala Tyr His Pro Glu Glu Phe 260 265 270

Val Cys Ser Gln Cys Gly Lys Val Leu Glu Glu Gly Gly Phe Phe Glu

275 280 285

Glu Lys Gly Ala Ile Phe Cys Pro Pro Cys Tyr Asp Val Arg Tyr Ala 290 295 300

Pro Ser Cys Ala Lys Cys Lys Lys Lys Ile Thr Gly Glu Ile Met His 305 310 315 320

Ala Leu Lys Met Thr Trp His Val Leu Cys Phe Thr Cys Ala Ala Cys
325 330 335

Lys Thr Pro Ile Arg Asn Arg Ala Phe Tyr Met Glu Glu Gly Val Pro

Tyr Cys Glu Arg Asp Tyr Glu Lys Met Phe Gly Thr Lys Cys Gln Trp 355 360 365

Cys Asp Phe Lys Ile Asp Ala Gly Asp Arg Phe Leu Glu Ala Leu Gly 370 375 380

Phe Ser Trp His Asp Thr Cys Phe Val Cys Ala Ile Cys Gln Ile Asn 385 390 395 400

Leu Glu Gly Lys Thr Phe Tyr Ser Lys Lys Asp Arg Pro Leu Cys Lys
405 410 415

Ser His Ala Phe Ser His Val

<210> 39

<211> 1575

<212> DNA

<213> Homo sapiens

<400> 39

cgacgcagag cagcgccctg gccgggccaa gcaggagccg gcatcatgga ttccttcaag 60 gtagtgctgg aggggccagc accttggggc ttccggctgc aagggggcaa ggacttcaat 120 gtgccctct ccatttcccg gctcactcct gggggcaaag cggcgcaggc cggagtggcc 180 gtgggtgact gggtgctgag catcgatggc gagaatgcgg gtagcctcac acacatcgaa 240 qctcagaaca agatccgggc ctgcggggag cgcctcagcc tgggcctcag cagggcccag 300 ceggtteaga geaaacegea gaaggeetee geeceegeeg eggaceetee geggtacace 360 tttgcaccca gcgtctccct caacaagacg gcccggcct ttggggcgcc cccgcccgct 420 gacagegeee egeaacagaa tgggtgeaga eeeetgacaa acageegete egacegetgg 480 teccagatge cageaageag eggetgatgg agaacacaga ggaetggegg eegeggeegg 540 qqacaqqcca gtcqcqttcc ttccqcatcc ttqcccacct cacaqqcacc gagttcatqc 600 aagacccgga tgaggagcac ctgaagaaat caagccaggt gcccaggaca gaagccccag 660 ccccagcctc atctacaccc caggagccct ggcctggccc taccgccccc agccctacca 720 geogeoegee etgggetgtg gaccetgegt ttgeegageg etatgeeceg gacaaaacga 780 gcacagtgct gacceggcac agceageegg ceaegeecac geegetgeag ageegeacet 840 ccattqtqca gqcaqctqcc ggaggggtqc caggaggggg cagcaacaac ggcaagactc 900 ccqtgtgtca ccagtgccac aaggtcatcc ggggccgcta cctggtggcg ttgggccacg 960 cgtaccaccc ggaggagttt gtgtgtagcc agtgtgggaa ggtcctggaa gagggtggct 1020 tetttgagga gaagggegee atettetgee caccatgeta tgacgtgege tatgcaccca 1080 qctqtqccaa qtqcaaqaag aagattacag gcgagatcat gcacgccctg aagatgacct 1140 qqcacqtqca ctqctttacc tqtqctgcct gcaagacqcc catccggaac agggccttct 1200

acatggagga gggcgtgccc tattgcgagc gagactatga gaagatgttt ggcacgaaat 1260 gccatggctg tgacttcaag atcgacgctg gggaccgctt cctggaggcc ctgggcttca 1320 qctggcatga cacctgcttc gtctgtgcga tatgtcagat caacctggaa ggaaagacct 1380 tctactccaa gaaggacagg cctctctgca agagccatgc cttctctcat gtgtgagccc 1440 cttctgccca cagctgccgc ggtggcccct agcctgaggg gcctggagtc gtggccctgc 1500 atttctgggt agggctggca atggttgcct taaccetggc teetggeeeg agectggget 1560 cccgggccct gccca <210> 40 <211> 153 <212> PRT <213> Homo sapiens <400> 40 Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp 40 45 Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly 75 Leu Ser Arg Ala Gln Pro Val Gln Ser Lys Pro Gln Lys Ala Ser Ala 85 Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Val Ser Leu 105 Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala Pro Gln Gln Asn Gly Cys Arg Pro Leu Thr Asn Ser Arg Ser Asp Arg .135 Trp Ser Gln Met Pro Ala Ser Ser Gly 145 150 <210> 41 <211> 24740 <212> DNA <213> Homo sapiens <400> 41 nnnnntgta ttttatcata ttttaaaaat caaaaaacaa aaggcagttg aggttaggca 60 tggaggttcg tgcctgtaat cccagcactt tgggaagccg aagcacgtgg atcacctgag 120

gtcaggagtt cgagaccagc ctgcccaata tggtaaaacc ctgtctctac taaaaataca 180 aaaaattagc caggcatggt ggtgggcacc tgtaatccca gctacttggg agactgaggc 240

aggagaatca	cttaaacccg	ggaggcgggc	tgggcgcggt	ggctcatgcc	tgtaatccca	300
					catcctggct	
					cctggtggcg	
					aacctgggag	
					caagagtgag	
					gccgagatct	
					aaaggcagtt	
					gttgctatgg	
					tgctgaaggc	
					aggtggctgt	
ggtgagatgg	aggaagacac	ctggggtact	gctctaggct	gtcagacata	cagaagctgg	900
cccagccaag	cccaggggct	gcaagggaca	tccttttgtg	tccccagtga	tctgcagctc	960
tcagacaccc	tcaagcacag	tgcctcttgc	ccagcccagc	actctcagtg	gggagccagg	1020
tgggagaaca	ggctcggaag	gggacctagg	cttatgcagc	gagccgggca	aagctggaac	1080
tggagcccag	gcccctggat	gccccctggc	ttgtggagtt	ctgggatact	gaggggaggg	1140
gacagggcat	gggagtgcgg	tgctctcacc	tttgacttga	actcattccc	caggggacag	1200
gggaggcctc	ctcaggatcc	acagatgccc	agtctcccaa	gaggggcctg	gtccccatgg	1260
aggaaaactc	catctactcc	tcctggcagg	aaggtaagtt	ggaggacgtg	caagggcagc	1320
					cctggccacc	
					cccgctgcca	
gctgaagggg	ccggccctgc	tggtgctggg	cccagacgcc	atccagctga	gggaggccaa	1500
					ctccgacaag	
					tgctaatccc	
					gctgccactc	
					gggctgtggc	
					gccctgcccc	
					ggagatgcca	
					accccagagc	
					cgcttcagtg	
					gtgtgtgctg	
					cggcagccgc	
					cggcccggcc	
					ggtggagggc	
					gctgagtcgt	
					agtggtggcc	
					cattgggccc	
					gcctgtggag	
					taataaagct	
					cattgcggcc	
					cctgccaccc	
					tccatgttgc	
					tacttgtcag	
					ctgggaggag	
					aaaacacttg	
					gtcagtctcc	
					gagagtttgc	
					agccttggtc	
					tccatgctgc	
					acgcaagcgt	
					gctgcacaca	
					tctctcctgt	
					gcggccagta	
					gaggcctctg	
					gtttcatctt	
					ctgcacacag	
					aacagtgcaa	
		_				

ggcaggcagt	ccctgccaca	agagaccctg	ttccctagta	gggcagcttt	tctcctcccc	3660
agaacctcct	gtctatcccc	acccaatgtc	tecteacagg	catattqqqq	aaacaggtca	3720
ggctctccca	ccgtatctgc	aagtgtactg	gcatccatct	gtcttcttcc	tacccctaca	3780
gtagaaacag	tatctatccc	cagctatact	ctgatcccgg	ctcctttcac	ctcagagctt	3840
ggaaaattga	actatececa	ctctctcctq	coccattca	tcctaccage	agcttttcca	3900
					ctcacattcc	
tetteggea	cctctatttc	tetatteete	ttcacagcaa	aaactottca	aaagagttgt	4020
taattacttt	catttccact	ttctcacccc	cattetetee	tcaattaact	ctccttcatc	4080
accatastac	cattatataa	cttttattag	agtcaccaac	cttattctcc	aaaacaaaag	4140
gaagaagaa	tttgacttct	carcarcact	carctetaat	tettgaaaca	ccccgttac	4200
ttactattcc	tectacetea	taacaatctc	cttcccagc	tctactacta	ccttctctga	4260
					acaaagaatc	
geeeeeea	gggtcctagg	atcastacst	attetatasa	tratctacca	agaaaataaa	4380
teetgaaage	cigiaaaaaa	gcccacgcac	accececacac	aggggggtgtg	atttcatcag	4440
aaattttaaa	aaccaaacgc	ceatgeetgg	geceacatge	aggggccccg	attatataca	4500
					gtagtatgca	
gcctaggctg	ggaaccactg	geeteageaa	tacatacaca	ccccaggige	cacagaccet	4520
ctaggtgcta	atgaccccga	aggicigici	cagigeaca	ecceeccig	agctccagat	4620
ttaggaatcc	cactgcacac	gagacatetg	gatgtggaaa	agacatetee	agatcccatg	4740
ggtgaaaggg	ggttggggga	atggagactc	gtgttettee	aggatgtgtg	tggacacaga	4740
atgcaaagcc	tggagggatg	ctagagccat	agggaggaag	attteggete	acttattcat	4800
gcaagcactt	cctgatgggt	aaggtcttag	agcaagctga	ggccaagagg	cgggcagtcg	4860
aggtgctgct	gcaggcaccc	ccactcccta	cagtggcaag	cccaagccca	gcccttggca	4920
gctcaaatcc	caggacacgc	tgaaggtcac	-ccagagagtc	aggggcatgg	-ctagaaccag-	4980
aacccaggac	tctggggacc	cagcatggca	teettteett	cattacaaat	ctgagctgct	5040
ttgtttccta	gggatttctg	tgatattcca	aggggactgt	gggaaagaaa	gtccttggaa	5100
accaccagga	cgctagaggc	ctggcctgga	gcctcaggag	tctcggccac	cagagggcgc	5160
tgggtccttg	tccaggtcca	gttgctacgc	aggggctgcc	tgtgctggga	ggctccccag	5220
gggacacaga	ccagagcctt	gcaccagccc	aaggaatggg	agcctggggt	cctctctgct	5280
ggaggactgc	caggaccccc	aggctgccgc	ctcttccttt	gctcatttgc	tgtttcactt	5340
tgtcaatcct	tectttette	gtgtgttcat	tcacatccac	tgtgtgctgg	ccctggggaa	5400
atgttagata	agacacatta	gctgtgtgtc	ttcattgtcc	taacaaagaa	cacaccctgg	5460
aaagagcacc	gcagagagtc	cccattcccc	catctccctc	cacacatgga	atctggagat	5520
gccttttcca	catccagatg	tctctggtgc	tgtgggattc	ttaaataaac	aaacatttca	5580
tacagaatgt	gagatgatgg	agatgctatg	gggaaaagta	aagcagaggg	agggcctagt	5640
gtgtgatgcg	ggtgaggcat	ccagggattg	ctgtttcagc	tgtgatcagg	aaaggccctg	5700
ggaggaggcc	acatctgagc	agagacctaa	ataaagttgg	aaacctgttg	ctgagatatc	5760
tggagaagtg	tttcaagggc	cgggcaccgg	gcatggtggc	tcacgcctgt	aatcccagca	5820
ctttgggagg	ccaaggcagg	tggatcgctg	gaggtcagga	gtttgagagc	agcctgacca	5880
acatggagaa	accccatctc	tactaaacat	ataaaaatta	tccgggcatg	gtggttcatg	5940
cctgtagtcc	cagctactcg	ggaggttgag	gcaggagaat	cacttgaacg	tgggaggcag	6000
					agcgagactc	
cgtctcaaaa	aaaaaaaaga	aaagaaaaaa	gaaaaaaaaa	gaaaagtgtt	tcaagcaggg	6120
gaactggcaa	gtggagaggc	cctgaggcag	aaatatgctt	ggcctgctgg	aggaaatgtg	6180
					gacccagttt	
attcatattc	tgtaggtctt	aaggacttca	gttttatttt	gagtgcaata	tgagcccact	6300
ggaatgctaa	aagctgagag	tgacatggtg	ctgtgattct	ggctttaaaa	atatcacttt	6360
					cgttaggaga	
					agcagcggtt	
					agagaaagga	
					ttgaagtgag	
gaaggagtta	ggaaggtggg	aaaggcacaa	gtttgaattq	ggccatgatg	gtctgagata	6660
cctagtacag	tggttcccca	acctttttgg	cagaagggac	cgctttcatg	gaagacaatt	6720
tttccacaga	ctagaaataa	qqtqqqqatq	gtttcagggt	ggttcgagtq	cagtacattt	6780
atcattagac	tettttttt	tttttttt	tgagatagag	tetegeteta	tcacccacac	6840
					tcaagtcatt	
ctcctacctc	agcctctcaa	qtaqctqqqa	ttataggcat	atgcqccacc	acgcccagct	6960
	J	J. J. J.J.	33			

				ggccaggatg		
				gtgctggggt		
				ggaatgagca		
				gagtctaatg		
				gcagcccagt		
ccacaaacgg	ggagttgggg	acccctgatc	tagtaaacat	ctaggcaggg	ttttggataa	7320
tggagttaga	gttcctgggg	agaggtcagg	ctggccatga	aacatgggat	gcctttgcat	7380
ataggtggtg	ttgaaagcca	caggacagta	cggggtctca	gggggtgagc	ataaagagag	7440
gcgacatcag	atggccaagg	ccagaggcag	aggaggatgg	gaaggagggg	ccagtggggc	7500
agggggaagc	tgtgaagcca	gggaaaaagg	gtgtttcgcg	gaaaaggatc	aacctggacc	7560
				acggattcca		
				gggacgggtg		
				agtggagacg		
				aagggtagta		
				cacggcggtt		
				cagcgactct		
				cgccccttcc		
				cctgggctga		
				ccaaggtgga		
				catccggcgt		
				tttaaacgcc		
				cagtccccga		
				atcgtcccat		
				ctggtttgaa		
				ggcagcgctt		
				gcagggcccg		
				aggacggaag		
				gacgagggcg		
				aaaaacaaaa		
				agcagcgccc		
aagcaggtat	cgacgaccgc	gcggggcgtc	ttgggctgga	ccaggcgggc	gcccggggcc	8820
tgctgaggac	cacaaagggc	actgggggtc	gtggtccagg	ctgtgcttcc	tcccgctggc	8880
cctggcccct	gcctccgccc	ccgcccccgc	cttcctgccg	ctaagccggc	tgcggcgggg	8940
ccgattggcg	cctgccggct	tcctgcgccg	gggccagtct	aatgcatggg	gcccgggcgg	9000
gggactaagg	ggaaactgag	tcacgtcggt	gtgggagcag	ttctgtgtgg	gaggcaccac	9060
ccccactgg	gctcggggaa	ggatccccct	ccaagctatg	cttgagggtc	ccagccccca	9120
tctgtctcca	caggggccgc	accccactcc	cgccttcccc	ttcttcagca	cccaggggtc	9180
				ctaggaggat		
				ccctcccctc		
				tctccgcaca		
				gctgccggga		
		•		gaggetecag		
				ttgcaaagaa		
				ggggctgctc		
				aaaatacctg		
				tgtccaaggc		
				ccactcacct		
				ctggaggggc		
				ctctccattt		
				gcttaggtct		
				gccaggagat		
				gaggtggata		
				ttctcccaag		
				gagcctaaca		
				ggcaggacac		
ggatttcact	tcagcctcac	rgccgcccag	ggaatagcaa	tcctcatccc	gtttttccag	10320

						ggtgaagcag	
						gtctctcccc	
						tttgacccca	
	agcagaagtg	taggacaggg	ctccattcta	ggggcttaac	ttcagcttcc	aagagcctgc	10560
	cctggtgtgg	gtggagctgg	aggctggctc	ctccctgtag	cagggggatt	gccttataag	10620
						agagctgaaa	
						tggtcaccag	
						cccagagtgg	
						ggggtgggga	
						tgggcaggca	
						tcgtggcctg	
						gggtgggcag	
						gccccgccag	
						cctcacacag	
						tgtctgggga	
						gttgggggca	
						ctagctgatc	
						tctgagttag	
						gcaagggctg	
						gggctgctca	
						tccttggttc	
						ccacccgccc	
						catcctctgt	
						ccctctccag	
						tcccaggagc	
						accatgacct	
						ggggttgatt	
						ggctgacccc	
						agggtcagag	
	acacttcctg	gctctgcctc	ttactagctg	ggtgacttga	ggcaagttgt	ttaacctctc	12120
	tgtgtacatt	tgcaagtgca	aaatgggtaa	aatcccagat	tactccacaa	ggttgttgga	12180
	agattcagtg	tcaatatgta	gcatagttgg	tgctcaataa	actgaagcaa	gtcttcttat	12240
	ttagcgagtg	aggaaggggc	cgccgagctc	tcttagcctt	ctgacctcct	acgcaagcaa	12300
						ctgtgcctgg	
						gcctcagctc	
						ggacccccc	
•						ctacttgatg	
						cagtcaagct	
						agggtcaccc	
						ccctctacct	
						gaactctcac	
						aaagtcattc	
						ctctactcat	
						aggggaaatt	
						gctgtcaact	
						gttcctggct	
						ggggtgagag	
						gccagagggc	
						aggcaatggt	
						atagcaattt	
						gggaagetea	
						tgactgggaa	
						agcagctgcc	
						ttgggtgggg	
						gtcactttgc	
	recreteree	acageteact	cergggggca	aaycygcgca	ggeeggageg	gccgtgggtg	T3080

actgggtgct	gagcatcgat	ggcgagaatg	cgggtagcct	cacacacatc	gaagctcaga	13740
acaagatccg	ggcctgcggg	gagcgcctca	gcctgggcct	cagcaggtat	gcgggtggac	13800
atggatgggt	gegeeegege	tggcagtggg	gatccctgcg	gcccggcccg	ctgtcacgct	13860
ttccttctcc	tccagggccc	agccggttca	gagcaaaccg	cagaaggtac	gaggetggee	13920
gggacatccg	ggcggtgggc	ggtgtgggct	tggacggcca	ggcctgctcg	ccctcctggc	13980
acattctcgg	taccccaatc	cctggccggg	agtggagggc	agaaaccgga	gctaaggcgg	14040
	ctggagttga					
	cgcctgtctg					
	cagccctcgg					
	gtcgtcctct					
tggggggctg	gagctgccgc	tgtcctctgc	tgcaggcgcc	ccacttccac	ccaggccccc	14340
accttaccct	geeegeeege	cctgcccggc	tgtgtctctg	cccaggcctc	cgccccgcc	14400
gcggaccctc	cgcggtacac	ctttgcaccc	agcgtctccc	tcaacaagac	ggcccggcct	14460
ttgggcgccc	ccgcccgctg	acagcgcccc	gcagcagaat	gggtacgtcg	gcccctgccc	14520
gcccgcgccc	acgccatcag	gcccactgtg	gccccacgcc	cgctgcccgc	tgctgctcag	14580
tctgtgctgc	gccccagccc	ggcggaaccg	tgcggcacgc	cccctggcgg	ccggggtggg	14640
gctgcaggca	cagggcccct	cccgaggctg	tggcgccttg	cagggcaccg	cctggggagg	14700
ggtctctgaa	tgacgccgcg	cccctgctg	gcggctgggg	gttgggttgt	ggtgtcgggc	14760
cagctgagcc	ccagacactc	agtgccgcct	tgtccccggc	tgttctgacc	cctccccgtc	14820
tttcttcctc	tcctgtgtct	gtccctttgt	ccctttatct	gtctgtctgt	cttatttcct	14880
tcacaggtgc	agacccctga	caagtcagtg	agcccccctc	tgcctgtgcc	tttcttcttc	14940
cttttggcac	tetgggtgge	ggcccctccc	caccctggct	gccctcctct	ccacttcgcc	15000
ctcctgtcct	ctcacctacc	-cgcccagcag	-ggctcctggc	ctcaccctta	-cccactccct	15060
cccatcactg	taacccaaac	ccacatgcac	caaatcctgg	gaggggctgc	cccaccgcc	15120
cacccccagt	gtggggttct	gagccacacc	ctccccacag	acagccgctc	cgaccgctgg	15180
teccagatge	cagcaagcag	cggctgatgg	agaacacaga	ggactggcgg	ccgcggccgg	15240
ggacaggcca	gtcgcgttcc	ttccgcatcc	ttgcccacct	cacaggcacc	gagttcagta	15300
agtgccagcc	cagggcaggg	ggtactttcc	tegeececag	cccaggcgtg	atccctgacc	15360
ctgtgtcttt	tttggtcaat	gcctgcctct	gctctctcag	tgcaagaccc	ggatgaggag	15420
cacctgaaga	aatcaaggta	cagggacggg	caccagcccc	tctcccacct	cctgcctctt	15480
	tactgccctg					
	ttccttcctt					
cccttcctcc	ttccttcctt	ctttcatttc	ttccctccct	ccttccttcc	ctcctccctc	15660
cctgcctccc	ttccatctct	ccttccttcc	acttcttcct	ccctctctct	ctgcccctca	15720
gggaaaagta	tgtcctggag	ctgcagagcc	cacgctacac	ccgcctccgg	gactggcacc	15780
accagcgctc	tgcccacgtg	ctcaacgtgc	agtcgtagcc	cggccctctc	cagccggctg	15840
ccctctctgc	ctccctctt	ctgttcctcc	tgcccagggc	acccccttag	tgcctccagc	15900
ttctgcctac	ctcacccccc	ctttcgtgcc	cctggcctga	geeteetget	ggcctggccc	15960
tggccgccca	cctgggttca	tctgacactg	ccttccctct	ttgccctgtg	gtactgctgt	16020
ctgccaggtc	tgtgctgcct	tgggcatgga	ataaacattc	tcagccctgc	ttgctctgcc	16080
tgtcttctat	ctttgtggac	ctggtttgca	tttggggtgt	gggggtgttt	cgtggttcgg	16140
actgtttggg	ccctgccgtc	cttgttttca	gtgggagggg	gtacctggca	aaggggccct	16200
gccctgccat	cacagatggc	ttcctggcat	gaggggagcc	ccaggagctg	cctcagaagc	16260
gggagccctg	cctcgtctcc	cagctagaga	ccgcacacca	gctaactgga	cattgctagg	16320
agaagctgcc	cttcccatcc	ctaccccagt	gggacctgga	atccaactcg	gcagtttcca	16380
cgcccccagt	catctcccgt	ggggccagca	ggacccaggt	tggggggtgg	ggccatgtca	16440
ggaagctcag	ccatgcaggg	ccttgaatgg	cagatcttgc	agccaggtgc	ccaggacaga	16500
agccccagcc	ccagcctcat	ctacacccca	ggagccctgg	cctggtgaga	gggagtgggc	16560
tegggeetgg	gcaagggtgg	gcagcctcca	ggggcatggg	ggtggtgggc	ttctctcagc	16620
	tccacccccg					
ccggcaggcc	ctaccgcccc	cagccctacc	agccgcccgc	cctgggctgt	ggaccctgcg	16740
tttgccgagc	gctatgcccc	ggacaaaacg	agcacagtgc	tgacccggca	cagccagccg	16800
gccacgccca	cgccgctgca	gagccgcacc	tccattgtgc	aggcagctgc	cggaggggtg	16860
ccaggagggg	gcagcaacaa	cggcaagact	cccgtgtgtc	accagtgcca	caaggtcatc	16920
	cctgttcctg					
gtctgcccac	ctgtcttgcc	tcagctgcga	ctggggggaa	taaggattca	gttctcagct	17040

```
ggagtaggag tagggacctg ggctgggtcc teccattett aateccaege tacetaecce 17100
agcccaccca caacaactgc tagcagcatc tgccgtggcg aaatagccga agggccaacc 17160
ataggetgaa getgeaccee tacetttget getetetggg caaagagggg cetgeeceet 17220
eccagegegt etgeceetee etcetgetet etgeteteet etgeteteag ageatacagg 17280
cctggagcca ctccctctgt gcactgcccc gtggggccaa gcagcatcaa acacccccca 17340
tgtcagctct ttttttttt tttttgaaac agagtctcac tgtgttgccc acgttggcgt 17460
gcagtggcgc gatctcggct cactgcaacc tctgcctcct gggttcaaga gattctcctg 17520
cctcagcctc ctgagtagct gggattacag gcaccegcca ccatgcctgg ctaattttgt 17580
atttttagta gagacggggt tttaccatgt tggccaggct ggtctcaaac tcctcacctc 17640
aggtgatete aggeetgeet tggeeteeca aagtgetggg actacaggtg tgagecactg 17700
tgcccagccg actctatcag ctcttgccag gtagaacagg caggccagca ggacagggca 17760
getecagggt ttgcccaggg geggeteage ttttatgagg etecagtegt eagecettee 17820
teceggggte etecetgete taaagetgee teteetgtea ceageagtte agtgtggegg 17880
actggetetg taagetteat ggetgecacg gteactteec aageetgtet tetateetat 17940
gtggaaaatg gggagaatga actgtccctc ccaaggcctc ctggtgggtg gtcagtcaac 18000
ctgaaggggg ccaagacccc cacctctctg cgtgtgctcc ctctgaccgc tctcgcctcc 18060
ctgcaggggc cgctacctgg tggcgctggg ccacgcgtac cacccggagg agtttgtgtg 18120
tagccagtgt gggaaggtcc tggaagaggg tggcttcttt gaggagaagg gcgccatctt 18180
ctgcccacca tgctatgacg tgcgctatgc acccagctgt gccaagtgca agaagaagat 18240
tacaggcgtg agtagggctg gctggcgggg aggtggtccc aagcctgtca gtgggaacga 18300
gggctgctgg gaaacccaca gtccaggtct ctccccgagt gagcctccgg gtccttacca 18360
gcgtaataaa-tgggctgctg-tactggcctc-accctgcatt-agtcaggatg-ctcttaacaa-18420
atgaccatgt tcctgctcag aaaccgccca aggctgcaaa gagcaggagg accaagccag 18480
gagaageeet gggeeeteet gaeteeeaet ttgggetete eetgeeetgg tgaaatgaea 18540
gaacggccaa cttgacacgc tgaagctgct ctgtctcatg cgtcctcctc atttctggat 18600
ccagagccag ggctgccagg agtagccaga gagctctgtg tggtgatgtt catattagtg 18660
aggtttacct tgaccacgag cagtgggaaa ctcaaaataa tggtggctta tttctcatct 18720
aaaaacatcc cggggtgggt ggtctgggac tgatctggtg gacccaggct ccgccttgtt 18780
gettgaetgt tggeageace tgettaetta ceacteatgg tgeaagatga caetteagee 18840
teegecaaaa tgeteacett eeagecagea ggaagtegga aggagaagaa aggggacaga 18900
geeceatgge gtecateett agaggatget geeacetgaa eetetgettt cateetgttg 18960
gtcagaaccc agtcacatga ccacacccag tggcaacgga ggctgggaaa tatagtcttt 19020
attttgggca cccatgtgtc cagcaaaact gggggttcca tcagtcggca agaacgggag 19080
agtggccgat gcagtggctg atgcttgtat cccagcactt tgggaggtcg aggtgggcag 19140
atcacctgag gtcaggagtt caagaccagc ctggccaata tggtgaaacc ctgtctctac 19200
taaaaataaa aaaattagct gggtgtgctg gcgcacctgt agtcccagct acttgggagg 19260
ctgaggcagg agaatcgctt gatcttgaga ggtggaggtt gcagtgagcc aagattgtgc 19320
aggeaeggtg geteaeacet gtaateceag eactttggga ggeegagatg ggeggateae 19440
gaggtcagga gattgagacc atcctggcta acacggtgaa accccatctc tactaaaaat 19500
acaaaaaaat tggccgggca tggtggagta gtcccagcta ctcgggaggc tgaggcagga 19560
gaatggcgtg aacctgggag gcagagcttg cagtgagccg agatcgcgcc actgcactcc 19620
agcctgggca acagagcgag actcttgtct caaaaagaaa aaaagaaaga gaaatctgcc 19680
teccageett gggeteetge eetaceagee cacaceeetg gtagageete eteteceaee 19740
ageteaaage ceaagtteet teaetgtgae ettgtetget cetetaaaac aggeaacace 19800
agacagtgag aagagccagc cagacatggg cagaaaacct atttctgtga tctactggct 19860
gtgtgageag gggetagttg etetetetgg geeteactga agagaagggt ggeactatge 19920
tagggccggc acggttgcaa ggtagatgta agatggggta caggtgttgt ggagggcaga 19980
aatgcaccat ccgaaggcta catgtccccc acacttatgt cttgcttggc ccacactgtt 20040
tcattttaaa atcagtagca aacaatttaa aaaatcagaa gatttgcctg catgatgcag 20100
tggctcatgc ctgtaatccc agcactttgg gaggccaagg tgggaggatt gcttgagccc 20160
attagaaaat tagccaagtg tggtggcatg cacctgtggt cccagctact tgggaggcag 20280
agggaaagtg agatotootg otttttattt otttatgtat aatgataggg tottgototg 20340
ttgcccaggc tggagtgcag tggcatgatc actgctcact gcagccttga tctcctgggc 20400
```

	tcagaggatc	ctcccacctc	agcctcccaa	atagctagga	ctagaggtgc	ccaccagcat	20460
	gctcagcaga	tttttaaatc	tttttgtaga	gatgaggttt	tgctatgttg	cccaggctgg	20520
	tctcgaactc	ctggcctcga	gcgatcctcc	caccttggcc	tcccaaagca	ctgggattac	20580
	agacgtgagc	cactgcgccc	agcagatttc	tctttaacac	ctagatttca	gcctgagcca	20640
	ggcaggcatt	cctgaatgaa	ccagtagtac	tgctcccaga	agaagaggtc	ctcctccgtg	20700
	tgacacagtc	cccacttggc	ccttgcaggg	attggatctg	ggatccctgg	atttaaactc	20760
	agggccatcc	tcataacagc	ctcacaaggc	tgggattagc	ttcccagttc	acaagggaag	20820
	aaaccaagac	ttgagaaggt	caaggtctgg	ccagacccac	acatcttgga	ccctcatacc	20880
	gcctcgaggc	cccatgctgc	cctctgcctg	ctccagatgt	gaatactgct	ggccctggct	20940
	ggccccggct	ggccccgagg	gtcctaggga	tgaacagccc	agcccaggga	gagctcagcc	21000
	ccttgtgcct	ctgccccttc	ccacctcctg	cggaggccag	tcgactcacc	cacaaagggc	21060
	caggcactgt	ggggatagat	cagctaacaa	aacagttgat	gcttcctgcc	cttctgggcc	21120
	ttacattttg	gctggaagaa	gagggagag	gcagactgta	agcaataagc	gcaataagta	21180
	ggttgcctgg	aagtaatgtt	agatcacgtt	acggaaaaca	ggaaagagca	gagcgacaag	21240
	tgctggggtg	cgtggtgcag	ggaaggcagc	tggctgctgc	tggtgtggtc	agagtgggcc	21300
	ctcatggaga	agactgcatt	cgagcagaaa	cttgaagggg	gtgaggggtg	agcctagaga	21360
	tatctqqqqc	agagcagtcc	aggcagaggg	gacagccggt	gtcaagccca	ggacaggagt	21420
	gtgcctggtg	tgccagtttc	aggcaagagg	ccagtgtgca	gaggcaaggt	gagaacgcaa	21480
	qqqaqaqcaq	tggcggagac	gggtgggaac	gaggtcagac	ctgctggcct	ccagcctctg	21540
	catqqqqctt	ggctcttgct	gggagcaatg	ggaagcagta	cacagtttca	tgcaggggga	21600
	gaaggcctgt	cttgggttgc	aggggcacgc	tgtggcagct	gggatcagag	agaggagctt	21660
	gtaggccagt	tgttatgtgg	tcccacgggc	cagatggcca	tggcttacct	cacttcaggg	21720
_	aggctgtgag=	aagcactcag	aatetggatg	tgccttgggg	gtgggcccca-	ctggatttcc	21780
	tqqtqqacct	ggtgtggggt	gtgagaggag	ggtgtgtttg	gctgcagcag	acaggagaat	21840
	ggagttgcca	teegegtgat	ggggatggct	gtgggaggag	aggtttgggg	tgagggaatc	21900
	aggaactgag	tgctggacat	ggcaagtctg	aaggcgcagt	ggtcgtccac	tcagagacct	21960
	tagaattaga	gatggaggtg	tgggagtcct	gaacagttag	atgtagtgtt	taccgcgaga	22020
	aggaacaggg	cttqcqqcca	gccctcctgt	gttcccgtga	cccagggcag	ggcaggaggg	22080
	gcctgagcct	gccgagtgac	tgggacctcc	ttccaggaga	tcatgcacgc	cctgaagatg	22140
	acctggcacg	tgcactgctt	tacctgtgct	gcctgcaaga	cgcccatccg	gaacagggcc	22200
	ttctacatgg	aggagggcgt	gccctattgc	gagcgaggta	cccactggcc	agtgagggtg	22260
	aggagggatg	gtgcatgggg	caggcatgaa	tccaggtcct	ctttctctct	gccccattc	22320
	tcagactatg	agaagatgtt	tggcacgaaa	tgccatggct	gtgacttcaa	gatcgacgct	22380
	ggggaccgct	tcctggaggc	cctgggcttc	agctggcatg	acacctgctt	cgtctgtgcg	22440
	gtgagagccc	cgccctcga	actgagcccc	aageccaceg	gccctctgtt	cattccccag	22500
	gagatgcagg	agaagttggg	aaggggcctc	tcctgctgcc	cccaacccca	tgtgactggg	22560
	cctttqctqt	ccttagatat	gtcagatcaa	cctggaagga	aagaccttct	actccaagaa	22620
	ggacaggcct	ctctgcaaga	gccatgcctt	ctctcatgtg	tgagcccctt	ctgcccacag	22680
	ctgccgcggt	ggcccctagc	ctgaggggcc	tggagtcgtg	gccctgcatt	tctgggtagg	22740
	gctggcaatg	gttgccttaa	ccctggctcc	tggcccgagc	ctggggctcc	ctgggccctg	22800
	cccacccac	cttatcctcc	caccccactc	cctccaccac	cacagcacac	cgatgctggc	22860
	cacaccagcc	ccctttcacc	tccagtgcca	caataaacct	gtacccagct	gtgtcttgtg	22920
	tgcccttccc	ctgtgcatcc	ggaggggcag	aatttgaggc	acgtggcagg	gtggagagta	22980
	agatggtttt	cttgggctgg	ccatctgggt	ggtcctcgtg	atgcagacat	ggcgggctca	23040
						cacagacatg	
	aqqqqaqcca	ctggtctggc	ctggcttgga	ggttagagaa	gggtagttag	gaagggtagt	23160
	tagcatggtg	gctcatgcct	gtgatcccag	cactttggaa	ggccaaggtg	ggcagatcgc	23220
	ttgaggtcag	gagttcgaga	cctcatggcc	aacacggtga	aacagcgtct	ctagtaaaaa	23280
	tacaaaaatt	agccgagtgt	ggtggggcat	gcctgtaatc	ccagccactc	aggaggctga	23340
	ggcgggaaaa	tcacttgaac	ctgggaagtg	gaggttgcag	tgagctgaga	tcacaccact	23400
	qcqcqcqaqc	ctgggtggca	gatggcagag	cgagaccctg	cttcaaaaaa	aaaaaaaaa	23460
	aaaaaaaaaa	gaagggtagt	tgtagttggg	ggtggatctg	cagagatatg	gtgtggaaaa	23520
	cagcaatqqc	cacagcaaaq	tcctggaggg	gccagctgcc	gtccaaacag	aagaaggcag	23580
	ggctggagag	ggtagccctt	aggtcctggg	aagccacgag	tgccaggcag	tagagctggg	23640
	gctgtctctt	gaggttaggg	cagggcaagg	cacagcagag	tttgaaatag	gtttgtgttg	23700
	tattgcagaa	aagaggcccc	agaacactga	gggagtgcag	gagggaggct	gggaggagga	23760

```
gttgcagcag ggcctagggg cgggggccag gcaagggagg ggcagagagt aatatggcag 23820
agatgggacc cagtggcagg tccgggggat gagggatgga gagaaggaca ggagcgttgc 23880
caggeatetg geetatacca gacatgetea egetgtetee egegaacete etageaacet 23940
ttttaagaga caggatetee etaggttgee egggetggtt teaaaeteet gggeteaage 24060
aattetteet eettageee aaagtgetgg cattacaggt gtgagecace atgeetggee 24120
cacttatttt ctagatgagg cacagaaaga ttgggagact tgaccaaggt cacgctgtca 24180
ttgagccatg agccagacta gaatccaggc ctgaagctgg gtgcgctgtc ccaggactgg 24240
ctggcactga gtaccatttg ccagcgagca tetetetggg aagetgaett etgeeeggta 24300
cctggaggac tgtagacctt ggtggtggcg ccgtcactct ggggcttcct gcctcccact 24360
gatgcccgca ccaccctaga gggactgtca tetetectgt cccaagcctg gactggaaag 24420
actgaagaga agccttaagt aggccaggac agctcagtgt gccatggctg cccgtccttc 24480
aqtqqtccct qgcatgagga cctgcaacac atctgttagt cttctcaaca ggcccttggc 24540
ccggtcccct ttaagagacg agaagggctg ggcacggtga ctcacacctc taatcccagc 24600
actttggaag getgaggetg gagaaggget ceagettagg agtteaggae cageetggge 24660.
aacatggtga gaccctgttt tgttttgttt tttgttttt tgagatggag tcttgctctg 24720
                                                              24740
tcgcccaggc tggagtgcag
```

<210> 42

<211> 25

<212> DNA

<213> Rattus norvegicus

<400> 42

gcactacctt gaaggaatcc atggt

# INTERNATIONAL SEARCH REPORT

International application No. PCT/US00/11664

A. CLA	SSIFICATION OF SUBJECT MATTER				
	:Please See Extra Sheet.	14.4			
	:435/320.1, 325; 530/350, 388.1; 536/23.5; 514 to International Patent Classification (IPC) or to bot				
<del></del>	DS SEARCHED				
	locumentation searched (classification system follow	ed by classification symbols)			
i					
U.S. :	435/320.1, 325; 530/350, 388.1; 536/23.5; 514/	44			
Documenta	tion searched other than minimum documentation to th	ne extent that such documents are included	in the fields searched		
Flectronic o	data base consulted during the international search (	name of data base and, where practicable	e search terms used)		
	e Extra Sheet.		o, source: to the uses,		
c. Doc	UMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where a	ppropriate, of the relevant passages	Relevant to claim No.		
A	BODEN et al. LMP-1, a LIM-domain protein, mediates BMP-6 effects on bone formation. Endocrinology. 1998. Vol. 139, No. 12, pages 5125-5134, entire document.				
A	HOGAN et al. Bone morphogenetic proteins: multifunctional regulators of vertebrate development. Genes and Development. 1996. Vol. 10, pages 1580-1594, entire document.				
A	KINGSLEY, DM What do BMPs do in mammals? Clues from the mouse short-ear mutation. Trends Genet. January 1994. Vol. 10, No. 1, pages 16-21, entire document.				
·					
Further documents are listed in the continuation of Box C. See patent family annex.					
Special categories of cited documents:  A* document defining the general state of the art which is not considered  T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention					
to t	to be of particular relevance				
	cartier document published on or after the miernational filing date considered novel or cannot be considered to involve an inventive step				
cito	document which may throw doubts on priority claim(s) or which is when the document is taken alone cited to establish the publication date of another citation or other				
*O* doc	pocial reason (as specified)  Ye  document of particular relevance; the claimed invention cannot be considered to inventive as inventive step when the document is  locument referring to an oral disclosure, use, exhibition or other  Combined with one or more other such documents, such combination				
*P* doc	means being obvious to a person skilled in the art  document published prior to the international filing date but later than *& document member of the same patent family the priority date claimed				
	actual completion of the international search	Date of mailing of the international sea	arch report		
16 JULY	2000	29 AUG 2800			
Commission Box PCT	nailing address of the ISA/US ner of Patents and Trademarks , D.C. 20231	Authorized officer ANNE-MARIE BAKER	ellusfor		
Consimile M	~ (702) 205 2220	Telephone No. (703) 308-0196	$\nu$		

Form PCT/ISA/210 (second sheet) (July 1998)★

# INTERNATIONAL SEARCH REPORT

International application No. PCT/US00/11664

A. CLASSIFICATION OF SUBJECT MATTER: IPC (7):
A61K 48/00; C07H 21/04; C07K 14/00, 16/00; C12N 15/63, 15/85, 15/86
B. FIELDS SEARCHED  Electronic data bases consulted (Name of data base and where practicable terms used):
WEST Dialog (file: medicine) search terms: gene(w)transfer, LMP, bone, LIM, mineral?, human, osteogen?
Taran da se em esta de manera de la compansión de la compansión de la compansión de la compansión de la compan

Form PCT/ISA/210 (extra sheet) (July 1998)\*

# CORRECTED VERSION

#### (19) World Intellectual Property Organization International Bureau





(43) International Publication Date 9 November 2000 (09.11.2000)

**PCT** 

# (10) International Publication Number WO 00/66178 A1

- (51) International Patent Classification?: A61K 48/00, C07H 21/04, C07K 14/00, 16/00, C12N 15/63, 15/85, 15/86
- (21) International Application Number: PCT/US00/11664
- (22) International Filing Date: 28 April 2000 (28.04.2000)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/132,021

30 April 1999 (30.04.1999) US

- (71) Applicant (for all designated States except US): EMORY UNIVERSITY [US/US]; 1380 South Oxford Road, Atlanta, GA 30322 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): BODEN, Scott, D. [US/US]; 2842 Cravey Drive, N.E., Atlanta, GA 30345 (US). HAIR, Gregory, A. [US/US]; 1887 Chrysler Drive, N.E., Atlanta, GA 30345 (US).
- (74) Agents: GARRETT, Arthur, S. et al.; Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P., 1300 I Street, N.W., Washington, DC 20005-3315 (US).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published:

- With international search report.
- With (an) indication(s) in relation to deposited hiological material furnished under Rule 13his separately from the description.
- (48) Date of publication of this corrected version:

15 March 2001

(15) Information about Correction: see PCT Gazette No. 11/2001 of 15 March 2001, Section 11

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

#### . 7 (

# (54) Title: LIM MINERALIZATION PROTEIN SPLICE VARIANTS

(57) Abstract: The present invention is directed to isolated nucleic acid molecules that encode LIM mineralization protein, or LMP. The invention further provides vectors comprising splice variants of nucleotide sequences that encode LMP, as well as host cells comprising those vectors. Moreover, the present invention relates to methods of inducing bone formation by transfecting osteogenic precursor cells with an isolated nucleic acid molecule comprising a nucleotide sequence encoding splice variants of LIM mineralization protein. The transfection may occur ex vivo or in vivo by direct injection of virus or naked plasmid DNA. In a particular embodiment, the invention provides a method of fusing a spine by transfecting osteogenic precursor cells with an isolated nucleic acid molecule having a nucleotide sequence encoding LIM mineralization protein, admixing the transfected osteogenic precursor cells with a matrix and contacting the matrix with the spine. Finally, the invention relates to methods for inducing systemic bone formation by stable transfection of host cells with the vectors of the invention.

# LIM Mineralization Protein Splice Variants

#### BACKGROUND OF THE INVENTION

#### 5 1. Field of the Invention

10

15

20

25

30

The field of the invention relates generally to osteogenic cells and the formation of bone and boney tissue in mammalian species. Specifically, the invention concerns a novel family of proteins, and nucleic acids encoding those proteins, that enhances the efficacy of bone mineralization *in vitro* and *in vivo*. The invention provides methods for treating a variety of pathological conditions associated with bone and boney tissue, such as, for example, spine fusion, fracture repair and osteoporosis.

# 2. Description of the Related Art

Osteoblasts are thought to differentiate from pluripotent mesenchymal stem cells. The maturation of an osteoblast results in the secretion of an extracellular matrix which can mineralize and form bone. The regulation of this complex process is not well understood but is thought to involve a group of signaling glycoproteins known as bone morphogenetic proteins (BMPs). These proteins have been shown to be involved with embryonic dorsal-ventral patterning, limb bud development, and fracture repair in adult animals. B. L. Hogan, Genes & Develop., 10:1580 (1996). This group of transforming growth factor-beta superfamily secreted proteins has a spectrum of activities in a variety of cell types at different stages of differentiation; differences in physiological activity between these closely related molecules have not been clarified. D. M. Kingsley, Trends Genet., 10:16 (1994).

To better discern the unique physiological role of different BMP signaling proteins, we recently compared the potency of BMP-6 with that of BMP-2 and BMP-4, for inducing rat calvarial osteoblast differentiation. Boden *et al.*, Endocrinology, 137:3401 (1996). We studied this process in first passage (secondary) cultures of fetal rat calvaria that require BMP or glucocorticoid for

initiation of differentiation. In this model of membranous bone formation, glucocorticoid (GC) or a BMP will initiate differentiation to mineralized bone nodules capable of secreting osteocalcin, the osteoblast-specific protein. This secondary culture system is distinct from primary rat osteoblast cultures which undergo spontaneous differentiation. In this secondary system, glucocorticoid resulted in a ten-fold induction of BMP-6 mRNA and protein expression which was responsible for the enhancement of osteoblast differentiation. Boden *et al.*, Endocrinology, 138:2920 (1997).

In addition to extracellular signals, such as the BMPs, intracellular signals or regulatory molecules may also play a role in the cascade of events leading to formation of new bone. One broad class of intracellular regulatory molecules are the LIM proteins, which are so named because they possess a characteristic structural motif known as the LIM domain. The LIM domain is a cysteine-rich structural motif composed of two special zinc fingers that are joined by a 2-amino acid spacer. Some proteins have only LIM domains, while others contain a variety of additional functional domains. LIM proteins form a diverse group, which includes transcription factors and cytoskeletal proteins. The primary role of LIM domains appears to be in mediating protein-protein interactions, through the formation of dimers with identical or different LIM domains, or by binding distinct proteins.

In LIM homeodomain proteins, that is, proteins having both LIM domains and a homeodomain sequence, the LIM domains function as negative regulatory elements. LIM homeodomain proteins are involved in the control of cell lineage determination and the regulation of differentiation, although LIM-only proteins may have similar roles. LIM-only proteins are also implicated in the control of cell proliferation since several genes encoding such proteins are associated with oncogenic chromosome translocations.

Humans and other mammalian species are prone to diseases or injuries that require the processes of bone repair and/or regeneration. For example, treatment of fractures would be improved by new treatment regimens that could stimulate the natural bone repair mechanisms, thereby reducing the time

5

10

15

20

25

required for the fractured bone to heal. In another example, individuals afflicted with systemic bone disorders, such as osteoporosis, would benefit from treatment regimens that would results in systemic formation of new bone. Such treatment regimens would reduce the incidence of fractures arising from the loss of bone mass that is a characteristic of this disease.

For at least these reasons, extracellular factors, such as the BMPs, have been investigated for the purpose of using them to stimulate formation of new bone *in vivo*. Despite the early successes achieved with BMPs and other extracellular signalling molecules, their use entails a number of disadvantages. For example, relatively large doses of purified BMPs are required to enhance the production of new bone, thereby increasing the expense of such treatment methods. Furthermore, extracellular proteins are susceptible to degradation following their introduction into a host animal. In addition, because they are typically immunogenic, the possibility of stimulating an immune response to the administered proteins is ever present.

Due to such concerns, it would be desirable to have available treatment regimens that use an intracellular signalling molecule to induce new bone formation. Advances in the field of gene therapy now make it possible to introduce into osteogenic precursor cells, that is, cells involved in bone formation, or peripheral blood leukocytes, nucleotide fragments encoding intracellular signals that form part of the bone formation process. Gene therapy for bone formation offers a number of potential advantages: (1) lower production costs; (2) greater efficacy, compared to extracellular treatment regiments, due to the ability to achieve prolonged expression of the intracellular signal; (3) it would by-pass the possibility that treatment with extracellular signals might be hampered due to the presence of limiting numbers of receptors for those signals; (4) it permits the delivery of transfected potential osteoprogenitor cells directly to the site where localized bone formation is required; and (5) it would permit systemic bone formation, thereby providing a treatment regimen for osteoporosis and other metabolic bone diseases.

5

10

15

20

25

#### SUMMARY OF THE INVENTION

The present invention seeks to overcome the drawbacks in the prior art by providing novels compositions and methods for inducing bone formation using an intracellular signalling molecule that participates early in the cascade of events that leads to bone formation. Applicants have discovered 10-4/RLMP (SEQ ID NO: 1, SEQ ID NO: 2), a novel LIM gene with a sequence originally isolated from stimulated rat calvarial osteoblast cultures. The gene has been cloned, sequenced and assayed for its ability to enhance the efficacy of bone mineralization in vitro. The protein RLMP affects mineralization of bone matrix as well as differentiation of cells into the osteoblast lineage. Unlike other known cytokines, for example, BMPs, RLMP is not a secreted protein, but is instead an intracellular signaling molecule. This feature has the advantage of providing intracellular signaling amplification as well as easier assessment of transfected cells. It is also suitable for more efficient and specific in vivo applications. Suitable clinical applications include enhancement of bone repair in fractures, bone defects, bone grafting, and normal homeostasis in patients presenting with osteoporosis.

Applicants have also cloned, sequenced and deduced the amino acid sequence of a corresponding human protein, named human LMP-1. The human protein demonstrates enhanced efficacy of bone mineralization *in vitro* and *in vivo*.

In addition, the applicants have characterized a truncated (short) version of LMP-1, termed HLMP-1s. This short version resulted from a point mutation in one source of a cDNA clone, providing a stop codon which truncated the protein. The short version (LMP-1s) is fully functional when expressed in cell culture and *in vivo*.

Using PCR analysis of human heart cDNA library, Applicants have identified two alternative splice variants (referred to as HLMP-2 and HLMP-3) that differ from HLMP-1 in a region between base pairs 325 and 444 in the

5

10

15

20

25

nucleotide sequence encoding HLMP-1. The HLMP-2 sequence has a 119 base pair deletion and an insertion of 17 base pairs in this region. Compared to HLMP-1, the nucleotide sequence encoding HLMP-3 has no deletions, but it does have the same 17 base pairs as HLMP-2, which are inserted at position 444 in the HLMP-1 sequence.

Additional features and advantages of the invention will be set forth in the description which follows, and in part will be apparent from the description, or may be learned by practice of the invention. The objectives and other advantages of the invention will be realized and attained by the methods and compositions of matter particularly pointed out in the written description and claims hereof.

In one broad aspect, the invention relates to an isolated nucleic acid molecule comprising a nucleic acid sequence encoding any LIM mineralization protein, wherein the nucleic acid molecule hybridizes under standard conditions to a nucleic acid molecule complementary to the full length of SEQ. ID NO: 25, and wherein the molecule hybridizes under highly stringent conditions to a nucleic acid molecule complementary to the full length of SEQ. ID NO: 26. In a specific aspect, the isolated nucleic acid molecule encodes HLMP-1, HLMP-1s, RLMP, HLMP-2, or HLMP-3. In addition, the invention is directed to vectors comprising these nucleic acid molecules, as well as host cells comprising the vectors. In another specific aspect, the invention relates to the proteins themselves.

In a second broad aspect, the invention relates to antibody that is specific for LIM mineralization protein, including HLMP-1, HLMP-1s, RLMP, HLMP-2, and HLMP-3. In one specific aspect, the antibody is a polyclonal antibody. In another specific aspect, the antibody is a monoclonal antibody.

In a third broad aspect, the invention relates to method of inducing bone formation wherein osteogenic precursor cells are transfected with an isolated nucleic acid molecule comprising a nucleotide sequence encoding LIM mineralization protein. In one specific aspect, the isolated nucleic acid molecule is in a vector, which may be a plasmid or a virus, such as adenovirus

5

10

15

20

25

or retrovirus. The transfection may occur *ex vivo* or *in vivo* by direct injection of the isolated nucleic acid molecule. The transfected isolated nucleic acid molecule may encode HLMP-1, HLMP-1s, RLMP, HLMP-2, or HLMP-3.

In a further aspect, the invention relates to methods of fusing a spine by transfecting osteogenic precursor cells with an isolated nucleic acid molecule having a nucleotide sequence encoding LIM mineralization protein, admixing the transfected osteogenic precursor cells with a matrix and contacting the matrix with the spine.

In yet another aspect, the invention relates to methods for inducing systemic bone formation by stable transfection of host cells with the vectors of the invention.

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory and are intended to provide further explanation of the invention as claimed.

15

10

5

#### ABBREVIATIONS AND DEFINITIONS

	BMP	Bone Morphogenetic Protein
	HLMP-1	Human LMP-1, also
20		designated as Human LIM
		Protein or HLMP
	HLMP-1s	Human LMP-1 Short
		(truncated) protein
	HLMPU	Human LIM Protein Unique
25		Region
	LMP	LIM mineralization protein
	MEM	Minimal essential medium
	Trm	Triamcinolone
	-GlyP	Beta-glycerolphosphate
30	RACE	Rapid Amplification of cDNA
		Ends

	RLMP	Rat LIM mineralization protein,
		also designated as RLMP-1
	RLMPU	Rat LIM Protein Unique
		Region
5	RNAsin	RNase inhibitor
	ROB	Rat Osteoblast
	10-4	Clone containing cDNA
		sequence for RLMP (SEQ ID
		NO: 2)
10	UTR	Untranslated Region
	HLMP-2	Human LMP Splice Variant 2
	HLMP-3	Human LMP Splice Variant 3

# DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to novel mammalian LIM proteins, herein designated LIM mineralization proteins, or LMP. The invention relates more particularly to human LMP, known as HLMP or HLMP-1, or alternative splice variants of human LMP, which are known as HLMP-2 or HLMP-3. The applicants have discovered that these proteins enhance bone mineralization in mammalian cells grown *in vitro*. When produced in mammals, LMP also induces bone formation *in vivo*.

Ex vivo transfection of bone marrow cells, osteogenic precursor cells, peripheral blood leukocytes, or mesenchymal stem cells with nucleic acid that encodes LMP or HLMP, followed by reimplantation of the transfected cells in the donor, is suitable for treating a variety of bone-related disorders or injuries. For example, one can use this method to: augment long bone fracture repair; generate bone in segmental defects; provide a bone graft substitute for

15

20

25

fractures; facilitate tumor reconstruction or spine fusion; and provide a local treatment (by injection) for weak or osteoporotic bone, such as in osteoporosis of the hip, vertebrae, or wrist. Transfection with LMP or HLMP-encoding nucleic acid is also useful in: the percutaneous injection of transfected marrow cells to accelerate the repair of fractured long bones; treatment of delayed union or non-unions of long bone fractures or pseudoarthrosis of spine fusions; and for inducing new bone formation in avascular necrosis of the hip or knee.

In addition to ex vivo-based methods of gene therapy, transfection of a recombinant DNA vector comprising a nucleic acid sequence that encodes LMP or HLMP can be accomplished in vivo. When a DNA fragment that encodes LMP or HLMP is inserted into an appropriate viral vector, for example, an adenovirus vector, the viral construct can be injected directly into a body site were endochondral bone formation is desired. By using a direct, percutaneous injection to introduce the LMP or HLMP sequence stimulation of bone formation can be accomplished without the need for surgical intervention either to obtain bone marrow cells (to transfect ex vivo) or to reimplant them into the patient at the site where new bone is required. Alden et al., Neurosurgical Focus (1998), have demonstrated the utility of a direct injection method of gene therapy using a cDNA that encodes BMP-2, which was cloned into an adenovirus vector.

It is also possible to carry out *in vivo* gene therapy by directly injecting into an appropriate body site, a naked, that is, unencapsulated, recombinant plasmid comprising a nucleic acid sequence that encodes HLMP. In this embodiment of the invention, transfection occurs when the naked plasmid DNA is taken up, or internalized, by the appropriate target cells, which have been described. As in the case of *in vivo* gene therapy using a viral construct, direct injection of naked plasmid DNA offers the advantage that little or no surgical intervention is required. Direct gene therapy, using naked plasmid DNA that encodes the endothelial cell mitogen VEGF (vascular endothelial growth factor), has been successfully demonstrated in human patients. Baumgartner *et al.*, Circulation, 97(12):1114-23 (1998).

5

10

15

20

25

By using an adenovirus vector to deliver LMP into osteogenic cells, transient expression of LMP is achieved. This occurs because adenovirus does not incorporate into the genome of target cells that are transfected. Transient expression of LMP, that is, expression that occurs during the lifetime of the transfected target cells, is sufficient to achieve the objects of the invention. Stable expression of LMP, however, can occur when a vector that incorporates into the genome of the target cell is used as a delivery vehicle. Retrovirus-based vectors, for example, are suitable for this purpose.

Stable expression of LMP is particularly useful for treating various systemic bone-related disorders, such as osteoporosis and osteogenesis imperfecta. For this embodiment of the invention, in addition to using a vector that integrates into the genome of the target cell to deliver an LMP-encoding nucleotide sequence into target cells, LMP expression is placed under the control of a regulatable promoter. For example, a promoter that is turned on by exposure to an exogenous inducing agent, such as tetracycline, is suitable. Using this approach, one can stimulate formation of new bone on a systemic basis by administering an effective amount of the exogenous inducing agent. Once a sufficient quantity of bone mass is achieved, administration of the exogenous inducing agent is discontinued. This process may be repeated as needed to replace bone mass lost, for example, as a consequence of osteoporosis.

Antibodies specific for HLMP are particularly suitable for use in methods for assaying the osteoinductive, that is, bone-forming, potential of patient cells. In this way one can identify patients at risk for slow or poor healing of bone repair. Also, HLMP-specific antibodies are suitable for use in marker assays to identify risk factors in bone degenerative diseases, such as, for example, osteoporosis.

Following well known and conventional methods, the genes of the present invention are prepared by ligation of nucleic acid segments that encode LMP to other nucleic acid sequences, such as cloning and/or expression vectors. Methods needed to construct and analyze these

10

15

20

25

recombinant vectors, for example, restriction endonuclease digests, cloning protocols, mutagenesis, organic synthesis of oligonucleotides and DNA sequencing, have been described. For DNA sequencing DNA, the dieoxyterminator method is the preferred.

Many treatises on recombinant DNA methods have been published, including Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, 2nd edition (1988), Davis et al., Basic Methods in Molecular Biology, Elsevier (1986), and Ausubel et al., Current Protocols in Molecular Biology, Wiley Interscience (1988). These reference manuals are specifically incorporated by reference herein.

Primer-directed amplification of DNA or cDNA is a common step in the expression of the genes of this invention. It is typically performed by the polymerase chain reaction (PCR). PCR is described in U.S. Patent No. 4,800,159 to Mullis *et al.* and other published sources. The basic principle of PCR is the exponential replication of a DNA sequence by successive cycles of primer extension. The extension products of one primer, when hybridized to another primer, becomes a template for the synthesis of another nucleic acid molecule. The primer-template complexes act as substrate for DNA polymerase, which in performing its replication function, extends the primers. The conventional enzyme for PCR applications is the thermostable DNA polymerase isolated from *Thermus aquaticus*, or Taq DNA polymerase.

Numerous variations of the basic PCR method exist, and a particular procedure of choice in any given step needed to construct the recombinant vectors of this invention is readily performed by a skilled artisan. For example, to measure cellular expression of 10-4/RLMP, RNA is extracted and reverse transcribed under standard and well known procedures. The resulting cDNA is then analyzed for the appropriate mRNA sequence by PCR.

The gene encoding the LIM mineralization protein is expressed in an expression vector in a recombinant expression system. Of course, the constructed sequence need not be the same as the original, or its complimentary sequence, but instead may be any sequence determined by the

5

10

15

20

25

degeneracy of the DNA code that nonetheless expresses an LMP having bone forming activity. Conservative amino acid substitutions, or other modifications, such as the occurrance of an amino-terminal methionine residue, may also be employed.

A ribosome binding site active in the host expression system of choice is ligated to the 5' end of the chimeric LMP coding sequence, forming a synthetic gene. The synthetic gene can be inserted into any one of a large variety of vectors for expression by ligating to an appropriately linearized plasmid. A regulatable promoter, for example, the *E. coli* lac promoter, is also suitable for the expression of the chimeric coding sequences. Other suitable regulatable promoters include trp, tac, recA, T7 and lambda promoters.

DNA encoding LMP is transfected into recipient cells by one of several standard published procedures, for example, calcium phosphate precipitation, DEAE-Dextran, electroporation or protoplast fusion, to form stable transformants. Calcium phosphate precipitation is preferred, particularly when performed as follows.

DNAs are coprecipitated with calcium phosphate according to the method of Graham and Van Der, <u>Virology</u>, 52:456 (1973), before transfer into cells. An aliquot of 40-50 g of DNA, with salmon sperm or calf thymus DNA as a carrier, is used for  $0.5 \times 10^6$  cells plated on a 100 mm dish. The DNA is mixed with 0.5 ml of 2X Hepes solution (280 mM NaCl, 50 mM Hepes and 1.5 mM Na<sub>2</sub>HPO<sub>4</sub>, pH 7.0), to which an equal volume of 2x CaCl<sub>2</sub> (250 mM CaCl<sub>2</sub> and 10 mM Hepes, pH 7.0) is added. A white granular precipitate, appearing after 30-40 minutes, is evenly distributed dropwise on the cells, which are allowed to incubate for 4-16 hours at 37°C. The medium is removed and the cells shocked with 15% glycerol in PBS for 3 minutes. After removing the glycerol, the cells are fed with Dulbecco's Minimal Essential Medium (DMEM) containing 10% fetal bovine serum.

DNA can also be transfected using: the DEAE-Dextran methods of Kimura *et al.*, Virology, 49:394 (1972) and Sompayrac *et al.*, Proc. Natl. Acad. Sci. USA, 78:7575 (1981); the electroporation method of Potter, Proc. Natl.

5

10

15

20

25

Acad. Sci. USA, 81:7161 (1984); and the protoplast fusion method of Sandri-Goddin *et al.*, Molec. Cell. Biol., 1:743 (1981).

Phosphoramidite chemistry in solid phase is the preferred method for the organic synthesis of oligodeoxynucleotides and polydeoxynucleotides. In addition, many other organic synthesis methods are available. Those methods are readily adapted by those skilled in the art to the particular sequences of the invention.

The present invention also includes nucleic acid molecules that hybridize under standard conditions to any of the nucleic acid sequences encoding the LIM mineralization proteins of the invention. "Standard hybridization conditions" will vary with the size of the probe, the background and the concentration of the nucleic acid reagents, as well as the type of hybridization, for example, *in situ*, Southern blot, or hybrization of DNA-RNA hybrids (Northern blot). The determination of "standard hybridization conditions" is within the level of skill in the art. For example, see U.S. Patent 5,580,775 to Fremeau *et al.*, herein incorporated by reference for this purpose. See also, Southern, E. M., J. Mol. Biol., 98:503 (1975), Alwine *et al.*, Meth. Enzymol., 68:220 (1979), and Sambrook *et al.*, Molecular Cloning: A laboratory Manual, 2nd edition, pp. 7.19-7.50, Cold Spring Harbor Press (1989).

One preferred set of standard hybrization conditions involves a blot that is prehybridized at 42°C for 2 hours in 50% formamide, 5X SSPE (150 nM NaCl, 10 mM Na H<sub>2</sub>PO<sub>4</sub> [pH 7.4], 1 mM EDTA [pH 8.0]), 5X Denhardt's solution (20 mg Ficoll, 20 mg polyvinylpyrrolidone and 20 mg BSA per 100 ml water), 10% dextran sulphate, 1% SDS and 100 g/ml salmon sperm DNA. A <sup>32</sup>P-labelled cDNA probe is added, and hybridization is continued for 14 hours. Afterward, the blot is washed twice with 2X SSPE, 0.1% SDS for 20 minutes at 22°C, followed by a 1 hour wash at 65°C in 0.1X SSPE, 0.1 %SDS. The blot is then dried and exposed to x-ray film for 5 days in the presence of an intensifying screen.

5

10

15

20

25

Under "highly stringent conditions," a probe will hybridize to its target sequence if those two sequences are substantially identical. As in the case of standard hybridization conditions, one of skill in the art can, given the level of skill in the art and the nature of the particular experiment, determine the conditions under which only susbstantially identical sequences will hybridize.

Another aspect of the invention includes the proteins encoded by the nucleic acid sequences. In still another embodiment, the inventon relates to the identification of such proteins based on anti-LMP antibodies. In this embodiment, protein samples are prepared for Western blot analysis by lysing cells and separating the proteins by SDS-PAGE. The proteins are transferred to nitrocellulose by electroblotting as described by Ausubel *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons (1987). After blocking the filter with instant nonfat dry milk (1 gm in 100 ml PBS), anti-LMP antibody is added to the filter and incubated for 1 hour at room temperature. The filter is washed thoroughly with phosphate buffered saline (PBS) and incubated with horseradish peroxidase (HRPO)-antibody conjugate for 1 hour at room temperature. The filter is again washed thoroughly with PBS and the antigen bands are identified by adding diaminobenzidine (DAB).

Monospecific antibodies are the reagent of choice in the present invention, and are specifically used to analyze patient cells for specific characteristics associated with the expression of LMP. "Monospecific antibody" as used herein is defined as a single antibody species or multiple antibody species with homogenous binding characteristics for LMP. "Homogeneous binding" as used herein refers to the ability of the antibody species to bind to a specific antigen or epitope, such as those associated with LMP, as described above. Monospecific antibodies to LMP are purified from mammalian antisera containing antibodies reactive against LMP or are prepared as monoclonal antibodies reactive with LMP using the technique of Kohler and Milstein, Nature, 256:495-97 (1975). The LMP specific antibodies are raised by immunizing animals such as, for example, mice, rats, guinea

5

10

15

20

25

pigs, rabbits, goats or horses, with an appropriate concentration of LMP either with or without an immune adjuvant.

In this process, preimmune serum is collected prior to the first immunization. Each animal receives between about 0.1 mg and about 1000 mg of LMP associated with an acceptable immune adjuvant, if desired. Such acceptable adjuvants include, but are not limited to, Freund's complete, Freund's incomplete, alum-precipitate, water in oil emulsion containing Corynebacterium parvum and tRNA adjuvants. The initial immunization consists of LMP in, preferably, Freund's complete adjuvant injected at multiple sites either subcutaneously (SC), intraperitoneally (IP) or both. Each animal is bled at regular intervals, preferably weekly, to determine antibody titer. The animals may or may not receive booster injections following the initial immunization. Those animals receiving booster injections are generally given an equal amount of the antigen in Freund's incomplete adjuvant by the same route. Booster injections are given at about three week intervals until maximal titers are obtained. At about 7 days after each booster immunization or about weekly after a single immunization, the animals are bled, the serum collected, and aliquots are stored at about -20° C.

Monoclonal antibodies (mAb) reactive with LMP are prepared by immunizing inbred mice, preferably Balb/c mice, with LMP. The mice are immunized by the IP or SC route with about 0.1 mg to about 10 mg, preferably about 1 mg, of LMP in about 0.5 ml buffer or saline incorporated in an equal volume of an acceptable adjuvant, as discussed above. Freund's complete adjuvant is preferred. The mice receive an initial immunization on day 0 and are rested for about 3-30 weeks. Immunized mice are given one or more booster immunizations of about 0.1 to about 10 mg of LMP in a buffer solution such as phosphate buffered saline by the intravenous (IV) route. Lymphocytes from antibody-positive mice, preferably splenic lymphocytes, are obtained by removing the spleens from immunized mice by standard procedures known in the art. Hybridoma cells are produced by mixing the splenic lymphocytes with an appropriate fusion partner, preferably myeloma cells, under conditions

5

10

15

20

25

which will allow the formation of stable hybridomas. Fusion partners may include, but are not limited to: mouse myelomas P3/NS1/Ag 4-1; MPC-11; S-194 and Sp 2/0, with Sp 2/0 being preferred. The antibody producing cells and myeloma cells are fused in polyethylene glycol, about 1000 mol. wt., at concentrations from about 30% to about 50%. Fused hybridoma cells are selected by growth in hypoxanthine, thymidine and aminopterin in supplemented Dulbecco's Modified Eagles Medium (DMEM) by procedures known in the art. Supernatant fluids are collected from growth positive wells on about days 14, 18, and 21, and are screened for antibody production by an immunoassay such as solid phase immunoradioassay (SPIRA) using LMP as the antigen. The culture fluids are also tested in the Ouchterlony precipitation assay to determine the isotype of the mAb. Hybridoma cells from antibody positive wells are cloned by a technique such as the soft agar technique of MacPherson, "Soft Agar Techniques", in Tissue Culture Methods and Applications, Kruse and Paterson (eds.), Academic Press (1973). See, also, Harlow et al., Antibodies: A Laboratory Manual, Cold Spring Laboratory (1988).

Monoclonal antibodies may also be produced *in vivo* by injection of pristane- primed Balb/c mice, approximately 0.5 ml per mouse, with about 2x10<sup>6</sup> to about 6x10<sup>6</sup> hybridoma cells about 4 days after priming. Ascites fluid is collected at approximately 8-12 days after cell transfer and the monoclonal antibodies are purified by techniques known in the art.

In vitro production in anti-LMP mAb is carried out by growing the hydridoma cell line in DMEM containing about 2% fetal calf serum to obtain sufficient quantities of the specific mAb. The mAb are purified by techniques known in the art.

Antibody titers of ascites or hybridoma culture fluids are determined by various serological or immunological assays, which include, but are not limited to, precipitation, passive agglutination, enzyme-linked immunosorbent antibody (ELISA) technique and radioimmunoassay (RIA) techniques. Similar assays

10

15

20

25

are used to detect the presence of the LMP in body fluids or tissue and cell extracts.

It is readily apparent to those skilled in the art that the above described methods for producing monospecific antibodies may be utilized to produce antibodies specific for polypeptide fragments of LMP, full-length nascent LMP polypeptide, or variants or alleles thereof.

In another embodiment, the invention is directed to alternative splice variants of HLMP-1. PCR analysis of human heart cDNA revealed mRNA for two HLMP alternative splice variants, named HLMP-2 and HLMP-3, that differ from HLMP-1 in a region between base pairs 325 and 444 in the hLMP-1 sequence. The HLMP-2 sequence has a 119 base pair deletion and an insertion of 17 base pairs in this region. These changes preserve the reading frame, resulting in a 423 amino acid protein, which compared to HLMP-1, has a net loss of 34 amino acids (40 amino acids deleted plus 6 inserted amino acids). HLMP-2 contains the c-terminal LIM domains that are present in HLMP-1.

Compared to HLMP-1, HLMP-3 has no deletions, but it does have the same 17 base pair insertion at position 444. This insertion shifts the reading frame, causing a stop codon at base pairs 459-461. As a result, HLMP-3 encodes a protein of 153 amino acids. This protein lacks the c-terminal LIM domains that are present in HLMP-1 and HLMP-2. The predicted size of the proteins encoded by HLMP-2 and HLMP-3 was confirmed by western blot analysis.

PCR analysis of the tissue distribution of the three splice variants revealed that they are differentially expressed, with specific isoforms predominating in different tissues. HLMP-1 is apparently the predominant form expressed in leukocytes, spleen, lung, placenta, and fetal liver. HLMP-2 appears to be the predominant isoform in skeletal muscle, bone marrow, and heart tissue. HLMP-3, however, was not the predominant isoform in any tissue examined.

5

10

15

20

25

Overexpression of HLMP-3 in secondary rat osteoblast cultures induced bone nodule formation (287±56) similar to the effect seen for glucicorticoid (272±7) and HLMP-1 (232±200). Since HLMP-3 lacks the C-terminal LIM domains, there regions are not required for osteoinductive activity. Overexpression of HLMP-2, however, did not induce nodule formation (11±3). These data suggest that the amino acids encoded by the deleted 119 base pairs are necessary for osteoinduction. The data also suggest that the distribution of HLMP splice variants may be important for tissue-specific function. Surprisingly, we have shown that HLMP-2 inhibits steroid-induced osteoblast formation in secondary rat osteoblast cultures. Therefore, HLMP-2 will have therapeutic utility in clinical situations where bone formation is not desirable.

On July 22, 1997, a sample of 10-4/RLMP in a vector designated pCMV2/RLMP (which is vector pRc/CMV2 with insert 10-4 clone/RLMP) was deposited with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, MD 20852. The culture accession number for that deposit is 209153. On March 19, 1998, a sample of the vector pHis-A with insert HLPM-1s was deposited at the American Type Culture Collection ("ATCC"). The culture accession number for that deposit is 209698. On April 14, 2000, samples of plasmids pHAhLMP-2 (vector pHisA with cDNA insert derived from human heart muscle cDNA with HLMP-2) and pHAhLMP-3 (vector pHisA with cDNA insert derived from human heart muscle cDNA with HLMP-3) were deposited with the ATCC, 10801 University Blvd., Manassas, VA, 20110-2209, USA, under the conditions of the Budapest treaty. The accession numbers for these deposits are \_\_\_\_\_ and \_\_\_\_, respectively. These deposits, as required by the Budapest Treaty, will be maintained in the ATCC for at least 30 years and will be made available to the public upon the grant of a patent disclosing them. It should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by government action.

5

10

15

20

25

In assessing the nucleic acids, proteins, or antibodies of the invention, enzyme assays, protein purification, and other conventional biochemical methods are employed. DNA and RNA are analyzed by Southern blotting and Northern blotting techniques, respectively. Typically, the samples analyzed are size fractionated by gel electrophoresis. The DNA or RNA in the gels are then transferred to nitrocellulose or nylon membranes. The blots, which are replicas of sample patterns in the gels, were then hybridized with probes. Typically, the probes are radiolabelled, preferably with <sup>32</sup>P, although one could label the probes with other signal-generating molecules known to those in the art. Specific bands of interest can then be visualized by detection systems, such as autoradiography.

For purposes of illustrating preferred embodiments of the present invention, the following, non-limiting examples are included. These results demonstrate the feasibility of inducing or enhancing the formation of bone using the LIM mineralization proteins of the invention, and the isolated nucleic acid molecules encoding those proteins.

Rat calvarial cells, also known as rat osteoblasts ("ROB"), were obtained

#### Example 1: Calvarial Cell Culture

20-day pre-parturition rats as previously described. Boden *et al.*, Endocrinology, 137(8):3401-07 (1996). Primary cultures were grown to confluence (7 days), trypsinized, and passed into 6-well plates (1 x 10<sup>5</sup> cells/35 mm well) as first subculture cells. The subculture cells, which were confluent at day 0, were grown for an additional 7 days. Beginning on day 0, media were changed and treatments (Trm and/or BMPs) were applied, under a laminar flow hood, every 3 or 4 days. The standard culture protocol was as follows: days 1-7, MEM, 10% FBS, 50 g/ml ascorbic acid, ± stimulus; days 8-14, BGJb medium, 10% FBS, 5mM -GlyP (as a source of inorganic phosphate to permit mineralization). Endpoint analysis of bone nodule formation and osteocalcin secretion was performed at day 14. The dose of BMP was chosen as 50 ng/ml

5

10

15

20

25

based on pilot experiments in this system that demonstrated a mid-range effect on the dose-response curve for all BMPs studied.

#### EXAMPLE 2: Antisense Treatment and Cell Culture

5

10

15

20

25

To explore the potential functional role of LMP-1 during membranous bone formation, we synthesized an antisense oligonucleotide to block LMP-1 mRNA translation and treated secondary osteoblast cultures that were undergoing differentiation initiated by glucocorticoid. Inhibition of RLMP expression was accomplished with a highly specific antisense oligonucleotide (having no significant homologies to known rat sequences) corresponding to a 25 bp sequence spanning the putative translational start site (SEQ ID NO: 42). Control cultures either did not receive oligonucleotide or they received sense oligonucleotide. Experiments were performed in the presence (preincubation) and absence of lipofectamine. Briefly, 22 g of sense or antisense RLMP oligonucleotide was incubated in MEM for 45 minutes at room temperature. Following that incubation, either more MEM or pre-incubated lipofectamine/MEM (7% v/v; incubated 45 minutes at room temperature) was added to achieve an oligonucleotide concentration of 0.2 M. The resulting mixture was incubated for 15 minutes at room temperature. Oligonucleotide mixtures were then mixed with the appropriate medium, that is, MEM/Ascorbate/±Trm, to achieve a final oligonucleotide concentration of 0.1 M.

Cells were incubated with the appropriate medium (±stimulus) in the presence or absence of the appropriate oligonucleotides. Cultures originally incubated with lipofectamine were re-fed after 4 hours of incubation (37°C; 5% CO<sub>2</sub>) with media containing neither lipofectamine nor oligonucleotide. All cultures, especially cultures receiving oligonucleotide, were re-fed every 24 hours to maintain oligonucleotide levels.

30

LMP-1 antisense oligonucleotide inhibited mineralized nodule formation and osteocalcin secretion in a dose-dependent manner, similar to the effect of

BMP-6 oligonucleotide. The LMP-1 antisense block in osteoblast differentiation could not be rescued by addition of exogenous BMP-6, while the BMP-6 antisense oligonucleotide inhibition was reversed with addition of BMP-6. This experiment further confirmed the upstream position of LMP-1 relative to BMP-6 in the osteoblast differentiation pathway. LMP-1 antisense oligonucleotide also inhibited spontaneous osteoblast differentiation in primary rat osteoblast cultures.

# **EXAMPLE 3: Quantitation of Mineralized Bone Nodule Formation**

10

15

20

5

Cultures of ROBs prepared according to Examples 1 and 2 were fixed overnight in 70% ethanol and stained with von Kossa silver stain. A semi-automated computerized video image analysis system was used to quantitate nodule count and nodule area in each well. Boden et al., Endocrinology, 137(8):3401-07 (1996). These values were then divided to calculate the area per nodule values. This automated process was validated against a manual counting technique and demonstrated a correlation coefficient of 0.92 (p < 0.000001). All data are expressed as the mean ± standard error of the mean (S.E.M.) calculated from 5 or 6 wells at each condition. Each experiment was confirmed at least twice using cells from different calvarial preparations.

#### **EXAMPLE 4: Quantitation of Osteocalcin Secretion**

Osteocalcin levels in the culture media were measured using a

competitive radioimmunoassay with a monospecific polyclonal antibody (Pab) raised in our laboratory against the C-terminal nonapeptide of rat osteocalcin as described in Nanes *et al.*, Endocrinology, 127:588 (1990). Briefly, 1 g of nonapeptide was iodinated with 1 mCi <sup>125</sup>I-Na by the lactoperoxidase method. Tubes containing 200 I of assay buffer (0.02 M sodium phosphate, 1 mM

EDTA, 0.001% thimerosal, 0.025% BSA) received media taken from cell cultures or osteocalcin standards (0 - 12,000 fmole) at 100 I/tube in assay

buffer. The Pab (1:40,000; 100 I) was then added, followed by the iodinated peptide (12,000 cpm; 100 I). Samples tested for non-specific binding were prepared similarly but contained no antibody.

Bound and free PAbs were separated by the addition of 700 I goat antirabbit IgG, followed by incubation for 18 hours at 4°C. After samples were centrifuged at 1200 rpm for 45 minutes, the supernatants were decanted and the precipitates counted in a gamma counter. Osteocalcin values were reported in fmole/100 I, which was then converted to pmole/ml medium (3-day production) by dividing those values by 100. Values were expressed as the mean ± S.E.M. of triplicate determinations for 5-6 wells for each condition. Each experiment was confirmed at least two times using cells from different calvarial preparations.

#### EXAMPLE 5: Effect of Trm and RLMP on Mineralization In Vitro

15

20

25

30

10

5

There was little apparent effect of either the sense or antisense oligonucleotides on the overall production of bone nodules in the non-stimulated cell culture system. When ROBs were stimulated with Trm, however, the antisense oligonucleotide to RLMP inhibited mineralization of nodules by > 95%. The addition of exogenous BMP-6 to the oligonucleotide-treated cultures did not rescue the mineralization of RLMP-antisense-treated nodules.

Osteocalcin has long been synonymous with bone mineralization, and osteocalcin levels have been correlated with nodule production and mineralization. The RLMP-antisense oligonucleotide significantly decreases osteocalcin production, but the nodule count in antisense-treated cultures does not change significantly. In this case, the addition of exogenous BMP-6 only rescued the production of osteocalcin in RLMP-antisense-treated cultures by 10-15%. This suggests that the action of RLMP is downstream of, and more specific than, BMP-6.

#### **EXAMPLE 6: Harvest and Purification of RNA**

Cellular RNA from duplicate wells of ROBs (prepared according to Examples 1 and 2 in 6-well culture dishes) was harvested using 4M guanidine isothiocyanate (GIT) solution to yield statistical triplicates. Briefly, culture supernatant was aspirated from the wells, which were then overlayed with 0.6 ml of GIT solution per duplicate well harvest. After adding the GIT solution, the plates were swirled for 5-10 seconds (being as consistent as possible). Samples were saved at -70°C for up to 7 days before further processing.

RNA was purified by a slight modification of standard methods according to Sambrook *et al.*, Molecular Cloning: a Laboratory Manual, 2nd Ed., chapter 7.19, Cold Spring Harbor Press (1989). Briefly, thawed samples received 60 1 2.0 M sodium acetate (pH 4.0), 550 1 phenol (water saturated) and 150 1 chloroform:isoamyl alcohol (49:1). After vortexing, the samples were centrifuged (10000 x g; 20 minutes; 4°C), the aqueous phase transferred to a fresh tube, 600 1 isopropanol was added and the RNA precipitated overnight at -20°C.

Following the overnight incubation, the samples were centrifuged (10000 x g; 20 minutes) and the supernatant was aspirated gently. The pellets were resuspended in 400 TDEPC-treated water, extracted once with phenol:chloroform (1:1), extracted with chloroform:isoamyl alcohol (24:1) and precipitated overnight at -20°C after addition of 40 T sodium acetate (3.0 M; pH 5.2) and 1.0 ml absolute ethanol. To recover the cellular RNA, the samples were centrifuged (10000 x g; 20 min), washed once with 70% ethanol, air dried for 5-10 minutes and resuspended in 20 T of DEPC-treated water. RNA concentrations were calculated from optical densities that were determined with a spectrophotometer.

30

5

10

15

20

# **EXAMPLE 7: Reverse Transcription-Polymerase Chain Reaction**

Heated total RNA (5 g in 10.5 I total volume DEPC-H<sub>2</sub>O at 65°C for 5 minutes) was added to tubes containing 4 I 5X MMLV-RT buffer, 2 I dNTPs, 2 I dT17 primer (10 pmol/ml), 0.5 I RNAsin (40U/ml) and 1 I MMLV-RT (200 units/I). The samples were incubated at 37°C for 1 hour, then at 95°C for 5 minutes to inactivate the MMLV-RT. The samples were diluted by addition of 80 I of water.

Reverse-transcribed samples (5 I) were subjected to polymerase-chain reaction using standard methodologies (50 I total volume). Briefly, samples were added to tubes containing water and appropriate amounts of PCR buffer, 25 mM MgCl<sub>2</sub>, dNTPs, forward and reverse primers for glyceraldehyde 3-phosphate dehydrogenase (GAP, a housekeeping gene) and/or BMP-6), <sup>32</sup>P-dCTP, and Taq polymerase. Unless otherwise noted, primers were standardized to run consistently at 22 cycles (94°C, 30"; 58°C, 30"; 72°C, 20").

# EXAMPLE 8: Quantitation of RT-PCR Products by Polyacrylamide Gel Electrophoresis (PAGE) and Phosphorlmager Analysis

20

5

10

15

RT-PCR products received 5 I/tube loading dye, were mixed, heated at 65°C for 10 min and centrifuged. Ten I of each reaction was subjected to PAGE (12% polyacrylamide:bis; 15 V/well; constant current) under standard conditions. Gels were then incubated in gel preserving buffer (10% v/v glycerol, 7% v/v acetic acid, 40% v/v methanol, 43% deionized water) for 30 minutes, dried (80°C) *in vacuo* for 1-2 hours and developed with an electronically-enhanced phosphoresence imaging system for 6-24 hours. Visualized bands were analyzed. Counts per band were plotted graphically.

30

## **EXAMPLE 9: Differential Display PCR**

RNA was extracted from cells stimulated with glucocorticoid (Trm, 1 nM). Heated, DNase-treated total RNA (5 g in 10.5 I total volume in DEPC-H<sub>2</sub>O at 65°C for 5 minutes) was reverse transcribed as described in Example 7, but H-T<sub>11</sub>M (SEQ ID. NO: 4) was used as the MMLV-RT primer. The resulting cDNAs were PCR-amplified as described above, but with various commercial primer sets (for example, H-T<sub>11</sub>G (SEQ ID NO: 4) and H-AP-10 (SEQ ID. NO: 5); GenHunter Corp. Nashville, TN). Radiolabelled PCR products were fractionated by get 10 electrophoresis on a DNA sequencing gel. After electrophoresis, the resulting gels were dried in vacuo and autoradiographs were exposed overnight. Bands representing differentially-expressed cDNAs were excised from the gel and reamplified by PCR using the method of Conner et al., Proc. Natl. Acad. Sci. <u>USA</u>, 88:278 (1983). The products of PCR reamplification were cloned into 15 the vector PCR-II (TA cloning kit; InVitrogen, Carlsbad, CA).

# EXAMPLE 10: Screening of a UMR 106 Rat Osteosarcoma Cell cDNA Library

A UMR 106 library (2.5 x 10<sup>10</sup> pfu/ml) was plated at 5 x 10<sup>4</sup> pfu/ml onto agar plates (LB bottom agar) and the plates were incubated overnight at 37°C. Filter membranes were overlaid onto plates for two minutes. Once removed, the filters were denatured, rinsed, dried and UV cross-linked. The filters were then incubated in pre-hyridization buffer (2X PIPES [pH 6.5], 5% formamide, 1% SDS and 100 g/ml denatured salmon sperm DNA) for 2 h at 42°C. A 260 base-pair radiolabelled probe (SEQ ID NO: 3; <sup>32</sup>P labelled by random priming) was added to the entire hybridization mix/filters, followed by hybridization for 18 hours at 42°C. The membranes were washed once at room temperature (10 min, 1 x SSC, 0.1% SDS) and three times at 55°C (15 min, 0.1 x SSC, 0.1% SDS).

After they were washed, the membranes were analyzed by autoradiography as described above. Positive clones were plaque purified. The procedure was repeated with a second filter for four minutes to minimize spurious positives. Plaque-purified clones were rescued as lambda SK(-) phagemids. Cloned cDNAs were sequenced as described below.

## **EXAMPLE 11: Sequencing of Clones**

5

10

15

20

25

30

Cloned cDNA inserts were sequenced by standard methods. Ausubel *et al.*, <u>Current Protocols in Molecular Biology</u>, Wiley Interscience (1988). Briefly, appropriate concentrations of termination mixture, template and reaction mixture were subjected to an appropriate cycling protocol (95°C,30s; 68°C,30s; 72°C,60s; x 25). Stop mixture was added to terminate the sequencing reactions. After heating at 92°C for 3 minutes, the samples were loaded onto a denaturing 6% polyacrylamide sequencing gel (29:1 acrylamide:bisacrylamide). Samples were electrophoresed for about 4 hours at 60 volts, constant current. After electrophoresis, the gels were dried *in vacuo* and autoradiographed.

The autoradiographs were analyzed manually. The resulting sequences were screened against the databases maintained by the National Center for Biotechnology Information (NIH, Bethesda, MD; http://www.ncbi.nlm.nih.gov/) using the BLASTn program set with default parameters. Based on the sequence data, new sequencing primers were prepared and the process was repeated until the entire gene had been sequenced. All sequences were confirmed a minimum of three times in both orientations.

Nucleotide and amino acid sequences were also analyzed using the PCGENE software package (version 16.0). Per cent homology values for nucleotide sequences were calculated by the program NALIGN, using the following parameters: weight of non-matching nucleotides, 10; weight of non-matching gaps, 10; maximum number of nucleotides considered, 50; and minimum number of nucleotides considered, 50.

For amino acid sequences, per cent homology values were calculated using PALIGN. A value of 10 was selected for both the open gap cost and the unit gap cost.

## 5 EXAMPLE 12: Cloning of RLMP cDNA

The differential display PCR amplification products described in Example 9 contained a major band of approximately 260 base pairs. This sequence was used to screen a rat osteosarcoma (UMR 106) cDNA library. Positive clones were subjected to nested primer analysis to obtain the primer sequences necessary for amplifying the full length cDNA. (SEQ. ID NOs: 11, 12, 29, 30 and 31) One of those positive clones selected for further study was designated clone 10-4.

Sequence analysis of the full-length cDNA in clone 10-4, determined by nested primer analysis, showed that clone 10-4 contained the original 260 base-pair fragment identified by differential display PCR. Clone 10-4 (1696 base pairs; SEQ ID NO: 2) contains an open reading frame of 1371 base pairs encoding a protein having 457 amino acids (SEQ ID NO: 1). The termination codon, TGA, occurs at nucleotides 1444-1446. The polyadenylation signal at nucleotides 1675-1680, and adjacent poly(A)\* tail, was present in the 3' noncoding region. There were two potential N-glycosylation sites, Asn-Lys-Thr and Asn-Arg-Thr, at amino acid positions 113-116 and 257-259 in SEQ ID NO: 1, respectively. Two potential cAMP- and cGMP-dependent protein kinase phosphorylation sites, Ser and Thr, were found at amino acid positions 191 and 349, respectively. There were five potential protein kinase C phosphorylation sites, Ser or Thr, at amino acid positions 3, 115, 166, 219, 442. One potential ATP/GTP binding site motif A (P-loop), Gly-Gly-Ser-Asn-Asn-Gly-Lys-Thr, was determined at amino acid positions 272-279.

In addition, two highly conserved putative LIM domains were found at amino acid positions 341-391 and 400-451. The putative LIM domains in this newly identified rat cDNA clone showed considerable homology with the LIM

10

15

20

25

domains of other known LIM proteins. However, the overall homology with other rat LIM proteins was less than 25%. RLMP (also designated 10-4) has 78.5% amino acid homology to the human enigma protein (see U.S. Patent No. 5,504,192), but only 24.5% and 22.7% amino acid homology to its closest rat homologs, CLP-36 and RIT-18, respectively.

## **EXAMPLE 13: Northern Blot Analysis of RLMP Expression**

Thirty g of total RNA from ROBs, prepared according to Examples 1 and 2, was size fractionated by formaldehyde gel electrophoresis in 1% agarose flatbed gels and esmotically transblotted to nylon membranes. The blot was probed with a 600 base pair EcoR1 fragment of full-length 10-4 cDNA labeled with <sup>32</sup>P-dCTP by random priming.

Northern blot analysis showed a 1.7 kb mRNA species that hybridized with the RLMP probe. RLMP mRNA was up-regulated approximately 3.7-fold in ROBs after 24 hours exposure to BMP-6. No up-regulation of RMLP expression was seen in BMP-2 or BMP-4-stimulated ROBs at 24 hours.

### **EXAMPLE 14: Statistical Methods**

20

25

30

10

15

For each reported nodule/osteocalcin result, data from 5-6 wells from a representative experiment were used to calculate the mean ± S.E.M. Graphs may be shown with data normalized to the maximum value for each parameter to allow simultaneous graphing of nodule counts, mineralized areas and osteocalcin.

For each reported RT-PCR, RNase protection assay or Western blot analysis, data from triplicate samples of representative experiments, were used to determine the mean ± S.E.M. Graphs may be shown normalized to either day 0 or negative controls and expressed as fold-increase above control values.

Statistical significance was evaluated using a one-way analysis of variance with post-hoc multiple comparison corrections of Bonferroni as appropriate. D. V. Huntsberger, "The Analysis of Variance," in Elements of Statistical Variance, P. Billingsley (ed.), pp. 298-330, Allyn & Bacon Inc., Boston, MA (1977) and Sigmastat, Jandel Scientific, Corte Madera, CA. Alpha levels for significance were defined as p < 0.05.

# EXAMPLE 15: Detection of Rat LIM Mineralization Protein by Western Blot Analysis

10

Polyclonal antibodies were prepared according to the methods of England *et al.*, <u>Biochim.Biophys. Acta</u>, 623:171 (1980) and Timmer *et al.*, <u>J. Biol. Chem.</u>, 268:24863 (1993).

HeLa cells were transfected with pCMV2/RLMP. Protein was harvested from the transfected cells according to the method of Hair et al., Leukemia Research, 20:1 (1996). Western Blot Analysis of native RLMP was performed as described by Towbin et al., Proc. Natl. Acad. Sci. USA, 76:4350 (1979).

# EXAMPLE 16: Synthesis of the Rat LMP-Unique (RLMPU) derived Human 20 PCR product

Based on the sequence of the rat LMP-1 cDNA, forward and reverse PCR primers (SEQ ID NOs: 15 and 16) were synthesized and a unique 223 base-pair sequence was PCR amplified from the rat LMP-1 cDNA. A similar PCR product was isolated from human MG63 osteosarcoma cell cDNA with the same PCR primers.

RNA was harvested from MG63 osteosarcoma cells grown in T-75 flasks. Culture supernatant was removed by aspiration and the flasks were overlayed with 3.0 ml of GIT solution per duplicate, swirled for 5-10 seconds, and the resulting solution was transferred to 1.5 ml eppendorf tubes (5 tubes with 0.6 ml/tube). RNA was purified by a slight modification of standard

25

methods, for example, see Sambrook et al., Molecular Cloning: A Laboratory Manual, chapter 7, page 19, Cold Spring Harbor Laboratory Press (1989) and Boden et al., Endocrinology, 138:2820-28 (1997). Briefly, the 0.6 ml samples received 60 12.0 M sodium acetate (pH 4.0), 550 I water saturated phenol and 150 I chloroform:isoamyl alcohol (49:1). After addiiton of those reagents, the samples were vortexed, centrifuged (10000 x, q; 20 min; 4C) and the aqueous phase transferred to a fresh tube. Isopropanol (600 I) was added and the RNA was precipitated overnight at -20°C. The samples were centrifuged (10000 x g; 20 minutes) and the supernatant was aspirated gently. The pellets were resuspended in 400 I of DEPC-treated water, extracted once with phenol:chloroform (1:1), extracted with chloroform; isoamyl alcohol (24:1) and precipitated overnight at -20°C in 40 I sodium acetate (3.0 M; pH 5.2) and 1.0 ml absolute ethanol. After precipitation, the samples were centrifuged (10000 x g; 20 min), washed once with 70% ethanol, air dried for 5-10 minutes and resuspended in 20 I of DEPC-treated water. RNA concentrations were derived from optical densities.

Total RNA (5 g in 10.5 L total volume in DEPC-H<sub>2</sub>O) was heated at 65°C for 5 minutes, and then added to tubes containing 4 I 5X MMLV-RT buffer, 2 I dNTPs, 2 I dT17 primer (10 pmol/ml), 0.5 I RNAsin (40 U/ml) and 1 I MMLV-RT (200 units/I). The reactions were incubated at 37°C for 1 hour. Afterward, the MMLV-RT was inactivated by heating at 95°C for 5 minutes. The samples were diluted by addition of 80 L water.

Transcribed samples (5 I) were subjected to polymerase-chain reaction using standard methodologies (50 I total volume). Boden *et al.*, Endocrinology, 138:2820-28 (1997); Ausubel *et al.*, "Quantitation of rare DNAs by the polymerase chain reaction", *in* Current Protocols in Molecular Biology, chapter 15.31-1, Wiley & Sons, Trenton, NJ (1990). Briefly, samples were added to tubes containing water and appropriate amounts of PCR buffer (25 mM MgC1<sub>2</sub>, dNTPs, forward and reverse primers (for RLMPU; SEQ ID NOs: 15 and 16), <sup>32</sup>P-dCTP, and DNA polymerase. Primers were designed to run consistently at

30

10

15

20

22 cycles for radioactive band detection and 33 cycles for amplification of PCR product for use as a screening probe (94°C, 30 sec, 58°C, 30 sec; 72°C, 20 sec).

Sequencing of the agarose gel-purified MG63 osteosarcoma-derived PCR product gave a sequence more than 95% homologous to the RLMPU PCR product. That sequence is designated HLMP unique region (HLMPU; SEQ ID NO: 6).

# EXAMPLE 17: Screening of reverse-transcriptase-derived MG63 cDNA

10

15

5

Screening was performed with PCR using specific primers (SEQ ID NOs: 16 and 17) as described in Example 7. A 717 base-pair MG63 PCR product was agarose gel purified and sequenced with the given primers (SEQ. ID NOs: 12, 15, 16, 17, 18, 27 and 28). Sequences were confirmed a minimum of two times in both directions. The MG63 sequences were aligned against each other and then against the full-length rat LMP cDNA sequence to obtain a partial human LMP cDNA sequence (SEQ ID NO: 7).

# EXAMPLE 18: Screening of a Human Heart cDNA Library

20

25

30

Based on Northern blot experiments, it was determined that LMP-1 is expressed at different levels by several different tissues, including human heart muscle. A human heart cDNA library was therefore examined. The library was plated at 5 x 10<sup>4</sup> pfu/ml onto agar plates (LB bottom agar) and plates were grown overnight at 37° C. Filter membranes were overlaid onto the plates for two minutes. Afterward, the filters denatured, rinsed, dried, UV cross-linked and incubated in pre-hyridization buffer (2X PIPES [pH 6.5]; 5% formamide, 1% SDS, 100 g/ml denatured salmon sperm DNA) for 2 h at 42°C. A radiolabelled, LMP-unique, 223 base-pair probe (<sup>32</sup>P, random primer labelling; SEQ ID NO: 6) was added and hybridized for 18 h at 42°C. Following hybridization, the membranes were washed once at room temperature (10 min,

1 x SSC, 0.1% SDS) and three times at 55°C (15 min, 0.1 x SSC, 0.1% SDS). Double-positive plaque-purified heart library clones, identified by autoradiography, were rescued as lambda phagemids according to the manufacturers' protocols (Stratagene, La Jolla, CA).

Restriction digests of positive clones yielded cDNA inserts of varying sizes. Inserts greater than 600 base-pairs in length were selected for initial screening by sequencing. Those inserts were sequenced by standard methods as described in Example 11.

One clone, number 7, was also subjected to automated sequence analysis using primers corresponding to SEQ ID NOs: 11-14, 16 and 27. The sequences obtained by these methods were routinely 97-100% homologous. Clone 7 (Partial Human LMP-1 cDNA from a heart library; SEQ. ID NO: 8) contained sequence that was more than 87% homologous to the rat LMP cDNA sequence in the translated region.

15

20

25

30

10

5

## EXAMPLE 19: Determination of Full-Length Human LMP-1 cDNA

Overlapping regions of the MG63 human osteosarcoma cell cDNA sequence and the human heart cDNA clone 7 sequence were used to align those two sequences and derive a complete human cDNA sequence of 1644 base-pairs. NALIGN, a program in the PCGENE software package, was used to align the two sequences. The overlapping regions of the two sequences constituted approximately 360 base-pairs having complete homology except for a single nucleotide substitution at nucleotide 672 in the MG63 cDNA (SEQ ID NO: 7) with clone 7 having an "A" instead of a "G" at the corresponding nucleotide 516 (SEQ ID NO: 8).

The two aligned sequences were joined using SEQIN, another subprogram of PCGENE, using the "G" substitution of the MG63 osteosarcoma cDNA clone. The resulting sequence is shown in SEQ ID NO: 9. Alignment of the novel human-derived sequence with the rat LMP-1 cDNA was accomplished with NALIGN. The full-length human LMP-1 cDNA sequence

(SEQ. ID NO: 9) is 87.3% homologous to the translated portion of rat LMP-1 cDNA sequence.

# EXAMPLE 20: Determination of Amino Acid Sequence of Human LMP-1

5

10

15

20

25

30

The putative amino acid sequence of human LMP-1 was determined with the PCGENE subprogram TRANSL. The open reading frame in SEQ ID NO: 9 encodes a protein comprising 457 amino acids (SEQ. ID NO: 10). Using the PCGENE subprogram Palign, the human LMP-1 amino acid sequence was found to be 94.1% homologous to the rat LMP-1 amino acid sequence.

# EXAMPLE 21: Determination of the 5 Prime Untranslated Region of the Human LMP cDNA

MG63 5' cDNA was amplified by nested RT-PCR of MG63 total RNA using a 5' rapid amplification of cDNA ends (5' RACE) protocol. This method included first strand cDNA synthesis using a lock-docking oligo (dT) primer with two degenerate nucleotide positions at the 3' end (Chenchik *et al.*, CLONTECHniques, X:5 (1995); Borson *et al.*, PC Methods Applic., 2:144 (1993)). Second-strand synthesis is performed according to the method of Gubler *et al.*, Gene, 25:263 (1983), with a cocktail of *Escherichia coli* DNA polymerase I, RNase H, and *E. coli* DNA ligase. After creation of blunt ends with T4 DNA polymerase, double-stranded cDNA was ligated to the fragment (5'-CTAATACGACTCACTATAGGGCTCGAGCGGCCGCCCGGGCAGGT-3') (SEQ.ID NO: 19). Prior to RACE, the adaptor-ligated cDNA was diluted to a concentration suitable for Marathon RACE reactions (1:50). Adaptor-ligated double-stranded cDNA was then ready to be specifically cloned.

First-round PCR was performed with the adaptor-specific oligonucleotide, 5'-CCATCCTAATACGACTCACTATAGGGC- 3' (AP1) (SEQ.ID NO: 20) as sense primer and a Gene Specific Primer (GSP) from the unique region described in Example 16 (HLMPU). The second round of PCR was

performed using a nested primers GSP1-HLMPU (antisense/reverse primer) (SEQ. ID NO: 23) and GSP2-HLMPUF (SEQ. ID NO: 24) (see Example 16; sense/forward primer). PCR was performed using a commercial kit (Advantage cDNA PCR core kit; CloneTech Laboratories Inc., Palo Alto, CA) that utilizes an antibody-mediated, but otherwise standard, hot-start protocol. PCR conditions for MG63 cDNA included an initial hot-start denaturation (94°C, 60 sec) followed by: 94°C, 30 sec; 60°C, 30 sec; 68°C, 4 min; 30 cycles. The first-round PCR product was approximately 750 base-pairs in length whereas the nested PCR product was approximately 230 base-pairs. The first-round PCR product was cloned into linearized pCR 2.1 vector (3.9 Kb). The inserts were sequenced in both directions using M13 Forward and Reverse primers (SEQ. ID NO: 11; SEQ. ID NO: 12)

EXAMPLE 22: Determination of Full-length Human LMP-1 cDNA with 5 Prime

UTR

Overlapping MG63 human osteosarcoma cell cDNA 5'-UTR sequence (SEQ ID NO: 21), MG63 717 base-pair sequence (Example 17; SEQ ID NO: 8) and human heart cDNA clone 7 sequence (Example 18) were aligned to derive a novel human cDNA sequence of 1704 base-pairs (SEQ.ID NO: 22). The alignment was accomplished with NALIGN, (both PCGENE and Omiga 1.0; Intelligenetics). Over-lapping sequences constituted nearly the entire 717 base-pair region (Example 17) with 100% homology. Joining of the aligned sequences was accomplished with SEQIN.

25

30

5

10

15

20

#### **EXAMPLE 23: Construction of LIM Protein Expression Vector**

The construction of pHIS-5ATG LMP-1s expression vector was carried out with the sequences described in Examples 17 and 18. The 717 base-pair clone (Example 17; SEQ ID NO: 7) was digested with <u>Clal</u> and <u>EcoRV</u>. A small fragment (~250 base-pairs) was gel purified. Clone 7 (Example 18; SEQ

ID NO: 8) was digested with <u>Clal</u> and <u>Xbal</u> and a 1400 base-pair fragment was gel purified. The isolated 250 base-pair and 1400 base-pair restriction fragments were ligated to form a fragment of ~1650 base-pairs.

Due to the single nucleotide substitution in Clone 7 (relative to the 717 base-pair PCR sequence and the original rat sequence) a stop codon at translated base-pair 672 resulted. Because of this stop codon, a truncated (short) protein was encoded, hence the name LMP-1s. This was the construct used in the expression vector (SEQ ID NO: 32). The full length cDNA sequence with 5' UTR (SEQ ID NO: 33) was created by alignment of SEQ ID NO: 32 with the 5' RACE sequence (SEQ ID NO: 21). The amino acid sequence of LMP-1s (SEQ ID NO: 34) was then deduced as a 223 amino acid protein and confirmed by Western blot (as in Example 15) to run at the predicted molecular weight of ~ 23.7 kD.

The pHis-ATG vector (InVitrogen, Carlsbad, CA) was digested with EcoRV and Xbal. The vector was recovered and the1650 base-pair restriction fragment was then ligated into the linearized pHis-ATG. The ligated product was cloned and amplified. The pHis-ATG-LMP-1s Expression vector, also designated pHIS-A with insert HLMP-1s, was purified by standard methods.

20 EXAMPLE 24: Induction of Bone Nodule Formation and Mineralization *In vitro* with LMP Expression Vector

Rat Calvarial cells were isolated and grown in secondary culture according to Example 1. Cultures were either unstimulated or stimulated with glucocorticoid (GC) as described in Example 1. A modification of the Superfect Reagent (Qiagen, Valencia, CA) transfection protocol was used to transfect 3 g/well of each vector into secondary rat calvarial osteoblast cultures according to Example 25.

Mineralized nodules were visualized by Von Kossa staining, as described in Example 3. Human LMP-1s gene product overexpression alone induced bone nodule formation (~203 nodules/well) in vitro. Levels of nodules were

5

10

15

25

approximately 50% of those induced by the GC positive control (~412 nodules/well). Other positive controls included the pHisA-LMP-Rat expression vector (~152 nodules/well) and the pCMV2/LMP-Rat-Fwd Expression vector (~206 nodules/well), whereas the negative controls included the pCMV2/LMP-Rat-Rev. Expression vector (~2 nodules/well) and untreated (NT) plates (~4 nodules/well). These data demonstrate that the human cDNA was at least as osteoinductive as the rat cDNA. The effect was less than that observed with GC stimulation, most likely due to suboptimal doses of Expression vector.

10

15

20

25

30

#### EXAMPLE 25: LMP-Induced Cell Differentiation In Vitro and In Vivo

The rat LMP cDNA in clone 10-4 (see Example 12) was excised from the vector by double-digesting the clone with Notl and Apal overnight at 37°C. Vector pCMV2 MCS (InVitrogen, Carlsbad, CA) was digested with the same restriction enzymes. Both the linear cDNA fragment from clone 10-4 and pCMV2 were gel purified, extracted and ligated with T4 ligase. The ligated DNA was gel purified, extracted and used to transform *E. coli* JM109 cells for amplification. Positive agar colonies were picked, digested with Notl and Apal and the restriction digests were examined by gel electrophoresis. Stock cultures were prepared of positive clones.

A reverse vector was prepared in analogous fashion except that the restriction enzymes used were Xbal and HindIII. Because these restriction enzymes were used, the LMP cDNA fragment from clone 10-4 was inserted into pRc/CMV2 in the reverse (that is, non-translatable) orientation. The recombinant vector produced is designated pCMV2/RLMP.

An appropriate volume of pCMV10-4 (60 nM final concentration is optimal [3 g]; for this experiment a range of 0-600 nM/well [0-30 g/well] final concentration is preferred) was resuspended in Minimal Eagle Media (MEM) to 450 I final volume and vortexed for 10 seconds. Superfect was added (7.5 I/ml final solution), the solution was vortexed for 10 seconds and then incubated at

room termperature for 10 minutes. Following this incubation, MEM supplemented with 10% FBS (1 ml/well; 6 ml/plate) was added and mixed by pipetting.

The resulting solution was then promptly pipetted (1 ml/well) onto washed ROB cultures. The cultures were incubated for 2 hours at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub>. Afterward, the cells were gently washed once with sterile PBS and the appropriate normal incubation medium was added.

Results demonstrated significant bone nodule formation in all rat cell cultures which were induced with pCMV10-4. For example, pCMV10-4 transfected cells produced 429 nodules/well. Positive control cultures, which were exposed to Trm, produced 460 nodules/well. In contrast, negative controls, which received no treatment, produced 1 nodule/well. Similarly, when cultures were transfected with pCMV10-4 (reverse), no nodules were observed.

For demonstrating *de novo* bone formation *in vivo*, marrow was aspirated from the hindlimbs of 4-5 week old normal rats (rnu/+; heterozygous for recessive athymic condition). The aspirated marrow cells were washed in alpha MEM, centrifuged, and RBCs were lysed by resuspending the pellet in 0.83% NH<sub>4</sub>Cl in 10 mM Tris (pH 7.4). The remaining marrow cells were washed 3x with MEM and transfected for 2 hours with 9 g of pCMV-LMP-1s (forward or reverse orientation) per 3 x 10<sup>6</sup> cells. The transfected cells were then washed 2X with MEM and resuspended at a concentration of 3 x 10<sup>7</sup> cells/ml.

The cell suspension (100 I) was applied via sterile pipette to a sterile 2 x 5 mm type I bovine collagen disc (Sulzer Orthopaedics, Wheat Ridge, CO). The discs were surgically implanted subcutaneously on the skull, chest, abdomen or dorsal spine of 4-5 week old athymic rats (rnu/rnu). The animals were scarified at 3-4 weeks, at which time the discs or surgical areas were excised and fixed in 70% ethanol. The fixed specimens were analyzed by radiography and undecalcified histologic examination was performed on 5 m

5

10

15

20

25

thick sections stained with Goldner Trichrome. Experiments were also performed using devitalized (guanidine extracted) demineralized bone matrix (Osteotech, Shrewsbury, NJ) in place of collagen discs.

Radiography revealed a high level of mineralized bone formation that conformed to the form of the original collagen disc containing LMP-1s transfected marrow cells. No mineralized bone formation was observed in the negative control (cells transfected with a reverse-oriented version of the LMP-1s cDNA that did not code for a translated protein), and absorption of the carrier appeared to be well underway.

Histology revealed new bone trabeculae lined with osteroblasts in the LMP-1s transfected implants. No bone was seen along with partial resorption of the carrier in the negative controls.

Radiography of a further experiment in which 18-sets (9 negative control pCMV-LMP-REV & 9 experimental pCMV-LMP-1s) of implants were added to sites alternating between lumbar and thoracic spine in athymic rats demonstrated 0/9 negative control implants exhibiting bone formation (spine fusion) between vertebrae. All nine of the pCMV-LMP-1s treated implants exhibited solid bone fusions between vertebrae.

EXAMPLE 26: The Synthesis of pHIS-5' ATG LMP-1s Expression Vector from the sequences Demonstrated in Examples 2 and 3.

The 717 base-pair clone (Example 17) was digested with <u>Clal</u> and <u>Eco</u>RV (New England Biologicals, city, MA). A small fragment (~250 base-pairs) was gel purified. Clone No. 7 (Example 18) was digested with <u>Clal</u> and <u>Xbal</u>. A 1400 base-pair fragment was gel purified from that digest. The isolated 250 base-pair and 1400 base-pair cDNA fragments were ligated by standard methods to form a fragment of ~1650 bp. The pHis-A vector (InVitrogen) was digested with <u>Eco</u>RV and *Xbal*. The linearized vector was recovered and ligated to the chimeric 1650 base-pair cDNA fragment. The liqated product was cloned and amplified by standard methods, and the pHis-

5

10

15

20

25

A-5' ATG LMP-1s expression vector, also denominated as the vector pHis-A with insert HLMP-1s, was deposited at the ATCC as previously described.

EXAMPLE 27: The Induction of Bone Nodule Formation and Mineralization In

Vitro With pHis-5' ATG LMP-1s Expression Vector

Rat calvarial cells were isolated and grown in secondary culture according to Example 1. Cultures were either unstimulated or stimulated with glucocorticoid (GC) according to Example 1. The cultures were transfected with 3 g of recombinant pHis-A vector DNA/well as described in Example 25. Mineralized nodules were visualized by Von Kossa staining according to Example 3.

Human LMP-1s gene product overexpression alone (*i.e.*, without GC stimulation) induced significant bone nodule formation (~203 nodules/well) *in vitro*. This is approximately 50% of the amount of nodules produced by cells exposed to the GC positive control (~412 nodules/well). Similar results were obtained with cultures transfected with pHisA-LMP-Rat Expression vector (~152 nodules/well) and pCMV2/LMP-Rat-Fwd (~206 nodules/well). In contrast, the negative control pCMV2/LMP-Rat-Rev yielded (~2 nodules/well), while approximately 4 nodules/well were seen in the untreated plates. These data demonstrate that the human LMP-1 cDNA was at least as osteoinductive as the rat LMP-1 cDNA in this model system. The effect in this experiment was less than that observed with GC stimulation; but in some the effect was comparable.

25

30

5

10

15

20

#### EXAMPLE 28: LMP Induces Secretion of a Soluble Osteoinductive Factor

Overexpression of RLMP-1 or HLMP-1s in rat calvarial osteoblast cultures as described in Example 24 resulted in significantly greater nodule formation than was observed in the negative control. To study the mechanism of action of LIM mineralization protein conditioned medium was harvested at

different time points, concentrated to 10 X, sterile filtered, diluted to its original concentration in medium containing fresh serum, and applied for four days to untransfected cells.

Conditioned media harvested from cells transfected with RLMP-1 or HLMP-1s at day 4 was approximately as effective in inducing nodule formation as direct overexpression of RLMP-1 in transfected cells. Conditioned media from cells transfected with RLMP-1 or HLMP-1 in the reverse orientation had no apparent effect on nodule formation. Nor did conditioned media harvested from LMP-1 transfected cultures before day 4 induce nodule formation. These data suggest that expression of LMP-1 caused the synthesis and/or secretion of a soluble factor, which did not appear in culture medium in effectie amounts until 4 days post transfection.

Since overexpression of rLMP-1 resulted in the secretion of an osteoinductive factor into the medium, Western blot analysis was used to determine if LMP-1 protein was present in the medium. The presence of rLMP-1 protein was assessed using antibody specific for LMP-1 (QDPDEE) and detected by conventional means. LMP-1 protein was found only in the cell layer of the culture and not detected in the medium.

Partial purification of the osteoinductive soluble factor was accomplished by standard 25% and 100% ammonium sulfate cuts followed by DE-52 anion exchange batch chromatography (100 mM or 500 mM NaCl). All activity was observed in the high ammonium sulfate, high NaCl fractions. Such localization is consistent with the possibility of a single factor being responsible for conditioning the medium.

25

30

20

5

10

15

EXAMPLE 29: Gene Therapy In Lumbar Spine Fusion Mediated by Low Dose Adenovirus

This study determined the optimal dose of adenoviral delivery of the LMP-1 cDNA (SEQ ID NO: 2) to promote spine fusion in normal, that is, immune competent, rabbits.

A replication-deficient human recombinant adenovirus was constructed with the LMP-1 cDNA (SEQ ID NO: 2) driven by a CMV promoter using the Adeno-Quest™ Kit (Quantum Biotechnologies, Inc., Montreal). A commercially available (Quantum Biotechnologies, Inc., Montreal) recombinant adenovirus containing the beta-galactosidase gene was used as a control.

Initially, an *in vitro* dose response experiment was performed to determine the optimal concentration of adenovirus-delivered LMP-1 ("AdV-LMP-1") to induce bone differentiation in rat calvarial osteoblast cultures using a 60-minute transduction with a multiplicity of infection ("MOI") of 0.025, 0.25, 2.5, or 25 plaque-forming units (pfu) of virus per cell. Positive control cultures were differentiated by a 7-day exposure to 10<sup>9</sup> M glucocorticoid ("GC"). Negative control cultures were left untreated. On-day 14, the number of mineralized bone nodules was counted after von Kossa staining of the cultures, and the level of osteocalcin secreted into the medium (pmol/mL) was measured by radioimmunoassay (mean ± SEM).

The results of this experiment are shown in Table I. Essentially no spontaneous nodules formed in the untreated negative control cultures. The data show that a MOI equal to 0.25 pfu/cell is most effective for osteoinducing bone nodules, achieving a level comparable to the positive control (GC).

20 Lower and higher doses of adenovirus were less effective.

TABLE I

Outcome				AdV-LMF	P-1 Dose (I	MOI)
	Neg. Ctrl.	GC	0.025	0.25	2.5	25
Bone Nodules	0.5±0.2	188±35	79.8±13	145.1±1 3	26.4±15	87.6±2
Osteocalcin	1.0±0.1	57.8±9	28.6±11	22.8±1	18.3±3	26.0±2

5

10

In vivo experiments were then performed to determine if the optimal in vitro dose was capable of promoting intertransverse process spine fusions in skeletally mature New Zealand white rabbits. Nine rabbits were anesthetized and 3 cc of bone marrow was aspirated from the distal femur through the intercondylar notch using an 18 gauge needle. The buffy coat was then isolated, a 10-minute transduction with AdV-LMP-1 was performed, and the cells were returned to the operating room for implantation. Single level posterolateral lumbar spine arthrodesis was performed with decortication of transverse processes and insertion of carrier (either rabbit devitalized bone matrix or a collagen sponge) containing 8-15 million autologous nucleated buffy coat cells transduced with either AdV-LMP-1 (MOI = 0.4) or AdV-BGal (MOI = 0.4). Rabbits were euthanized after 5 weeks and spine fusions were assessed by manual palpation, plain x-rays, CT scans, and undecalcified histology.

The spine fusion sites that received AdV-LMP-1 induced solid, continuous spine fusion masses in all nine rabbits. In contrast, the sites receiving AdV-BGal, or a lower dose of AdV-LMP-1 (MOI = 0.04) made little or no bone and resulted in spine fusion at a rate comparable to the carrier alone (< 40%). These results were consistent as evaluated by manual palpation, CT scan, and histology. Plain radiographs, however, sometimes overestimated the amount of bone that was present, especially in the control sites. LMP-1 cDNA delivery and bone induction was successful with both of the carrier materials tested. There was no evidence of systemic or local immune response to the adenovirus vector.

These data demonstrate consistent bone induction in a previously validated rabbit spine fusion model which is quite challenging. Furthermore, the protocol of using autogenous bone marrow cells with intraoperative *ex vivo* gene transduction (10 minutes) is a more clinically feasible procedure than other methods that call for overnight transduction or cell expansion for weeks in culture. In addition, the most effective dose of recombinant adenovirus (MOI=0.25) was substantially lower than doses reported in other gene therapy

10

15

20

25

applications (MOI-40-500). We believe this is due to the fact that LMP-1 is an intracellular signaling molecule and may have powerful signal amplification cascades. Moreover, the observation that the same concentration of AdV-LMP-1 that induced bone in cell culture was effective *in vivo* was also surprising given the usual required increase in dose of other growth factors when translating from cell culture to animal experiments. Taken together, these observations indicate that local gene therapy using adenovirus to deliver the LMP-1 cDNA is possible and the low dose required will likely minimize the negative effects of immune response to the adenovirus vector.

10

15

20

25

30

5

# EXAMPLE 30: Use of Peripheral Venous Blood Nucleated Cells (Buffy Coat) for Gene Therapy With LMP-1 cDNA To Make Bone

In four rabbits we performed spine fusion surgery as above (Example 29) except the transduced cells were the buffy coat from venous blood rather than bone marrow. These cells were transfected with Adeno-LMP or pHIS-LMP plasmid and had equivalent successful results as when bone marrow cells were used. This discovery of using ordinary venous blood cells for gene delivery makes gene therapy more feasible clinically since it avoids painful marrow harvest under general anesthesia and yields two times more cells per mL of starting material.

#### EXAMPLE 31: Isolation of Human LMP-1 Splice Variants

Intron/Exon mRNA transcript splice variants are a relatively common regulatory mechanism in signal transduction and cellular/tissue development. Splice variants of various genes have been shown to alter protein-protein, protein-DNA, protein-RNA, and protein-substrate interactions. Splice variants may also control tissue specificity for gene expression allowing different forms (and therefore functions) to be expressed in various tissues. Splice variants are a common regulatory phenomenon in cells. It is possible that the LMP

splice variants may result in effects in other tissues such as nerve regeneration, muscle regeneration, or development of other tissues.

To screen a human heart cDNA library for splice variants of the HLMP-1 sequence, a pair of PCR primer corresponding to sections of SEQ ID NO: 22 was prepared. The forward PCR primer, which was synthesized using standard techniques, corresponds to nucleotides 35-54 of SEQ ID NO: 22. It has the following sequence:

5' GAGCCGGCATCATGGATTCC 3' (SEQ ID NO: 35)

10

The reverse PCR primer, which is the reverse complement of nucleotides 820-839 in SEQ ID NO: 22, has the following sequence:

5' GCTGCCTGCACAATGGAGGT 3' (SEQ ID NO: 36)

15

20

25

The forward and reverse PCR primers were used to screen human heart cDNA (ClonTech, Cat No. 7404-1) for sequences similar to HLMP-1 by standard techniques, using a cycling protocol of 94°C for 30 seconds, 64°C for 30 seconds, and 72°C for 1 minute, repeated 30 times and followed by a 10 minute incubation at 72°C. The amplification cDNA sequences were gel-purified and submitted to the Emory DNA Sequence Core Facility for sequencing. The clones were sequenced using standard techniques and the sequences were examined with PCGENE (Intelligenetics; Programs SEQUIN and NALIGN) to determine homology to SEQ ID NO: 22. Two homologous nucleotide sequences with putative alternative splice sites compared to SEQ ID NO: 22 were then translated to their respective protein products with Intelligenetic's program TRANSL.

One of these two novel human cDNA sequences (SEQ ID NO: 37) comprises 1456 bp:

30

10 20 30 40 50 60

	CGACGCAGAG	CAGCGCCCTG	GCCGGGCCAA	GCAGGAGCCG	GCATCATGGA	TTCCTTCAAG
	70	. 80	90	100	110	120
	GTAGTGCTGG	AGGGCCAGC	ACCTTGGGGC	TTCCGGCTGC	AAGGGGGCAA	GGACTTCAAT
	130	140	150	160	170	180
5	GTGCCCCTCT	CCATTTCCCG	GCTCACTCCT	GGGGGCAAAG	CGGCGCAGGC	CGGAGTGGCC
	190	200	210	220	230	240
	GTGGGTGACT	GGGTGCTGAG	CATCGATGGC	GAGAATGCGG	GTAGCCTCAC	ACACATCGAA
	250	260	270	280	290	300
	GCTCAGAACA	AGATCCGGGC	CTGCGGGGAG	CGCCTCAGCC	TGGGCCTCAG	CAGGGCCCAG
10	310	320	* 330	340	* 350	360
	CCGGTTCAGA	GCAAACCGCA	GAAG <u>GTGCAG</u>	ACCCCTGACA	<u>A</u> ACAGCCGCT	CCGACCGCTG
-	370	380	390	400	410	420
	GTCCCAGATG	CCAGCAAGCA	GCGGCTGATG	GAGAACACAG	AGGACTGGCG	GCCGCGGCCG
	430	440	450	460	470	480
15	GGGACAGGCC	AGTCGCGTTC	CTTCCGCATC	CTTGCCCACC	TCACAGGCAC	CGAGTTCATG
	490	500	510	520	530	540
	CAAGACCCGG	ATGAGGAGCA	CCTGAAGAAA	TCAAGCCAGG	TGCCCAGGAC	AGAAGCCCCA
	550	560	570	580	590	600
	GCCCCAGCCT	CATCTACACC	CCAGGAGCCC	TGGCCTGGCC	CTACCGCCCC	CAGCCCTACC
20	610	620	630	640	650	660
	AGCCGCCCGC	CCTGGGCTGT	GGACCCTGCG	TTTGCCGAGC	GCTATGCCCC	GGACAAAACG
	670	680	690	700	710	720
	AGCACAGTGC	TGACCCGGCA	CAGCCAGCCG	GCCACGCCCA	CGCCGCTGCA	GAGCCGCACC
	730	740	750	760	770	780
25	TCCATTGTGC	AGGCAGCTGC	CGGAGGGGTG	CCAGGAGGGG	GCAGCAACAA	CGGCAAGACT
	790	800	810	820	830	840
	CCCGTGTGTC	ACCAGTGCCA	CAAGGTCATC	CGGGGCCGCT	ACCTGGTGGC	GTTGGGCCAC
	850	860	870	880	890	900

	GCGTACCACC	CGGAGGAGTT	TGTGTGTAGC	CAGTGTGGGA	AGGTCCTGGA	AGAGGGTGGC
	910	920	· 930	940	950	960
	TTCTTTGAGG	AGAAGGGCGC	CATCTTCTGC	CCACCATGCT	ATGACGTGCG	CTATGCACCC
	970	980	990	1000	1010	1020
5	AGCTGTGCCA	AGTGCAAGAA	GAAGATTACA	GGCGAGATCA	TGCACGCCCT	GAAGATGACC
	1030	1040	1050	1060	1070	1080
	TGGCACGTGC	ACTGCTTTAC	CTGTGCTGCC	TGCAAGACGC	CCATCCGGAA	CAGGGCCTTC
	1090	1100	1110	1120	1130	1140
	TACATGGAGG	AGGGCGTGCC	CTATTGCGAG	CGAGACTATG	AGAAGATGTT	TGGCACGAAA
10	1150	1160	1170	1180	1190	1200
	TGCCATGGCT	GTGACTTCAA	GATCGACGCT	GGGGACCGCT	TCCTGGAGGC	CCTGGGCTTC
	1210	1220	1230	1240	1250	1260
	AGCTGGCATG	ACACCTGCTT	CGTCTGTGCG	ATATGTCAGA	TCAACCTGGA	AGGAAAGACC
	1270	1280	1290	1300	1310	1320
15	TTCTACTCCA	AGAAGGACAG	GCCTCTCTGC	AAGAGCCATG	CCTTCTCTCA	TGTGTGAGCC
	1330	1340	1350	1360	1370	1380
	CCTTCTGCCC	ACAGCTGCCG	CGGTGGCCCC	TAGCCTGAGG	GGCCTGGAGT	CGTGGCCCTG
	1390	1400	1410	1420	1430	1440
	CATTTCTGGG	TAGGGCTGGC	AATGGTTGCC	TTAACCCTGG	CTCCTGGCCC	GAGCCTGGGC
20	1450					
	TCCCGGGCCC	TGCCCA				

Reading frame shifts caused by the deletion of a 119 bp fragment (between \*) and the addition of a 17 bp fragment (<u>underlined</u>) results in a truncated gene product having the following derived amino acid sequence (SEQ ID NO: 38):

10 20 30 40 50 60

	MDSFKVVLEG	PAPWGFRLQG	GKDFNVPLSI	SRLTPGGKAA	QAGVAVGDWV	LSIDGENAGS
	. 70	80	90	100	110	120
	LTHIEAQNKI	RACGERLSLG	LSRAQPVQNK	PQK <u>VQTPDK</u> Q	PLRPLVPDAS	KQRLMENTED
	130	140	150	160	170	180
5	WRPRPGTGQS	RSFRILAHLT	GTEFMQDPDE	EHLKKSSQVP	RTEAPAPASS	TPQEPWPGPT
	190	200	210	220	230	240
	APSPTSRPPW	AVDPAFAERY	APDKTSTVLT	RHSQPATPTP	LQSRTSIVQA	AAGGVPGGGS
	250	260	270	280	290	300
	NNGKTPVCHQ	CHQVIRARYL	VALGHAYHPE	EFVCSQCGKV	LEEGGFFEEK	GAIFCPPCYD
10	310	320	330	340	350	360
	VRYAPSCAKC	KKKITGEIMH				VPYCERDYEK
-	370			400		420
	MFGTKCQWCD	FKIDAGDRFL	EALGFSWHDT	CFVCAICQIN	LEGKTFYSKK	DRPLCKSHAF
	SHV					

This 423 amino acid protein demonstrates 100% homology to the protein shown in Sequence ID No. 10, except for the sequence in the highlighted area (amino acids 94-99), which are due to the nucleotide changes depicted above.

The second novel human heart cDNA sequence (SEQ ID NO: 39) comprises 1575 bp:

	10	20	30	40	50	60
	CGACGCAGAG	CAGCGCCCTG	GCCGGGCCAA	GCAGGAGCCG	GCATCATGGA	TTCCTTCAAG
	70	80	90	100	110	120
25	GTAGTGCTGG	AGGGCCAGC	ACCTTGGGGC	TTCCGGCTGC	AAGGGGGCAA	GGACTTCAAT
	130	140	150	160	170	180
	GTGCCCCTCT	CCATTTCCCG	GCTCACTCCT	GGGGCAAAG	CGGCGCAGGC	CGGAGTGGCC
	190	200	210	220	230	240

	GTGGGTGACT	GGGTGCTGAG	CATCGATGGC	GAGAATGCGG	GTAGCCTCAC	ACACATCGAA
	250	260	270	280	290	300
	GCTCAGAACA	AGATCCGGGC	CTGCGGGGAG	CGCCTCAGCC	TGGGCCTCAG	CAGGGCCCAG
	310	320	330	340	350	360
5	CCGGTTCAGA	GCAAACCGCA	GAAGGCCTCC	GCCCCGCCG	CGGACCCTCC	GCGGTACACC
	370	380	390	400	410	420
	TTTGCACCCA	GCGTCTCCCT	CAACAAGACG	GCCCGGCCCT	TTGGGGCGCC	CCCGCCCGCT
	430	440	450	460	470	480
	GACAGCGCCC	CGCAACAGAA	TGG <u>GTGCAGA</u>	CCCCTGACAA	ACAGCCGCTC	CGACCGCTGG
10	490	500	510	520	530	540
	TCCCAGATGC	CAGCAAGCAG	CGGC <u>TGA</u> TGG	AGAACACAGA	GGACTGGCGG	CCGCGGCCGG
	550	560	570	580	590	600
	GGACAGGCCA	GTCGCGTTCC	TTCCGCATCC	TTGCCCACCT	CACAGGCACC	GAGTTCATGC
	610	620	630	640	650	660
15	AAGACCCGGA	TGAGGAGCAC	CTGAAGAAAT	CAAGCCAGGT	GCCCAGGACA	GAAGCCCCAG
	670	680	690	700	710	720
	CCCCAGCCTC	ATCTACACCC	CAGGAGCCCT	GGCCTGGCCC	TACCGCCCCC	AGCCCTACCA
	730	740	750	760	770	780
	GCCGCCCGCC	CTGGGCTGTG	GACCCTGCGT	TTGCCGAGCG	CTATGCCCCG	GACAAAACGA
20	790	800	810	820	830	840
	GCACAGTGCT	GACCCGGCAC	AGCCAGCCGG	CCACGCCCAC	GCCGCTGCAG	AGCCGCACCT
	850	860	870	880	890	900
	CCATTGTGCA	GGCAGCTGCC	GGAGGGGTGC	CAGGAGGGGG	CAGCAACAAC	GGCAAGACTC
	910	920	930	940	950	960
25	CCGTGTGTCA	CCAGTGCCAC	AAGGTCATCC	GGGGCCGCTA	CCTGGTGGCG	TTGGGCCACG
	970	980	990	1000	1010	1020
	CGTACCACCC	GGAGGAGTTT	GTGTGTAGCC	AGTGTGGGAA	GGTCCTGGAA	GAGGGTGGCT
	1030	1040	1050	1060	1070	1080

	TCTTTGAGGA	GAAGGGCGCC	ATCTTCTGCC	CACCATGCTA	TGACGTGCGC	TATGCACCCA
	1090	1100	1110	1120	1130	1140
	GCTGTGCCAA	GTGCAAGAAG	AAGATTACAG	GCGAGATCAT	GCACGCCCTG	AAGATGACCT
	1150	1160	1170	1180	1190	1200
5	GGCACGTGCA	CTGCTTTACC	TGTGCTGCCT	GCAAGACGCC	CATCCGGAAC	AGGGCCTTCT
	1210	1220	1230	1240	1250	1260
	ACATGGAGGA	GGGCGTGCCC	TATTGCGAGC	GAGACTATGA	GAAGATGTTT	GGCACGAAAT
	1270	1280	1290	1300	1310	1320
	GCCATGGCTG	TGACTTCAAG	ATCGACGCTG	GGGACCGCTT	CCTGGAGGCC	CTGGGCTTCA
10	1330	1340	1350	1360	1370	1380
	GCTGGCATGA	CACCTGCTTC	GTCTGTGCGA	TATGTCAGAT	CAACCTGGAA	GGAAAGACCT
rale a résulto de la composição de la co	1390	1400	1410	1420	1430	1440
	TCTACTCCAA	GAAGGACAGG	CCTCTCTGCA	AGAGCCATGC	CTTCTCTCAT	GTGTGAGCCC
	1450	1460	1470	1480	1490	1500
15	CTTCTGCCCA	CAGCTGCCGC	GGTGGCCCCT	AGCCTGAGGG	GCCTGGAGTC	GTGGCCCTGC
	1510	1520	1530	1540	1550	1560
	ATTTCTGGGT	AGGGCTGGCA	ATGGTTGCCT	TAACCCTGGC	TCCTGGCCCG	AGCCTGGGCT
	1570					

CCCGGGCCCT GCCCA

Reading frame shifts caused by the addition of a 17 bp fragment (bolded, italicized and underlined) results in an early translation stop codon at position 565-567 (underlined).

The derived amino acid sequence (SEQ ID NO: 40) consists of 153 amino acids:

25

60	50	40	30	20	10
LSIDGENAGS	QAGVAVGDWV	SRLTPGGKAA	GKDFNVPLSI	PAPWGFRLQG	MDSFKVVLEG
120	110	100	90	80	70

LTHIEAQNKI RACGERLSLG LSRAQPVQSK PQKASAPAAD PPRYTFAPSV SLNKTARPFG

130 140 150

#### APPPADSAPO ONGCRPLTNS RSDRWSOMPA SSG

This protein demonstrates 100% homology to SEQ ID NO: 10 until amino acid 94, where the addition of the 17 bp fragment depicted in the nucleotide sequence results in a frame shift. Over amino acids 94-153, the protein is not homologous to SEQ ID NO: 10. Amino acids 154-457 in SEQ ID NO: 10 are not present due to the early stop codon depicted in nucleotide sequence.

## EXAMPLE 32: Genomic HLMP-1 Nucleotide Sequence

Applicants have identified the genomic DNA sequence encoding HLMP-1, including putative regulatory elements associated with HLMP-1 expression. The entire genomic sequence is shown in SEQ ID. NO: 41. This sequence was derived from AC023788 (clone RP11-564G9), Genome Sequencing Center, Washington University School of Medicine, St. Louis, MO.

The putative promoter region for HLMP-1 spans nucleotides 2,660-8,733 in SEQ ID NO: 41. This region comprises, among other things, at least ten potential glucocorticoid response elements ("GREs") (nucleotides 6148-6153, 6226-6231, 6247-6252, 6336-6341, 6510-6515, 6552-6557, 6727-6732, 6752-6757, 7738-7743, and 8255-8260), twelve potential Sma-2 homologues to Mothers against Drosophilla decapentaplegic ("SMAD") binding element sites (nucleotides 3569-3575, 4552-4558, 4582-4588, 5226-5232, 6228-6234, 6649-6655, 6725-6731, 6930-6936, 7379-7384, 7738-7742, 8073-8079, and 8378-8384), and three TATA boxes (nucleotides 5910-5913, 6932-6935, and 7380-7383). The three TATA boxes, all of the GREs, and eight of the SMAD binding elements ("SBEs") are grouped in the region spanning nucleotides 5,841-8,733 in SEQ ID NO: 41. These regulatory elements can be used, for example, to regulate expression of exogenous nucleotide sequences encoding proteins involved in the process of bone formation. This would permit systemic administration of therapeutic factors or genes relating to bone formation and repair, as well as factors or genes associated with tissue differentiation and development.

In addition to the putative regulatory elements, 13 exons corresponding to the nucleotide sequence encoding HLMP-1 have been identified. These exons span the following nucleotides in SEQ ID NO: 41:

	Exon 1	8733-8767	
30	Exon 2	9790-9895	
	Exon 3	13635-13787	

5

10

15

20

	Exon 4	13877-13907
	Exon 5	14387-14502
	Exon 6	15161-15297
	Exon 7	15401-15437
5	Exon 8	16483-16545
	Exon 9	16689-16923
	Exon 10	18068-18248
	Exon 11	22117-22240
	Exon 12	22323-22440
10	Exon 13	22575-22911

In HLMP-2 there is another exon (Exon 5A), which spans nucleotides 14887-14904.

All cited publications and patents are hereby incorporated by reference in their entirety.

15 While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be appreciated by one skilled in the art from reading this disclosure that various changes in form and detail can be made without departing from the true scope of the invention.

Applicant's or agent's file		International application ?	
reference number 6148	2-304	PCT/US00/11664	

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism reference on page 17 , line 20	erred to in the description				
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet				
Name of depositary institution American Type Culture Collection					
Address of depositary institution (including postal code and cou 10801 University Blvd.  Manassas, VA 20110-2209					
Date of deposit 14 April 2000	Accession Number PTA-1698				
C. ADDITIONAL INDICATIONS (leave blank if not applicate	ble) This information is continued on an additional sheet				
D. DESIGNATED STATES FOR WHICH INDICATIONS A	ARE MADE (if the indications are not for all designated States)				
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)  The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")					
This sheet was received with the international application	For International Bureau use only  This sheet was received by the International Bureau on:  1 7 JUL 2000				
Authorized officer	Authorized officer				

Form PCT/RO/134 (July 1992)

LegalStar 1997, Form PCTM5

PCT/US00/11664 WO 00/66178

Applicant's or agent's file	:	International application 1
reference number	61482-304	PCT/US00/11664

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microor	ganism referred	to in the description					
on page 17 , lir	ne <u>21</u>						
B. IDENTIFICATION OF DEPOSIT		Further deposits are identified on an additional sheet					
Name of depositary institution American Type Culture Collection							
Address of depositary institution (including postal co 10801 University Blvd. Manassas, VA 20110-2209	ode and country						
Day (day-ii)		Accession Number					
Date of deposit 14 April 2000		PTA-1699					
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet							
		E MADE (if the indications are not for all designated States)					
E. SEPARATE FURNISHING OF INDICATION							
The indications listed below will be submitted to a "Accession Number of Deposit")	the Internation	al Bureau later (specify the general nature of the indications e.g.,					
For receiving Office use only		For International Bureau use only					
This sheet was received with the international a	application	This sheet was received by the International Bureau on:					
Authorized officer	Authorized officer						

Form PCT/RO/134 (July 1992)

-51/2-

#### We claim:

10

15

20

25

 An isolated nucleic acid molecule comprising a SEQ ID NO: 37 or SEQ ID NO: 39.

- An isolated human LMP protein encoded by SEQ ID NO: 37 or SEQ
   ID NO: 39.
  - 3. A vector comprising the isolated nucleic acid molecule of claims 1.
  - 4. A host cell comprising the vector of claim 3.
  - 5. The host cell of claim 4, wherein the host cell is selected from the group consisting of prokaryotic cells, yeast cells and mammalian cells
  - 6. The isolated nucleic acid molecule of claim 1, further comprising a label.
  - 7. A human LIM mineralization protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 38 and SEQ ID NO: 40.
  - 8. A monoclonal antibody specific for a HLMP-2 (SEQ ID NO: 38) or HLMP-3 (SEQ ID NO: 40).
  - 9. A method of inducing bone formation comprising transfecting osteogenic precursor cells or peripheral blood leukocytes with an isolated nucleic acid molecule comprising SEQ ID NO: 39.
    - 10. The method of claim 9, wherein the isolated nucleic acid molecule is in a vector.
      - 11. The method of claim 10, wherein the vector is an expression vector.
      - 12. The method of claim 11, wherein the vector is a plasmid.
      - 13. The method of claim 11, wherein the vector is a virus.
      - 14. The method of claim 13, wherein the virus is an adenovirus
      - 15. The method of claim 13, wherein the virus is a retrovirus.
  - 16. The method of claim 9, wherein the osteogenic precursor cells or peripheral blood leukocytes are transfected *ex vivo*.
- 30 17. The method of claim 9, wherein the osteogenic precursor cells are transfected *in vivo* by direct injection of the isolated nucleic acid molecule.

18. The method of claim 9, wherein the LIM mineralization protein is HLMP-3 (SEQ ID NO: 40).

- 19. A method of fusing a spine, comprising:
- (a) transfecting osteogenic precursor cells or peripheral blood leukocytes with an isolated nucleic acid molecule comprising SEQ ID NO: 39;
  - (b) admixing the transfected osteogenic precursor cells or peripheral blood leukocytes with a matrix; and
- (c) contacting the matrix with the spine;
   wherein expression of the nucleotide sequence causes mineralized
   bone formation in the matrix.
  - 20. The method of claim 19, wherein the osteogenic precursor cells or peripheral blood cells are transfected *ex vivo*.
  - 21. A method of inducing systemic bone formation in a mammalian host in need thereof, comprising:
  - a) transfecting osteogenic precursor cells or peripheral blood leukocytes with a vector that is stablely maintained in the osteogenic precursor cells or peripheral blood leukocytes, the vector comprising SEQ ID NO: 39 operably linked to a regulatable promoter, wherein the regulatable promoter responds to an exogenous control compound; and
  - (b) administering to the host, as needed, an amount of the exogenous control substance effective to cause expression of SEQ ID NO: 39.

5

15

22. A method of stimulating production of an osteogenic soluble factor by an osteogenic cell, comprising:

- (a) transfecting the osteogenic cell or peripheral blood leukocyte with an isolated nucleic acid molecule comprising SEQ ID NO: 39; and
  - (b) overexpressing the isolated nucleic acid molecule.
  - 23. An osteogenic soluble factor produced by the method of claim 22.
- 24. The osteogenic soluble factor of claim 23, wherein the osteogenic factor is a protein.
- 25. A method of inhibiting the expression of HLMP-2 or HLMP-3
   10 comprising transfecting a cell wherein HLMP-2 or HLMP-3 is expressed with an antisense oligonucleotide.
  - 26. The method of claim of claim 17, wherein the isolated nucleic acid molecule is in a vector selected from the group consisting of a plasmid and a virus.
  - 27. The method of claim 26, wherein the vector is a plasmid, which plasmid is directly injected into muscle tissue.
    - 28. A method of inhibiting bone formation comprising transfecting osteogenic precursor cells or peripheral blood leukocytes with an isolated nucleic acid molecule comprising SEQ ID NO: 37.
    - 29. The method of claim 28, wherein the isolated nucleic acid molecule is in a vector.
      - 30. The method of claim 29, wherein the vector is an expression vector.
      - 31. The method of claim 30, wherein the vector is a plasmid.
      - 32. The method of claim 30, wherein the vector is a virus.
      - 33. The method of claim 32, wherein the virus is an adenovirus
      - 34. The method of claim 32, wherein the virus is a retrovirus.
    - 35. The method of claim 28, wherein the osteogenic precursor cells or peripheral blood leukocytes are transfected *ex vivo*.
- 36. The method of claim 28, wherein the osteogenic precursor cells are transfected *in vivo* by direct injection of the isolated nucleic acid molecule.

5

15

20

37. The method of claim 28, wherein the LIM mineralization protein is HLMP-2 (SEQ ID NO: 38).

#### SEQUENCE LISTING

- <110> Boden M.D., Scott D Hair Ph.D., Gregory A
- <120> LIM Mineralization Protein Splice Variants

- <130> splice variant LMP
- <140> 6148.0222-00304
- <141> 2000-04-28
- <150> 60/132,021
- <151> 1999-04-30
- <160> 42
- <170> PatentIn Ver. 2.1
- <210> 1
- <211> 457
- <212> PRT
- <213> Rattus norvegicus
- <400> 1
- Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe 1 5 10 15
- Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg
  20 25 30
- Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp 35 40 45
- Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
  50 55 60
- Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly 65 70 75 80
- Leu Ser Arg Ala Gln Pro Ala Gln Ser Lys Pro Gln Lys Ala Leu Thr 85 90 95
- Pro Pro Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Ala Ser Leu 100 105 110
- Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Thr Asp Ser Ala 115 120 125
- Leu Ser Gln Asn Gly Gln Leu Leu Arg Gln Leu Val Pro Asp Ala Ser 130 135 140
- Lys Gln Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly 145 150 155 160

Thr	Gly	Gln	Ser	Arg 165	Ser	Phe	Arg	Ile	Leu 170	Ala	His	Leu	Thr	Gly 175	Thr
Glu	Phe	Met	Gln 180	Asp	Pro	Asp	Glu	Glu 185	Phe	Met	Lys	Lys	Ser 190	Ser	Gln
Val	Pro	Arg 195	Thr	Glu	Ala	Pro	Ala 200	Pro	Ala	Ser	Thr	Ile 205	Pro	Gln	Glu
Ser	Trp 210	Pro	Gly	Pro	Thr	Thr 215	Pro	Ser	Pro	Thr	Ser 220	Arg	Pro	Pro	Trp
Ala 225	Val	Asp	Pro	Ala	Phe 230	Ala	Glu	Arg	Tyr	Ala 235	Pro	Asp	Lys	Thr	Ser 240
Thr	Val	Leu	Thr	Arg 245	His	Ser	Gln	Pro	Ala 250	Thr	Pro	Thr	Pro	Leu 255	Gln
Asn	Arg	Thr	Ser 260	Ile	Val	Gln	Ala	Ala 265	Ala	Gly	Gly	Gly	Thr 270	Gly	Gly
Gly	Ser	Asn 275	Asn	Gly	Lys	Thr	Pro 280	Val	Cys	His	Gln	Cys 285		Lys	Ile
Ile	Arg 290	Gly	Arg	Tyr	Leu	Val 295	Ala	Leu	Gly	His	Ala 300	Tyr	His	Pro	Glu
Glu 305	Phe	Val	Cys	Ser	Gln 310	Cys	Gly	Lys	Val	Leu 315	Glu	Glu	Gly	Gly	Phe 320
Phe	Glu	Glu	Lys	Gly 325	Ala	Ile	Phe	Cys	Pro 330	Ser	Cys	Tyr	Asp	Val 335	Arg
Tyr	Ala	Pro	Ser 340	Cys	Ala	Lys	Cys	Lys 345	Lys	Lys	Ile	Thr	Gly 350	Glu	Ile
Met	His	Ala 355	Leu	Lys	Met	Thr	Trp 360	His	Val	Pro	Cys	Phe 365	Thr	Cys	Ala
Ala	Cys 370	Lys	Thr	Pro	Ile	Arg 375	Asn	Arg	Ala	Phe	Tyr 380	Met	Glu	Glu	Gly
Ala 385	Pro	Tyr	Cys	Glu	Arg 390	Asp	Tyr	Glu	Lys	Met 395	Phe	Gly	Thr	Lys	Cys 400
Arg	Gly	Cys	Asp	Phe 405	Lys	Ile	Asp	Ala	Gly 410	Asp	Arg	Phe	Leu	Glu 415	Ala
Leu	Gly	Phe	Ser 420	Trp	His	Asp	Thr	Cys 425	Phe	Val	Cys	Ala	1le 430		Gln
Ile	Asn	Leu 435		Gly	Lys	Thr	Phe 440	Tyr	Ser	Lys	Lys	Asp 445		Pro	Leu
Cys	Lys 450		His	Ala	Phe	Ser 455	His	Val							

```
<210> 2
<211> 1696
<212> DNA
<213> Rattus norvegicus
<400> 2
gcacgaggat cccagcgcgg ctcctggagg ccgccaggca gccgcccagc cgggcattca 60
ggagcaggta ccatggattc cttcaaggta gtgctggagg gacctgcccc ttggggcttc 120
cgtctgcaag ggggcaagga cttcaacgtg cccctctcca tctctcggct cactcctgga 180
ggcaaggeeg cacaggeegg tgtggeegtg ggagaetggg tactgagtat egaeggtgag 240
aacgccggaa gcctcacaca cattgaagcc cagaacaaga tccgtgcctg tggggagcgc 300
ctcagcctgg gtcttagcag agcccagcct gctcagagca aaccacagaa ggccctgacc 360
cctcccgccg acccccgag gtacactttt gcaccaagcg cctccctcaa caagacggcc 420
eggeeetteg gggeaeeeee acetaetgae agegeeetgt egeagaatgg acagetgete 480
agacagetgg teeetgatge cageaageag eggetgatgg agaataetga agactggege 540
ccgcggccag ggacaggcca gtcccgttcc ttccgcatcc ttgctcacct cacgggcaca 600
gagttcatgc aagacccgga tgaggaattc atgaagaagt caagccaggt gcccaggaca 660
gaagecccag ecccageete aaceatacee caggaateet ggeetggeee caccacecee 720
agecceacea geogeocace etgggeogta gateetgeat ttgetgageg etatqeecea 780
gacaaaacca gcacagtgct gacccgacac agccagccag ccacacctac gcctctgcag 840
aaccgcacct ccatagttca ggctgcagct ggaggggca caggaggagg cagcaacaat 900
ggcaagacgc ctgtatgcca ccagtgccac aagatcatcc gcggccgata cctggtaqca 960
ctgggccacg cgtaccatcc tgaggaattt gtgtgcagcc agtgtgggaa ggtcctggaa 1020
gagggtggct tcttcgagga gaagggagct atcttttgcc cctcctgcta tgatgtgcgc 1080
tatgcaccca gctgtgccaa atgcaagaag aagatcactg gagagatcat gcatgcgctg 1140
aagatgacct ggcatgttcc ctgcttcacc tgtgcagcct gcaaaacccc tatccgcaac 1200
agggetttet acatggagga gggggetece tactgegage gagattaega gaagatgttt 1260
ggcacaaagt gtcgcggctg tgacttcaag atcgatgccg gggaccgttt cctggaagcc 1320
ctgggtttca gctggcatga tacgtgtttt gtttgcgcaa tatgtcaaat caacttggaa 1380
ggaaagacct tctactccaa gaaggacaag cccctgtgca agagccatgc cttttcccac 1440
gtatgagcac ctcctcacac tactgccacc ctactctgcc agaagggtga taaaatgaga 1500
gagetetete teectegace tttetgggtg gggetggeag ceattgteet ageettgget 1560
cctggccaga tcctggggct ccctcctcac agtccccttt cccacacttc ctccaccacc 1620
accaccgtca ctcacaggtg ctagcctcct agccccagtt cactctggtg tcacaataaa 1680
cctgtatgta gctgtg
                                                                  1696
<210> 3
<211> 260
<212> DNA
<213> Rattus norvegicus
<400> 3
ttctacatgg aggaggggc tccctactgc gagcgagatt acgagaagat gtttggcaca 60
aagtgtegeg getgtgaett caagategat geeggggaee gttteetgga ageeetgggt 120
ttcagctggc atgatacgtg ttttgtttgc gcaatatgtc aaatcaactt ggaaggaaag 180
accttctact ccaagaagga caagcccctg tgcaagagcc atgccttttc ccacgtatga 240
gcacctcctc acactactgc
                                                                  260
<210> 4
<211> 16
<212> DNA
```

<213> MMLV

```
<400> 4
aagctttttt tttttg
                                                                   16
<210> 5
<211> 13
<212> DNA
<213> MMLV
<400> 5
                                                                   13
aagcttggct atg
<210> 6
<211> 223
<212> DNA
<213> Homo sapiens
<400> 6
atcettgete aceteaeggg cacegagtte atgeaagace eggatgagga geacetgaag 60
aaatcaagcc aggtgcccag gacagaagcc ccagccccag cctcatctac accccaggag 120
ccctggcctg gccctaccgc ccccagccct accagccgcc cgccctgggc tgtggaccct 180
gcgtttgccg agcgctatgc cccagacaaa accagcacag tgc
                                                                  223
<210> 7
<211> 717
<212> DNA
<213> Homo sapiens
<400> 7
atggatteet teaaggtagt getggagggg ecageacett ggggetteeg getgeaaggg 60
ggcaaggact tcaatgtgcc cctctccatt tcccggctca ctcctggggg caaagcggcg 120
caggeeggag tggeegtggg tgaetgggtg etgageateg atggegagaa tgegggtage 180
ctcacacaca tegaagetea gaacaagate egggeetgeg gggagegeet eageetggge 240
etcagcaggg cccagccggt tcagagcaaa ccgcagaagg cctccgcccc cgccgcggac 300
cctccgcggt acacctttgc acccagcgtc tccctcaaca agacggcccg gccctttggg 360
gegeeeege eegetgacag egeeeegeaa eagaatggae ageegeteeg acegetggte 420
ccagatgcca gcaagcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 480
acaggecagt egegtteett eegeateett geecacetea eaggeacega gtteatgeaa 540
gacccggatg aggagcacct gaagaaatca agccaggtgc ccaggacaga agccccagcc 600
ecageeteat etacaeecca ggageeetgg eetggeeeta eegeeeccag eeetaeeage 660
cgcccgccct gggctgtgga ccctgcgttt gccgagcgct atgccccgga caaaacg
<210> 8
<211> 1488
<212> DNA
<213> Homo sapiens
<400> 8
atcgatggcg agaatgcggg tagcctcaca cacatcgaag ctcagaacaa gatccgggcc 60
tgcggggagc gcctcagcct gggcctcagc agggcccagc cggttcagag caaaccgcag 120
aaggesteeg eeccegeege ggaceeteeg eggtacacet ttgcacecag egteteeete 180
aacaagacgg cccggccctt tggggcgccc ccgcccgctg acagcgcccc gcaacagaat 240
```

ggacageege teegaceget ggteecagat geeageaage ageggetgat ggagaacaca 300

```
gaggactggc ggccgcggcc ggggacaggc cagtcgcgtt ccttccgcat ccttgcccac 360
ctcacaggca ccgagttcat gcaagacccg gatgaggagc acctgaagaa atcaagccag 420
cctaccgccc ccagccctac cagccgcccg ccctgagctg tggaccctgc gtttgccgag 540
cgctatgccc cggacaaaac gagcacagtg ctgacccggc acagccagcc ggccacgccc 600
acgccgctgc agagccgcac ctccattgtg caggcagctg ccggaggggt gccaggaggg 660
ggcagcaaca acggcaagac teeegtgtgt caccagtgee acaaggteat eeggggeege 720
tacctggtgg cgttgggcca cgcgtaccac ccggaggagt ttgtgtgtag ccagtgtggg 780
aaggtcctgg aagagggtgg cttctttgag gagaagggcg ccatcttctg cccaccatgc 840
tatgacgtgc gctatgcacc cagctgtgcc aagtgcaaga agaagattac aggcgagatc 900
atgcacgccc tgaagatgac ctggcacgtg cactgcttta cctgtgctgc ctgcaagacg 960
cccatccgga acagggcctt ctacatggag gagggcgtgc cctattgcga gcgagactat 1020
gagaagatgt ttggcacgaa atgccatggc tgtgacttca agatcgacgc tggggaccgc 1080
ttcctggagg ccctgggctt cagctggcat gacacctgct tcgtctgtgc gatatgtcag 1140
atcaacctgg aaggaaagac cttctactcc aagaaggaca ggcctctctg caagagccat 1200
gccttctctc atgtgtgagc cccttctgcc cacagctgcc gcggtggccc ctagcctgag 1260
gggcctggag tcgtggccct gcatttctgg gtagggctgg caatggttgc cttaaccctg 1320
getectggee egageetggg etecegggee eetgeecaee cacettatee teceaececa 1380
ctecetecae caccacagea caccggtget ggecacacca gececettte acetecagtg 1440
ccacaataaa cctgtaccca gctgaattcc aaaaaaatcca aaaaaaaa
```

<210> 9 <211> 1644 <212> DNA

<213> Homo sapiens

## <400> 9

atggatteet teaaggtagt getggagggg ceageacett ggggetteeg getgeaaggg 60 ggcaaggact tcaatgtgcc cctctccatt tcccggctca ctcctggggg caaagcggcg 120 caggccggag tggccgtggg tgactgggtg ctgagcatcg atggcgagaa tgcgggtagc 180 ctcacacaca tcgaagctca gaacaagatc cgggcctgcg gggagcgcct cagcctgggc 240 ctcagcaggg cccagccggt tcagagcaaa ccgcagaagg cctccgcccc cgccgcggac 300 cctccgcggt acacctttgc acccagcgtc tccctcaaca agacggcccg gccctttggg 360 gcgcccccgc ccgctgacag cgccccgcaa cagaatggac agccgctccg accgctggtc 420 ccagatgcca gcaagcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 480 acaggccagt cgcgttcctt ccgcatcctt gcccacctca caggcaccga gttcatgcaa 540 gacceggatg aggageacet gaagaaatea ageeaggtge eeaggacaga ageeecagee 600 ccagcctcat ctacacccca ggagccctgg cctggcccta ccgcccccag ccctaccagc 660 cgcccgccct gggctgtgga ccctgcgttt gccgagcgct atgccccgga caaaacgagc 720 acagtgetga eceggeacag ecageeggee acgeecacge egetgeagag ecgeacetee 780 attgtgcagg cagctgccgg aggggtgcca ggagggggca gcaacaacgg caagactccc 840 gtgtgtcacc agtgccacaa ggtcatccgg ggccgctacc tggtggcgtt gggccacgcg 900 taccacccgg aggagtttgt gtgtagccag tgtgggaagg tcctggaaga gggtggcttc 960 tttgaggaga agggcgccat cttctgccca ccatgctatg acgtgcgcta tgcacccagc 1020 tgtgccaagt gcaagaagaa gattacaggc gagatcatgc acgccctgaa gatgacctgg 1080 cacgtgcact getttacetg tgetgeetge aagacgeeca teeggaacag ggeettetae 1140 atggaggagg gcgtgcccta ttgcgagcga gactatgaga agatgtttgg cacgaaatgc 1200 catggctgtg acttcaagat cgacgctggg gaccgcttcc tggaggccct gggcttcagc 1260 tggcatgaca cotgettegt etgtgegata tgtcagatea acetggaagg aaagacette 1320 tactccaaga aggacaggcc tetetgcaag agecatgeet tetetcatgt gtgageeeet 1380 totgcccaca gotgccgcgg tggcccctag cotgaggggc ctggagtcgt ggccctgcat 1440 ttctgggtag ggctggcaat ggttgcctta accctggctc ctggcccgag cctgggctcc 1500 cgggcccctg cccacccacc ttatcctccc accccactcc ctccaccacc acagcacacc 1560 ggtgctggcc acaccagccc cctttcacct ccagtgccac aataaacctg tacccagctg 1620 aattccaaaa aatccaaaaa aaaa 1644

<210> 10

<211> 457

<212> PRT

<213> Homo sapiens

<400> 10

Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe
1 5 10 15

Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg 20 25 30

Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp.
35 40 45

Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
50 55 60

Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly 65 70 75 80

Leur-Serr-Arg-Ala Gln Pro Val Gln Ser Lys-Pro Gln-Lys-Ala-Ser Ala 85 90 95

Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Val Ser Leu 100 105 110

Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala 115 120 125

Pro Gln Gln Asn Gly Gln Pro Leu Arg Pro Leu Val Pro Asp Ala Ser 130 135 140

Lys Gln Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly 145 150 155 160

Thr Gly Gln Ser Arg Ser Phe Arg Ile Leu Ala His Leu Thr Gly Thr
165 170 175

Glu Phe Met Gln Asp Pro Asp Glu Glu His Leu Lys Lys Ser Ser Gln 180 185 190

Val Pro Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu

Pro Trp Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro Trp 210 215 220

Ala Val Asp Pro Ala Phe Ala Glu Arg Tyr Ala Pro Asp Lys Thr Ser 225 230 235 240

Thr Val Leu Thr Arg His Ser Gln Pro Ala Thr Pro Thr Pro Leu Gln 245 250 255

Ser Arg Thr Ser Ile Val Gln Ala Ala Gly Gly Val Pro Gly Gly

			260					265					270			
Gly	Ser	Asn 275	Asn	Gly	Lys	Thr	Pro 280	Val	Cys	His	Gln	Cys 285	His	Lys	Val	
Ile	Arg 290	Gly	Arg	Tyr	Leu	Val 295	Ala	Leu	Gly	His	Ala 300	Tyr	His	Pro	Glu	
Glu 305	Phe	Val	Cys	Ser	Gln 310	Cys	Gly	Lys	Val	Leu 315	Glu	Glu	Gly	Gly	Phe 320	
Phe	Glu	Glu	Lys	Gly 325	Ala	Ile	Phe	Cys	Pro 330	Pro	Cys	Tyr	Asp	Val 335	Arg	
Tyr	Ala	Pro	Ser 340	Cys	Ala	Lys	Cys	Lys 345	Lys	Lys	Ile	Thr	Gly 350	Glu	Ile	
Met	His	Ala 355	Leu	Lys	Met	Thr	Trp 360	His	Val	His	Cys	Phe 365	Thr	Cys	Ala	
Ala	Cys 370	Lys	Thr	Pro	Ile	Arg 375	Asn	Arg	Ala	Phe	Tyr 380	Met	Glu	Glu	Gly	
Val 385	Pro	Tyr	Cys	Glu	Arg 390	Asp	Tyr	Glu	Lys	Met 395	Phe	Gly	Thr	Lys	Cys 400	
His	Gly	Cys	Asp	Phe 405	Lys	Ile	Asp	Ala	Gly 410	Asp	Arg	Phe	Leu	Glu 415	Ala	
Leu	Gly	Phe	Ser 420	Trp	His	Asp	Thr	Cys 425	Phe	Val	Cys	Ala	Ile 430	Cys	Gln	
Ile	Asn	Leu 435	Glu	Gly	Lys	Thr	Phe 440	Tyr	Ser	Lys	Lys	Asp 445	Arg	Pro	Leu	
Cys	Lys 450	Ser	His	Ala	Phe	Ser 455	His	Val								
<211 <212	> 11 > 22 > DN > Ra		nor	vegi	cus								٠			
<400 gcca			ccca	gtca	c ga	i.										22
<210 <211 <212 <213	> 22 > DN		nor	vegi	cus											

22

<400> 12

gccagggttt tcccagtcac ga

WO 00/66178		PCT/US00/1166
<210> 13	•	
<211> 22		
<212> DNA		
<213> Homo sapiens		
<400> 13		22
tcttagcaga geceageetg et		22
<210> 14		
<211> 22		
<212> DNA		
<213> Homo sapiens		
<400> 14		22
gcatgaactc tgtgcccgtg ag		22
<210> 15		
<211> 20		
<212> DNA	·	
<213> Rattus norvegicus	and the second s	
<400> 15		
atccttgctc acctcacggg		20
<210> 16		,
<211> 22		
<212> DNA		
<213> Rattus norvegicus		
<400> 16		
gcactgtgct ggttttgtct gg		22
<210> 17		
<211> 23		
<212> DNA		
<213> Homo sapiens		
<400> 17		
catggattcc ttcaaggtag tgc		23
<210> 18		
<211> 20		
<212> DNA		
<213> Homo sapiens		
<400> 18		
gttttgtctg gggcagagcg		20
<210> 19 <211> 44		

```
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: adaptor for
      Marathon RACE reactions
<400> 19
ctaatacgac tcactatagg gctcgagcgg ccgcccgggc aggt
                                                                   44
<210> 20
<211> 27
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: PCR primer
      specific for Marathon RACE adaptor
<400> 20
ccatcctaat acgactcact atagggc
                                                                   27
<210> 21
<211> 765
<212> DNA
<213> Homo sapiens
<400> 21
ccgttgtttg taaaacgacg cagagcagcg ccctggccgg gccaagcagg agccggcatc 60
atggatteet teaaggtagt getggagggg ceageacett ggggetteeg getgeaaggg 120
ggcaaggact tcaatgtgcc ctcctccatt tcccggctca cctctggggg caaggccgtg 180
caggccggag tggccgtaag tgactgggtg ctgagcatcg atggcgagaa tgcgggtagc 240
ctcacacaca tcgaagctca gaacaagatc cgggcctgcg gggagcgcct cagcctgggc 300
ctcaacaggg cccagccggt tcagaacaaa ccgcaaaagg cctccgcccc cgccgcggac 360
ceteegeggt acacetttge accaagegte teceteaaca agaeggeeeg geeettgggg 420
gegeeceege eegetgacag egeecegeag cagaatggac ageegeteeg acegetggte 480
ccagatgcca gcaagcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 540
acaggccagt gccgttcctt tcgcatcctt gctcacctta caggcaccga gttcatgcaa 600
gacceggatg aggagcacet gaagaaatca agccaggtge ecaggacaga agccccagee 660
ccagoctcat ctacaccca ggagccctgg cctggcccta ccgccccag ccctaccagc 720
cgcccgccct gggctgtgga ccctgcgttt gccgagcgct atgcc
                                                                  765
<210> 22
<211> 1689
<212> DNA
<213> Homo sapiens
<400> 22
cgacgcagag cagcgccctg gccgggccaa gcaggagccg gcatcatgga ttccttcaag 60
gtagtgctgg aggggccagc accttggggc ttccggctgc aagggggcaa ggacttcaat 120
gtgcccctct ccatttcccg gctcactcct gggggcaaag cggcgcaggc cggagtggcc 180
gtgggtgact gggtgctgag catcgatggc gagaatgcgg qtaqcctcac acacatcgaa 240
```

geteagaaca agateeggge etgeggggag egeeteagee tggqeeteag eagqqeeeag 300

```
ceggtteaga geaaacegea gaaggeetee geeceegeeg eggaceetee geggtacaee 360
tttgcaccca gcgtctccct caacaagacg gcccggcct ttggggcgcc cccgcccgct 420
gacagegece egeaacagaa tggacageeg eteegacege tggteecaga tgecageaag 480
cageggetga tggagaacac agaggaetgg eggeegegge eggggaeagg eeagtegegt 540
teetteegea teettgeeca eeteaeagge acegagttea tgeaagaece ggatgaggag 600
cacctgaaga aatcaagcca ggtgcccagg acagaagccc cagccccagc ctcatctaca 660
ecceaggage cetggcetgg ceetacegee eccageceta ceageegeee geeetggget 720
gtggaccctg cgtttgccga gcgctatgcc ccggacaaaa cgagcacagt gctgacccgg 780
cacagecage eggecaegee caegeegetg cagageegea cetecattgt geaggeaget 840
gccggagggg tgccaggagg gggcagcaac aacggcaaga ctcccgtgtg tcaccagtqc 900
cacaaggtca teeggggeeg etacetggtg gegttgggee acgegtacea eeeggaggag 960
tttgtgtgta gccagtgtgg gaaggtcctg gaagagggtg gcttctttga ggagaagggc 1020
gccatcttct gcccaccatg ctatgacgtg cgctatgcac ccagctgtgc caagtgcaag 1080
aagaagatta caggcgagat catgcacgc ctgaagatga cctggcacgt gcactgcttt 1140
acctgtgctg cctgcaagac gcccatccgg aacagggcct tctacatgga ggagggcgtg 1200
ccctattgcg agcgagacta tgagaagatg tttggcacga aatgccatgg ctgtgacttc 1260
aagategaeg etggggaeeg etteetggag geeetggget teagetggea tgacacetge 1320
ttegtetgtg egatatgtea gateaacetg gaaggaaaga eettetaete caagaaqqae 1380
aggestetet geaagagsea tgestietet catgtgtgag ceesttetge ceasagetgs 1440
egeggtggec cetageetga ggggeetgga gtegtggeec tgcatttetg ggtagggetg 1500
gcaatggttg ccttaaccct ggctcctggc ccgagcctgg gctcccgggc ccctqcccac 1560
ccaccttatc ctcccaccc actccctcca ccaccacagc acaccggtgc tggccacacc 1620
agcccccttt cacctccagt gccacaataa acctgtaccc agctgaattc caaaaaatcc 1680
aaaaaaaa
                                                                   1689
<210> 23
<211> 22
<212> DNA
<213> Homo sapiens
<400> 23
gcactgtgct cgttttgtcc gg
                                                                  22
<210> 24
<211> 21
<212> DNA
<213> Homo sapiens
<400> 24
tccttgctca cctcacgggc a
                                                                  21
<210> 25
<211> 30
<212> DNA
<213> Homo sapiens
<400> 25
tcctcatccg ggtcttgcat gaactcggtg
                                                                  30
<210> 26
<211> 28
<212> DNA
```

WO 00/66178		PCT/US00/11664
<213> Homo sapiens		
<400> 26		
gececegece getgacageg cecegeaa		28
<210> 27		•
<211> 24		
<212> DNA		
<213> Homo sapiens		
<400> 27		•
tccttgctca cctcacgggc accg		24
<210> 28		
<211> 22		
<212> DNA		
<213> Homo sapiens		
<400> 28		
gtaatacgac tcactatagg gc	#1#.	 22
<210> 29		
<211> 23 <212> DNA		
<213> Rattus norvegicus		
<400> 29		
gcggctgatg gagaatactg aag		23
<210> 30		
<211> 23		
<212> DNA		
<213> Rattus norvegicus		
<400> 30		
atcttgtggc actggtggca tac		23
<210> 31		
<211> 22		
<212> DNA		
<213> Rattus norvegicus		
<400> 31		
tgtgtcgggt cagcactgtg ct		22
<210> 32	•	
<211> 1620		
<212> DNA		
<213> Homo sapiens		

```
<400> 32
atggatteet teaaggtagt getggagggg eeageacett ggggetteeg getgeaaggg 60
ggcaaggact teaatgtgee cetetecatt teeeggetea eteetggggg caaageggeg 120
caggccggag tggccgtggg tgactgggtg ctgagcatcg atggcgagaa tgcgggtagc 180
ctcacacaca tcgaagctca gaacaagatc cgggcctgcg gggagcgcct cagcctgggc 240
ctcagcaggg cccagccggt tcagagcaaa ccgcagaagg cctccgcccc cgccgcggac 300
ceteegeggt acacetttge acceagegte teceteaaca agaeggeeeg gecetttggg 360
gegeeeeege eegetgaeag egeeeegeaa eagaatggae ageegeteeg aeegetggte 420
ccagatgcca gcaagcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 480
acaggecagt egegtteett eegeateett geecaeetea eaggeacega gtteatgeaa 540
gacccggatg aggagcacct gaagaaatca agccaggtgc ccaggacaga agccccagcc 600
ccagcctcat ctacacccca ggagccctgg cctggcccta ccgcccccag ccctaccagc 660
cgcccgccct gagctgtgga ccctgcgttt gccgagcgct atgccccgga caaaacgagc 720
acagtgctga cccggcacag ccagccggcc acgcccacgc cgctgcagag ccgcacctcc 780
attgtgcagg cagctgccgg aggggtgcca ggagggggca gcaacaacgg caagactccc 840
gtgtgtcacc agtgccacaa ggtcatccgg ggccgctacc tggtggcgtt gggccacgcg 900
taccaccegg aggagtttgt gtgtagecag tgtgggaagg teetggaaga gggtggette 960
tttgaggaga agggcgccat cttctgccca ccatgctatg acgtgcgcta tgcacccagc 1020
tgtgccaagt gcaagaagaa gattacaggc gagatcatgc acgccctgaa gatgacctgg 1080
cacqtgcact gctttacctg tgctgcctgc aagacgccca tccggaacag ggccttctac 1140
atggaggagg gcgtgcccta ttgcgagcga gactatgaga agatgtttgg cacgaaatgc 1200
catggctgtg acttcaagat cgacgctggg gaccgcttcc tggaggccct gggcttcagc 1260
tggcatgaca cctgcttcgt ctgtgcgata tgtcagatca acctggaagg aaagaccttc 1320
tactccaaga aggacaggcc tctctgcaag agccatgcct tctctcatgt gtgagcccct 1380
tetgeccaca getgeegegg tggeecetag eetgagggge etggagtegt ggeeetgeat 1440
ttctgggtag ggctggcaat ggttgcctta accctggctc ctggcccgag cctgggctcc 1500
egggeeeetg eccaeceace ttatecteee accecaetee etecaecace acageacace 1560
ggtgctggcc acaccagccc cctttcacct ccagtgccac aataaacctg tacccagctg 1620
<210> 33
<211> 1665
<212> DNA
<213> Homo sapiens
<400> 33
cgacgcagag cagcgccctg gccgggccaa gcaggagccg gcatcatgga ttccttcaag 60
gtagtgctgg aggggccagc acettggggc ttccggctgc aagggggcaa ggacttcaat 120
gtgcccctct ccatttcccg gctcactcct gggggcaaag cggcgcaggc cggagtggcc 180
gtgggtgact gggtgctgag catcgatggc gagaatgcgg gtagcctcac acacatcgaa 240
gctcagaaca agatccgggc ctgcggggag cgcctcagcc tgggcctcag cagggcccag 300
ccggttcaga gcaaaccgca gaaggcctcc gccccgccg cggaccctcc gcggtacacc 360
tttgcaccca gcgtctccct caacaagacg gcccggccct ttggggcgcc cccgcccgct 420
gacagegeee egeaacagaa tggacageeg eteegacege tggteecaga tgeeageaag 480
cageggetga tggagaacae agaggaetgg eggeegegge eggggaeagg eeagtegegt 540
tectteegea teettgeeea eeteacagge acegagttea tgeaagacee ggatgaggag 600
cacctgaaga aatcaagcca ggtgcccagg acagaagccc cagccccagc ctcatctaca 660
ecceaggage cetggeetgg cectacegee eccageeeta ceageegeee geeetgaget 720
gtggaccctg cgtttgccga gcgctatgcc ccggacaaaa cgagcacagt gctgacccgg 780
cacagecage eggecacgee caegecgetg cagageegea cetecattgt geaggeaget 840
geeggagggg tgeeaggagg gggeageaac aaeggeaaga eteeegtgtg teaccagtge 900
cacaaggtca teeggggeeg etacetggtg gegttgggee aegegtaeca eeeggaggag 960
tttgtgtgta gccagtgtgg gaaggtcctg gaagagggtg gcttctttga ggagaagggc 1020
gccatcttct gcccaccatg ctatgacgtg cgctatgcac ccagctgtgc caagtgcaag 1080
aagaagatta caggcgagat catgcacgcc ctgaagatga cctggcacgt gcactgcttt 1140
```

acctgtgctg cctgcaagac gcccatccgg aacagggcct tctacatgga ggagggcgtg 1200

ccctattgcg agcgagacta tgagaagatg tttggcacga aatgccatgg ctgtgacttc 1260 aagatcgacg ctggggaccg cttcctggag gccctgggct tcagctggca tgacacctgc 1320 ttcgtctgtg cgatatgtca gatcaacctg gaaggaaaga ccttctactc caagaaggac 1380 aggcctctct gcaagagcca tgccttctct catgtgtgag ccccttctgc ccacagctgc 1440 cgcggtggcc cctagcctga ggggcctgga gtcgtggccc tgcattctg ggtagggctg 1500 gcaatggttg ccttaaccct ggctcctgc ccgagcctgg gctcccgggc ccctgccac 1560 ccaccttatc ctccacccc actcctca ccacacac acccggtgc tggccacacc 1620 agccccttt cacctccagt gccacaataa acctgtaccc agctg

<210> 34

<211> 223

<212> PRT

<213> Homo sapiens

<400> 34

Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe
1 5 10 15

Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg
20 25 30

Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp 35 40 45

Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
50 55 60

Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly
65 70 75 80

Leu Ser Arg Ala Gln Pro Val Gln Ser Lys Pro Gln Lys Ala Ser Ala 85 90 95

Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Val Ser Leu 100 105 110

Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala 115 120 125

Pro Gln Gln Asn Gly Gln Pro Leu Arg Pro Leu Val Pro Asp Ala Ser 130 135 140

Lys Gln Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly 145 150 155 160

Thr Gly Gln Ser Arg Ser Phe Arg Ile Leu Ala His Leu Thr Gly Thr 165 170 175

Glu Phe Met Gln Asp Pro Asp Glu Glu His Leu Lys Lys Ser Ser Gln
180 185 190

Val Pro Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu 195 200 205

Pro Trp Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro

20

210 215 220

<210> 35
<211> 20
<212> DNA
<213> Homo sapiens

<400> 35
gagccggcat catggattce

<210> 36

<212> DNA <213> Homo sapiens

<400> 36 gctgcctgca caatggaggt

gctgcctgca caatggaggt 20

<210> 37 <211> 1456 <212> DNA <213> Homo sapiens

<400> 37

<211> 20

cgacgcagag cagcgccttg gccgggccaa gcaggagccg gcatcatgga ttccttcaag 60 gtagtgctgg aggggccagc accttggggc ttccggctgc aagggggcaa ggacttcaat 120 gtgccctct ccatttcccg gctcactcct gggggcaaag cggcgcaggc cggagtggcc 180 gtgggtgact gggtgctgag catcgatggc gagaatgcgg gtagcctcac acacatcgaa 240 geteagaaca agateeggge etgeggggag egeeteagee tgggeeteag eagggeecag 300 ccggttcaga gcaaaccgca gaaggtgcag acccctgaca aacagccgct ccgaccgctg 360 gtcccagatg ccagcaagca gcggctgatg gagaacacag aggactggcg gccgcggccg 420 gggacaggcc agtcgcgttc cttccgcatc cttgcccacc tcacaggcac cgagttcatg 480 caagacccgg atgaggagca cetgaagaaa teaagccagg tgeccaggac agaagcccca 540 geoceageet catetacace ceaggageee tggeetggee etacegeece cagecetace 600 ageogeoege cetgggetgt ggaccetgeg tttgeegage getatgeece ggacaaaaeg 660 ageacagtgc tgacceggca cagecagecg gecacgecca egecgetgca gageegcace 720 tccattgtgc aggcagctgc cggaggggtg ccaggagggg gcagcaacaa cggcaagact 780 cccgtgtgtc accagtgcca caaggtcatc cggggccgct acctggtggc gttgggccac 840 gcgtaccacc cggaggagtt tgtgtgtagc cagtgtggga aggtcctgga agagggtggc 900 ttetttgagg agaagggege catettetge ceaecatget atgacgtgeg ctatgeaece 960 agetgtgeca agtgeaagaa gaagattaca ggegagatea tgeaegeeet gaagatgace 1020 tggcacgtgc actgctttac ctgtgctgcc tgcaagacgc ccatccggaa cagggccttc 1080 tacatggagg agggcgtgcc ctattgcgag cgagactatg agaagatgtt tggcacgaaa 1140 tgccatggct gtgacttcaa gatcgacgct ggggaccgct tcctggaggc cctgggcttc 1200 agetggcatq acacetgett egtetgtgeg atatgteaga teaacetgga aggaaagace 1260 ttctactcca agaaggacag gcctctctgc aagagccatg ccttctctca tgtgtgagcc 1320 cettetgeec acagetgeeg eggtggeece tageetgagg ggeetggagt egtggeeetg 1380 catttetggg tagggetgge aatggttgee ttaaccetgg etcetggeec gageetggge 1440 tcccgggccc tgccca

<210> 38 <211> 423

<212> PRT

<213> Homo sapiens

<400> 38

Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe

1 5 10 15

Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg 20 25 30

Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp 35 40 45

Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
50 55 60

Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly 65 70 75 80

Leu Ser Arg Ala Gln Pro Val Gln Asn Lys Pro Gln Lys Val Gln Thr 85 90 95

Pro Asp Lys Gln Pro Leu Arg Pro Leu Val Pro Asp Ala Ser Lys Gln
100 105 110

Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly Thr Gly
115 120 125

Gln Ser Arg Ser Phe Arg Ile Leu Ala His Leu Thr Gly Thr Glu Phe 130 135 140

Met Gln Asp Pro Asp Glu Glu His Leu Lys Lys Ser Ser Gln Val Pro 145 150 155 160

Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu Pro Trp 165 170 175

Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro Trp Ala Val

Asp Pro Ala Phe Ala Glu Arg Tyr Ala Pro Asp Lys Thr Ser Thr Val

Leu Thr Arg His Ser Gln Pro Ala Thr Pro Thr Pro Leu Gln Ser Arg 210 215 220

Thr Ser Ile Val Gln Ala Ala Ala Gly Gly Val Pro Gly Gly Gly Ser 225 230 235 240

Asn Asn Gly Lys Thr Pro Val Cys His Gln Cys His Gln Val Ile Arg 245 250 255

Ala Arg Tyr Leu Val Ala Leu Gly His Ala Tyr His Pro Glu Glu Phe 260 265 270

Val Cys Ser Gln Cys Gly Lys Val Leu Glu Glu Gly Gly Phe Phe Glu

275 280 285

Glu Lys Gly Ala Ile Phe Cys Pro Pro Cys Tyr Asp Val Arg Tyr Ala 290 295 300

Pro Ser Cys Ala Lys Cys Lys Lys Lys Ile Thr Gly Glu Ile Met His 305 310 315 320

Ala Leu Lys Met Thr Trp His Val Leu Cys Phe Thr Cys Ala Ala Cys 325 330 335

Lys Thr Pro Ile Arg Asn Arg Ala Phe Tyr Met Glu Glu Gly Val Pro 340 345 350

Tyr Cys Glu Arg Asp Tyr Glu Lys Met Phe Gly Thr Lys Cys Gln Trp 355 360 365

Cys Asp Phe Lys Ile Asp Ala Gly Asp Arg Phe Leu Glu Ala Leu Gly 370 375 380

Phe Ser Trp His Asp Thr Cys Phe Val Cys Ala Ile Cys Gln Ile Asn 385 390 395 400

Leu Glu Gly Lys Thr Phe Tyr Ser Lys Lys Asp Arg Pro Leu Cys Lys
405 410 415

Ser His Ala Phe Ser His Val 420

<210> 39

<211> 1575

<212> DNA

<213> Homo sapiens

<400> 39

cgacgcagag cagcgccctg gccgggccaa gcaggagccg gcatcatgga ttccttcaag 60 gtagtgctgg aggggccagc accttggggc ttccggctgc aagggggcaa ggacttcaat 120 gtgcccctct ccatttcccg gctcactcct gggggcaaag cggcgcaggc cggagtggcc 180 gtgggtgact gggtgctgag catcgatggc gagaatgcgg gtagcctcac acacatcgaa 240 geteagaaca agateeggge etgeggggag egeeteagee tgggeeteag eagggeecag 300 ccggttcaga gcaaaccgca gaaggcctcc gccccgccg cggaccctcc gcggtacacc 360 tttgcaccca gcgtctccct caacaagacg gcccggccct ttggggcgcc cccgcccgct 420 gacagegeee egeaacagaa tgggtgeaga eeeetgacaa acageegete egacegetgg 480 teccagatge cageaageag eggetgatgg agaacacaga ggaetggegg eegeggeegg 540 ggacaggcca gtcgcgttcc ttccgcatcc ttgcccacct cacaggcacc gagttcatgc 600 aagacccgga tgaggagcac ctgaagaaat caagccaggt gcccaggaca gaagccccag 660 ccccagcete atetacacce caggagecet ggcctggccc taccgccccc agccctacca 720 gccgcccgcc ctgggctgtg gaccctgcgt ttgccgagcg ctatgccccg gacaaaacga 780 gcacagtgct gacceggcac agccagcegg ccacgcecac gccgctgcag agccgcacct 840 ccattgtgca ggcagctgcc ggaggggtgc caggaggggg cagcaacaac ggcaagactc 900 ccgtgtgtca ccagtgccac aaggtcatcc ggggccgcta cctggtggcg ttgggccacg 960 cgtaccaccc ggaggagttt gtgtgtagcc agtgtgggaa ggtcctggaa gagggtggct 1020 tetttgagga gaagggegee atettetgee caccatgeta tgaegtgege tatgeaccea 1080 gctgtgccaa gtgcaagaag aagattacag gcgagatcat gcacgccctg aagatgacct 1140 ggcacgtgca ctgctttacc tgtgctgcct gcaagacgcc catccggaac agggccttct 1200

acatggagga gggcgtgccc tattgcgagc gagactatga gaagatgttt ggcacgaaat 1260 gccatggctg tgacttcaag atcgacgctg gggaccgctt cctggaggcc ctgggcttca 1320 gctggcatga cacctgcttc gtctgtgcga tatgtcagat caacctggaa ggaaagacct 1380 tctactccaa gaaggacagg cctctctgca agagccatgc cttctctcat gtgtgagccc 1440 cttctgcca cagctgcgc ggtggccct agcctgaggg gcctggagtc gtggccctgc 1500 atttctgggt agggctggca atggttgcct taaccctggc tcctggcccg agcctgggct 1560 cccgggccct gccca

<210> 40

<211> 153

<212> PRT

<213> Homo sapiens

<400> 40

Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe

Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg
20 25 30

Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp
35 40 45

Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile
50 . 55 60

Glu Ala Gln Asn Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly 65 70 75 80

Leu Ser Arg Ala Gln Pro Val Gln Ser Lys Pro Gln Lys Ala Ser Ala 85 90 95

Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Val Ser Leu 100 105 110

Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala 115 120 125

Pro Gln Gln Asn Gly Cys Arg Pro Leu Thr Asn Ser Arg Ser Asp Arg 130 135 140

Trp Ser Gln Met Pro Ala Ser Ser Gly
145

<210> 41

<211> 24740

<212> DNA

<213> Homo sapiens

<400> 41

nnnnntgta ttttatcata ttttaaaaat caaaaaacaa aaggcagttg aggttaggca 60 tggaggttcg tgcctgtaat cccagcactt tgggaagccg aagcacgtgg atcacctgag 120 gtcaggagtt cgagaccagc ctgcccaata tggtaaaacc ctgtctctac taaaaataca 180 aaaaaattagc caggcatggt ggtgggcacc tgtaatccca gctacttggg agactgaggc 240

aggagaatca	cttaaacccq	ggagggggg	tgggcgcggt	ggctcatgcc	totaatccca	300
			tgaggtcagg			
			tacaaaaaaa			
			tgaggcagga			
			ctgcactcca			
			ggaggcggaa			
			caagactcca			
			gtggcatttg			
gccagggtgg	tgctagtaga	ggagctgagt	gggaaagagc	acaggggaca	tgctgaaggc	780
ctgggtgtgg	ggatgaggca	gagattgggg	gcaccttgca	gggtcatagc	aggtggctgt	840
ggtgagatgg	aggaagacac	ctggggtact	gctctaggct	gtcagacata	cagaagctgg	900
cccagccaag	cccaggggct	gcaagggaca	tccttttgtg	tccccagtga	tctgcagctc	960
tcagacaccc	tcaagcacag	tgcctcttgc	ccagcccagc	actctcagtg	gggagccagg	1020
tgggagaaca	ggctcggaag	gggacctagg	cttatgcagc	gagccgggca	aagctggaac	1080
			ttgtggagtt			
			tttgacttga			
			agtctcccaa			
			aaggtaagtt			
			tttgggactg			
			gcagaggact			
			cccagacgcc			
			cactteetge			
			gctgacagcc			
			ctcctttgag			
			ctgtgcccct			
			ccagagctga			
			gacacccccg			
			atgcacctgg			
			gatctggcgt			
			gagcacctct			
			cctgagccgc			
agccccacaa	ccagtcccat	ctaccacaac	ggccaggact	tgagctggcc	cggcccggcc	2160
aacgacagta	ccctggaggc	ccagtaccgg	cggctgctgg	agctggatca	ggtggagggc	2220
acaggccgcc	ctgaccctca	ggcaggtttc	aaggccaagc	tggtgaccct	gctgagtcgt	2280
gagcggagga	agggcccagc	cccttgtgac	cggccctgaa	cgcccagcag	agtggtggcc	2340
agaggggaga	ggtgctcccc	ctgggacagg	agggtgggct	ggtgggcaaa	cattgggccc	2400
			ctgcaggaac			
			atagtgtgca			
			cacacgcccc			
			cacaggtgac			
			gggatggaaa			
			gcggagggag			
			tggagacccg			
			tttcagagac			
			tacttctggg			
			tatcaggcat			
			gacacageeg			
			agagaagttt			
			atcctaggag			
			ctcaccgcaa			
			gacggacaca			
			tccagctcac			
ccgcccctgc	ctaggcatgt	cgacctgcag	ggaccctttt	ctggctcttc	gaggcctctg	3420
			ccttccccct			
			gagccagacc			
ctgcccacag	ccagagaaat	ccatttaagc	agactgcctg	catccttctt	aacagtgcaa	3600

```
ggcaggcact ccctgccaca agagaccctg ttccctagta gggcagcttt tctcctcccc 3660
agaaceteet gtetateeec acceaatgte teeteacagg catattgggg aaacaggtea 3720
ggetetecea cegtatetge aagtgtactg geatecatet gtettettee taccectaca 3780
gtagaaacag tgtctgtccc cagctgtgct ctgatcccgg ctcctttcac ctcagagctt 3840
ggaaaattga gctgtcccca ctctctcctg cgcccattca tcctaccagc agcttttcca 3900
gecacaegea aacatgetet gtaattteae attttaaaee tteeettgae eteacattee 3960
tetteggeea cetetgttte tetgtteete tteacageaa aaactgttea aaagagttgt 4020
tgattacttt catttccact ttctcaccc cattctctc tcaattaact ctccttcatc 4080
cccatgatgc cattatgtgg cttttattag agtcaccaac cttattctcc aaaacaaaag 4140
caacaaggac tttgacttct cagcagcact cagctctggt tcttgaaaca cccccgttac 4200
ttgctattcc tcctacctca taacaatctc cttcccagcc tctactgctg ccttctctga 4260
gttcttccca gggtcctagg ctcagatgta gtgtagctca accetgctac acaaagaatc 4320
tcctgaaagc ctgtaaaaat gtccatgcat gttctgtgag tgatctacca agaaaataaa 4380
aaattttaaa aatcaaatgc ccatgcctgg gcccacacgc aggggctctg atttcatcag 4440
tetggtaggt gggttetggg catecacget caetggattt ceggatgatt gtagtatgca 4500
gectaggetg ggaaccaetg geeteageaa gecagteatt etecaggtgt caeagaeeet 4560
ctaggtgcta atgaccccga aggtctgtct tcagtgcaca cctccccctg agctccagat 4620
ttaggaatce cactgcacac gagacatetg gatgtggaaa agacatetee agateceatg 4680
ggtgaaaggg ggttggggga atggagactc gtgttcttcc aggatgtgtg tggacacaga 4740
atgcaaagcc tggagggatg ctagagccat agggaggaag atttcggctc acttattcat 4800
gcaagcactt cctgatgggt aaggtcttag agcaagctga ggccaagagg cgggcagtcg 4860
aggtgctgct gcaggcaccc ccactcccta cagtggcaag cccaagccca gcccttggca 4920
gctcaaatcc caggacacgc tgaaggtcac ccagagagtc aggggcatgg ctagaaccag 4980
ttgtttccta gggatttctg tgatattcca aggggactgt gggaaagaaa gtccttggaa 5100
accaccagga cgctagaggc ctggcctgga gcctcaggag tctcggccac cagagggcgc 5160
tgggtccttg tccaggtcca gttgctacgc aggggctgcc tgtgctggga ggctccccag 5220
gggacacaga ccagagcett gcaccagece aaggaatggg ageetggggt eetetetget 5280
ggaggactgc caggaccccc aggctgccgc ctcttccttt gctcatttgc tgtttcactt 5340
tgtcaatcct teetttette gtgtgtteat teacateeae tgtgtgetgg eeetggggaa 5400
atgttagata agacacatta gctgtgtgtc ttcattgtcc taacaaagaa cacaccctgg 5460
aaagagcacc gcagagagtc cccattcccc catctccctc cacacatgga atctggagat 5520
gccttttcca catccagatg tctctggtgc tgtgggattc ttaaataaac aaacatttca 5580
tacagaatgt gagatgatgg agatgctatg gggaaaagta aagcagaggg agggcctagt 5640
gtgtgatgcg ggtgaggcat ccagggattg ctgtttcagc tgtgatcagg aaaggccctg 5700
ggaggaggcc acatctgagc agagacctaa ataaagttgg aaacctgttg ctgagatatc 5760
tggagaagtg tttcaagggc cgggcaccgg gcatggtggc tcacgcctgt aatcccagca 5820
ctttgggagg ccaaggcagg tggatcgctg gaggtcagga gtttgagagc agcctgacca 5880
acatggagaa accccatctc tactaaacat ataaaaatta tccgggcatg gtggttcatg 5940
cctgtagtcc cagctactcg ggaggttgag gcaggagaat cacttgaacg tgggaggcag 6000
aggttgcage aageegagat cacaceactg cactecagee tggatgacag agegagaete 6060
cgtctcaaaa aaaaaaaaga aaagaaaaaa gaaaaaaaa gaaaagtgtt tcaagcaggg 6120
gaactggcaa gtggagaggc cctgaggcag aaatatgctt ggcctgctgg aggaaatgtg 6180
agtgaggagg tcagggtggc tggagtggag ggagcgagtg gtaggagtca gacccagttt 6240
attcatattc tgtaggtctt aaggacttca gttttatttt gagtgcaata tgagcccact 6300
ggaatgctaa aagctgagag tgacatggtg ctgtgattct ggctttaaaa atatcacttt 6360
ggctgcttcg tgaagactct ggaaggggca agggtgaaag cagggatgcc cgttaggaga 6420
ccgttacagg ggcgcaggca caaaatggca gtggctggga caatggtggc agcagcggtt 6480
agatgtgaac atgttgaagg tggaatttgc agaatctggg ggaggacaga agagaaagga 6540
taacttcatc gtttctgctg aaccagttgg ataaatgttg gtggcacttc ttgaagtgag 6600
gaaggagtta ggaaggtggg aaaggcacaa gtttgaattg ggccatgatg gtctgagata 6660
cctagtacag tggttcccca acctttttgg cagaagggac cgctttcatg gaagacaatt 6720
tttccacaga ctgggggtgg ggtggggatg gtttcagggt ggttcgagtg cagtacattt 6780
atcattagac tettttttt ttttttttt tgagatagag tetegetetg teacceaeae 6840
tggagtgcag tggagccatc ttggctcact acaacctctg ctgcccaggt tcaagtcatt 6900
ctcctgcctc agcctctcaa gtagctggga ttataggcat atgcgccacc acgcccagct 6960
```

aatttttgta	tttttagtag	agacggggtt	tcaccatatt	ggccaggatg	gtctcgaact	7020
					tacaggcgtg	
					acctagatcc	
					ctgccgctgc	
					tcctaacagg	
					ttttggataa	
					gcctttgcat	
					ataaagagag	
					ccagtggggc	
					aacctggacc	
					tggccccatg	
					cggacaagag	
					cgataaacag	
					gctggagagg	
					tgtgcagaac	
					gtcctaggag	
					ctcaccccga	
					acgcggggcc	
gggactctga	ggacgcctcc	caggcgcgca	gtccgtctgg	ccaaggtgga	gcgggacggc	8100
ngcttccgac	ggtgcgcggg	teggeteggg	gttgcaggga	cateeggegt	ccgctcctgc	8160
cctgttttcc	tgccttcgca	gagcgttgcg	caactctagc	tttaaacgcc	cctgtccccc	8220
tcaacttgtc	tcccccagcc	cctctgattt	acagattctg	cagtccccga	gggttgcgcc	8280
tacgataccg	acactcgcgg	cagcctgcga=	ggcgagtatg	atcgtcccat	ttttcggagt	8340
agcaaactaa	ggttcagaga	ctactatgtc	ccaggtcggt	ctggtttgaa	ggtccgcttt	8400
cctctccctc	cgccagcggg	cggtgcgagg	gactgggcga	ggcagcgctt	ccctaaggag	8460
gcgacccgca	gccccggccc	cctcccgact	ccgccccgtt	gcagggcccg	ggtcggcgag	8520
					agctgggtgg	
					tggcctggtg	
					tcagaacact	
					tggccgggcc	
					gcccggggcc	
					tcccgctggc	
					tgcggcgggg	
					gcccgggcgg	
					gaggcaccac	
					ccagccccca	
					cccaggggtc	
					ccgctgcagc	
					ccctcccctc	
					gtgtcagctt	
					cctgggggcg	
					gctggcttgg	
					cctcaagcct	
					ctgtctttcc	
					cctcctatgg	
					tggtatcctg	
					tctcttctct	
					cagcaccttg	
					cccgggtgag	
					ccagagggtg	
					gtctaggtcc	
					ctgggaggga	
					ctcctgtggc	
					tttcaggagg	
					tgtgactata	
ggatttcact	tcagectcae	tgccgcccag	ggaatagcaa	tcctcatccc	gtttttccag	10320

atgagagaag aactcatgga gaggtggcgg ggctcgctca tcgagtccat ggtgaagcag 10380 ggattggaat tgaggcacag catggcgtac attttttgtg ggtagaaggg gtctctcccc 10440 agcctatgta aggacccaca tecaetgtte ceatteagga tgtggtggee tttgacccca 10500 agcagaagtg taggacaggg ctccattcta ggggcttaac ttcagcttcc aagagcctgc 10560 cctggtgtgg gtggagctgg aggctggctc ctccctgtag cagggggatt gccttataag 10620 cccaagaatg cagccccacg ctgggatggc caacagtggc tgcggtctgc agagctgaaa 10680 agggctggcc taggcctggc cccctgaacc ccactggtgg gcctctcagc tggtcaccag 10740 gctgcagctc cagctgtatg gtccagttgt gagacacaac aaattgcctg cccagagtgg 10800 gtgaggccag cctgtcggct ggcatctctg actggcctgg gggtcaggag ggggtgggga 10860 ettectgece etatateege etgeceegag agacecacee aggegeeggg tgggeaggea 10920 gctgttgtca ggaagcccaa ggcaagccca gcctggaggg gcccagaggg tcgtggcctg 10980 aggaggggct caagctggag tctgtctgta ggagctgggc gtgggggtta gggtgggcag 11040 gccagcagtg ctcttctcag gggtcctttg atggcattct cctggaacct gccccgccag 11100 cagggtagtg aggcagtggt tgccctatga cacacgtccc actacatagc cctcacacag 11160 ccctgaaacc tacctgacgt cctgctccct gggaaagtgc tggcccagtg tgtctgggga 11220 gcctgaacct cagtttcttc cctgatggag atgactttca gatatggcct gttgggggca 11280 ctccgggctc cagctccctg gtcagcatcc ctggcatgtg ggcggggcca ctagctgatc 11340 ccagccctgg agttggacct gggcccacat gggtgggtga ggtgggcttt tctgagttag 11400 gccagccccc tcccctccc ctgaccccag aatggaggga ggtgggaggg gcaagggctg 11460 gctgtgggcc caggcctggg agatgaggta acgtctggga ctggggggct gggctgctca 11520 ggctgactca ccccacctc atgcagggtc cagcccctg gctttttccc tccttggttc 11580 ctctggcctt accctgcccc tggcttgagc ccctccctgc ctctctccag ccacccgccc 11640 agegetgtet tetgetetee tgetgeeete eecaegetet gaacaceet cateetetgt 11700 getteetgee etecteacte tgggaaggga ageegteece geeeceace ecetetecag 11760 gagecageta getgeacece aagaeeeeca eetegggete ageecacage teecaggage 11820 cagccctgtg ggcagggagt ggctgggcca ggtttccctt ctactgactc accatgacct 11880 tgagtaagtc acttcccctc tggggtgtca cttccccata cacagtataa ggggttgatt 11940 tagttggatt gaactaaagg tgagggagtg gctcagggtg tctccaggtg ggctgacccc 12000 teagttgggc ccccatgctc agcagaggtg gcccacagtg gtggagcctt agggtcagag 12060 acacttectg getetgeete ttactagetg ggtgaettga ggeaagttgt ttaacetete 12120 tgtgtacatt tgcaagtgca aaatgggtaa aatcccagat tactccacaa ggttgttgga 12180 agattcagtg tcaatatgta gcatagttgg tgctcaataa actgaagcaa gtcttcttat 12240 ttagcgagtg aggaaggggc cgccgagctc tcttagcctt ctgacctcct acgcaagcaa 12300 gaggtcatgt tgagcccagc tegectttet ttteccagtg etgtcaaget etgtgcetgg 12360 etgecetgee etetgacate tetetgaaac etettgeete eceteteeet geeteagete 12420 agtetgtgca etgacecace tgaggageet eetggggeea etggeageet ggaceceee 12480 agatecece cacceagtga aattgtette cageactgee teacaaaage etacttgatg 12540 cagtgccaģg cetettgcca gatggetggg tggtecetta ggettggace cagtcaaget 12600 gecetgeetg tgttgetggg getgggetag aggeetggaa ggggtttate agggteaece 12660 teteagggee tgggagatae ecaateeeag acattaaaae tgeeagtage ecetetaeet 12720 tcaaagccaa gtcctggtcc cttcccctgg cattcaaagc catcgtaagt gaactctcac 12780 ccgctaggca gcacacgcca ttctccttta ccgaggccca ccgcttcctc aaagtcattc 12840 ctgatggtct cagctcatgc tggtggcagc catttctccc agcctactgt ctctactcat 12900 tgccacagga accagggact cccagctcaa gagcctgaag gattggggtc aggggaaatt 12960 ggcagtcgag ggcttgggag tgacagccat gtatggccta cgaagtccca gctgtcaact 13020 taggteeeat teaggeagtg tteacaggga acegggagat aacagggeet gtteetgget 13080 ctcaaagggt cccagcagac ccctatagat ggcccccgac agggtgctgg ggggtgagag 13140 gtccataaga gcccccggtg gtttcgggga ggaagctgcc ccctgcatgg gccagagggc 13200 atatctggta ggtggagtgg cctgggcagg aggccagcag gagcctcaaa aggcaatggt 13260 cctcctgaaa cacttgggct ttagcctgag cgtggctgtt tgtggacatc atagcaattt 13320 ctggactgtg ggggagggtg gtggcggtga atagataagc atcgtgactg gggaagctca 13380 ggtgagcacc acctgaggga gagggtctgg cagtgaataa ataagcagtg tgactgggaa 13440 attgtgaagc tcaggtgagc gccaccacct cctgggttgc tttagtgtcc agcagctgcc 13500 tagaactatg ttgaatgaag agctctctgg gttctggaag tgggacagct ttgggtgggg 13560 cagtgttacc accgtcagcc tggcttgggt ctgcagggtc cagggcctcg gtcactttgc 13620 ttetetetee acageteaet eetgggggea aageggegea ggeeggagtg geegtgggtg 13680

actgggtgct gagcatcgat ggcgagaatg cgggtagcct cacacacatc gaagctcaga 13740 acaagateeg ggeetgeggg gagegeetea geetgggeet cageaggtat gegggtggae 13800 atggatgggt gegecegege tggcagtggg gatecetgeg geceggeeeg etgtcaeget 13860 tteettetee teeagggeee ageeggttea gageaaaceg cagaaggtae gaggetggee 13920 gggacatccg ggcggtgggc ggtgtgggct tggacggcca ggcctgctcg ccctcctggc 13980 acattetegg taccecaate cetggeeggg agtggaggge agaaacegga getaaggegg 14040 gtctagggcc ctggagttga gccaggggct gctgcacggt cctggcacca cgcatgtccg 14100 cetgtetgte egeetgtetg teegeetget geeteeegee geeggegetg egtgetegee 14160 cgcactcggt cagccctcgg tcctgcgtgg actgagatcg ccactcccaa atgggcccct 14220 tgaaacetga gtegteetet eeeegtagee teeaaataga tgtagggggt ggggtggggg 14280 tggggggetg gagetgeege tgteetetge tgeaggegee ceaetteeae eeaggeeeee 14340 acettacect geoegeooge cetgeoogge tgtgtetetg cecaggeete egeoecegee 14400 geggaeeete egeggtaeae etttgeaeee agegteteee teaacaagae ggeeeggeet 14460 ttgggcgccc ccgcccgctg acagcgcccc gcagcagaat gggtacgtcg gcccctgccc 14520 geoegegeee aegecateag geoeactgtg geoecaegee egetgeeege tgetgeteag 14580 tetgtgetge gececagece ggeggaaceg tgeggeacge cecetggegg ecggggtggg 14640 getgeaggea cagggeecet ecegaggetg tggegeettg cagggeaceg cetggggagg 14700 ggtctctgaa tgacgccgcg cccctgctg gcggctgggg gttgggttgt ggtgtcgggc 14760 cagctgagcc ccagacactc agtgccgcct tgtccccggc tgttctgacc cctccccgtc 14820 tttetteete teetgtgtet gteeetttgt eeetttatet gtetgtetgt ettattteet 14880 tcacaggtgc agacccctga caagtcagtg agcccccctc tgcctgtgcc tttcttcttc 14940 ettttggeac tetgggtgge ggeceteec caecetgget geceteetet ceaettegee 15000 ctectgteet eteacetace egeccageag ggeteetgge eteaceetta eccaetecet 15060 cccatcactg taacccaaac ccacatgcac caaatcctgg gaggggctgc ccccaccgcc 15120 caccccagt gtggggttct gagccacacc ctccccacag acagccgctc cgaccgctgg 15180 teccagatge cageaageag eggetgatgg agaacacaga ggaetggegg eegeggeegg 15240 ggacaggcca gtcgcgttcc ttccgcatcc ttgcccacct cacaggcacc gagttcagta 15300 agtgccagcc cagggcaggg ggtactttcc tcgcccccag cccaggcgtg atccctgacc 15360 ctgtgtcttt tttggtcaat gcctgcctct gctctctcag tgcaagaccc ggatgaggag 15420 cacctgaaga aatcaaggta cagggacggg caccagccc tctcccacct cctgcctctt 15480 ccattccage tactgccctg tgtctactcc tgaggctccc agctggggct ctcaattctc 15540 cetteettee tteetteett eetteettee tteetteett eetteettee tteetteett 15600 coeffected theoretect effication theoretect coeffected effected 15660 cetgeeteee ttecatetet cetteettee acttetteet ceetetete etgeecetea 15720 gggaaaagta tgtcctggag ctgcagagcc cacgctacac ccgcctccgg gactggcacc 15780 accagegete tgcccaegtg etcaaegtge agtegtagee eggecetete eageeggetg 15840 coctotetge etecetettt etgtteetee tgeecaggge accecettag tgeetecage 15900 ttetgeetae eteaececee etttegtgee eetggeetga geeteetget ggeetggeee 15960 tggccgccca cctgggttca tctgacactg ccttccctct ttgccctgtg gtactgctgt 16020 ctgccaggtc tgtgctgcct tgggcatgga ataaacattc tcagccctgc ttgctctgcc 16080 tgtcttctat ctttgtggac ctggtttgca tttggggtgt gggggtgttt cgtggttcgg 16140 actgtttggg ccctgccgtc cttgttttca gtgggagggg gtacctggca aaggggccct 16200 gecetgecat cacagatgge tteetggeat gaggggagee ecaggagetg ceteagaage 16260 gggagccctg cctcgtctcc cagctagaga ccgcacacca gctaactgga cattgctagg 16320 agaagetgee etteceatee etaececagt gggacetgga atecaacteg geagttteea 16380 cgccccagt cateteccgt ggggccagca ggacccaggt tgggggtgg ggccatgtca 16440 ggaagctcag ccatgcaggg ccttgaatgg cagatcttgc agccaggtgc ccaggacaga 16500 agccccagcc ccagcctcat ctacacccca ggagccctgg cctggtgaga gggagtgggc 16560 tegggeetgg geaagggtgg geageeteea ggggeatggg ggtggtggge tteteteage 16620 tgcctggggc tccacccccg tcctttgggg tccctgggca cccctttaga gtcactttcc 16680 ceggcaggcc ctacegeccc cagecetacc agecgecege cetgggetgt ggaccetgeg 16740 tttgccgagc gctatgcccc ggacaaaacg agcacagtgc tgacccggca cagccagccg 16800 gccacgccca cgccgctgca gagccgcacc tccattgtgc aggcagctgc cggaggggtg 16860 ccaggagggg gcagcaacaa cggcaagact cccgtgtgtc accagtgcca caaggtcatc 16920 eggtgggtgg cetgtteetg teegaceetg gettteecat cetgeageec ageceeacet 16980 gtctgcccac ctgtcttgcc tcagctgcga ctggggggaa taaggattca gttctcagct 17040

ggagtaggag	tagggacctg	ggctgggtcc	tcccattctt	aatcccacgc	tacctacccc	17100
agcccaccca	caacaactgc	tagcagcatc	tgccgtggcg	aaatagccga	agggccaacc	17160
ataggctgaa	gctgcacccc	tacctttgct	getetetggg	caaagagggg	cctgccccct	17220
cccagcgcgt	ctgccctcc	ctcctgctct	ctgtctccct	ctgctctcag	agcatacagg	17280
cctggagcca	ctccctctgt	gcactgccc	gtggggccaa	gcagcatcaa	acacccccca	17340
gcatcagcgt	gccggattct	agageettee	taattcgcag	gcctggcctg	ctctcatctc	17400
tgtcagctct	tttttttt	tttttgaaac	agagteteae	tgtgttgccc	acgttggcgt	17460
geagtggege	gatctcggct	cactgcaacc	tetgeeteet	gggttcaaga	gattctcctg	17520
cctcagcctc	ctgagtagct	gggattacag	gcacccgcca	ccatgcctgg	ctaattttgt	17580
	gagacggggt					
aggtgatete	aggcctgcct	tggcctccca	aagtgctggg	actacaggtg	tgagccactg	17700
tgcccagccg	actctatcag	ctcttgccag	gtagaacagg	caggccagca	ggacagggca	17760
geteeagggt	ttgcccaggg	geggeteage	ttttatgagg	ctccagtcgt	cagcccttcc	17820
teeeggggte	ctccctgctc	taaagetgee	teteetgtea	ccagcagttc	agtgtggcgg	17880
	taagcttcat					
	gggagaatga					
ctgaaggggg	ccaagacccc	cacctctctg	cgtgtgctcc	ctctgaccgc	tctcgcctcc	18060
ctgcaggggc	cgctacctgg	tggcgctggg	ccacgcgtac	cacccggagg	agtttgtgtg	18120
tagecagtgt	gggaaggtcc	tggaagaggg	tggcttcttt	gaggagaagg	gcgccatctt	18180
ctgcccacca	tgctatgacg	tgcgctatgc	acccagctgt	gccaagtgca	agaagaagat	18240
	agtagggctg					
	gaaacccaca					
gcgtaataaa	tgggctgctg	tactggcctc	-accctgcatt	agtcaggatg	ctcttaacaa	18420
atgaccatgt	tcctgctcag	aaaccgccca	aggctgcaaa	gagcaggagg	accaagccag	18480
gagaagccct	gggccctcct	gactcccact	ttgggctctc	cctgccctgg	tgaaatgaca	18540
	cttgacacgc					
	ggctgccagg					
aggtttacct	tgaccacgag	cagtgggaaa	ctcaaaataa	tggtggctta	tttctcatct	18720
aaaaacatcc	cggggtgggt	ggtctgggac	tgatctggtg	gacccagget	ccgccttgtt	18780
gcttgactgt	tggcagcacc	tgcttactta	ccactcatgg	tgcaagatga	cacttcagcc	18840
tccgccaaaa	tgctcacctt	ccagccagca	ggaagtcgga	aggagaagaa	aggggacaga	18900
	gtccatcctt					
	agtcacatga					
	cccatgtgtc					
	gcagtggctg					
	gtcaggagtt					
	aaaattagct					
ctgaggcagg	agaatcgctt	gatcttgaga	ggtggaggtt	gcagtgagcc	aagattgtgc	19320
cactgccttc	cagcctggga	gacagcaaaa	aaaaaaaaa	aaaaaaaaa	aaaaagggcc	19380
aggcacggtg	gctcacacct	gtaatcccag	cactttggga	ggccgagatg	ggcggatcac	19440
	gattgagacc					
	tggccgggca					
	aacctgggag					
	acagagcgag					
	gggctcctgc					
	ccaagttcct					
agacagtgag	aagagccagc	cagacatggg	cagaaaacct	atttctgtga	tctactggct	19860
gtgtgagcag	gggctagttg	ctctctctgg	gcctcactga	agagaagggt	ggcactatgc	19920
	acggttgcaa					
aatgcaccat	ccgaaggcta	catgtcccc	acacttatgt	cttgcttggc	ccacactgtt	20040
tcattttaaa	atcagtagca	aacaatttaa	aaaatcagaa	gatttgcctg	catgatgcag	20100
rggctcatgc	ctgtaatccc	agcactttgg	gaggccaagg	tgggaggatt	gcttgagccc	20160
aggagttcaa	gaccagcatg	ggcaccatag	caagacccct	gtttctacaa	aaaaaaaaa	20220
accagaaaat	tagccaagtg	tggtggcatg	cacctgtggt	cccagctact	tgggaggcag	20280
agggaaagtg	agatctcctg	cttttattt	ctttatgtat	aatgataggg	tettgetetg	20340
ttgcccaggc	tggagtgcag	tggcatgatc	actgctcact	gcagccttga	tctcctgggc	20400

```
tcagaggatc ctcccacctc agcctcccaa atagctagga ctagaggtgc ccaccagcat 20460
gctcagcaga tttttaaatc tttttgtaga gatgaggttt tgctatgttg cccaggctgg 20520
tetegaaete etggeetega gegateetee cacettggee teecaaagea etgggattae 20580
agacgtgage cactgegece ageagattte tetttaacae ctagatttea geetgageea 20640
ggcaggcatt cctgaatgaa ccagtagtac tgctcccaga agaagaggtc ctcctccgtg 20700
tgacacagte eccaettgge cettgeaggg attggatetg ggatecetgg atttaaacte 20760
agggecatec teataacage etcacaagge tgggattage tteecagtte acaagggaag 20820
aaaccaagac ttgagaaggt caaggtetgg ceagaceeac acatettgga ceeteatace 20880
gcctcgaggc cccatgctgc cctctgcctg ctccagatgt gaatactgct ggccctggct 20940
ggccccggct ggccccgagg gtcctaggga tgaacagccc agcccaggga gagctcagcc 21000
cettgtgcct etgeceette ceaecteetg eggaggecag tegaeteace cacaaaggge 21060
caqqcactgt ggggatagat cagctaacaa aacagttgat gcttcctgcc cttctgggcc 21120
ttacattttg gctggaagaa gaggggagag gcagactgta agcaataagc gcaataagta 21180
ggttgcctgg aagtaatgtt agatcacgtt acggaaaaca ggaaagagca gagcgacaag 21240
tgctggggtg cgtggtgcag ggaaggcagc tggctgctgc tggtgtggtc agagtgggcc 21300
ctcatggaga agactgcatt cgagcagaaa cttgaagggg gtgaggggtg agcctagaga 21360
tatctggggc agagcagtcc aggcagaggg gacagccggt gtcaagccca ggacaggagt 21420
gtgcctggtg tgccagtttc aggcaagagg ccagtgtgca gaggcaaggt gagaacgcaa 21480
gggagagcag tggcggagac gggtgggaac gaggtcagac ctgctggcct ccagcctctg 21540
catggggctt ggctcttgct gggagcaatg ggaagcagta cacagtttca tgcaggggga 21600
gaaggeetgt ettgggttge aggggeacge tgtggeaget gggateagag agaggagett 21660
gtaggccagt tgttatgtgg tcccacgggc cagatggcca tggcttacct cacttcaggg 21720
aggetgtgag aageaeteag aatetggatg tgeettgggg gtgggeeeca etggatttee 21780
tggtggacct ggtgtggggt gtgagaggag ggtgtgtttg gctgcagcag acaggagaat 21840
ggagttgcca tccgcgtgat ggggatggct gtgggaggag aggtttgggg tgagggaatc 21900
aggaactgag tgctggacat ggcaagtctg aaggcgcagt ggtcgtccac tcagagacct 21960
tggagttgga gatggaggtg tgggagteet gaacagttag atgtagtgtt taccgcgaga 22020
aggaacaggg cttgcggcca gccctcctgt gttcccgtga cccagggcag ggcaggaggg 22080
gcctgagcct gccgagtgac tgggacctcc ttccaggaga tcatgcacgc cctgaagatg 22140
acctggcacg tgcactgctt tacctgtgct gcctgcaaga cgcccatccg gaacagggcc 22200
ttctacatgg aggagggcgt gccctattgc gagcgaggta cccactggcc agtgagggtg 22260
aggagggatg gtgcatgggg caggcatgaa tccaggtcct ctttctctct gcccccattc 22320
tcagactatg agaagatgtt tggcacgaaa tgccatggct gtgacttcaa gatcgacgct 22380
ggggaccgct teetggagge cetgggette agetggeatg acacetgett egtetgtgeg 22440
gtgagagece egeceetega aetgagecee aageceaeeg gecetetgtt cattececag 22500
gagatgeagg agaagttggg aaggggeete teetgetgee cecaacecea tgtgactggg 22560
cctttgctgt ccttagatat gtcagatcaa cctggaagga aagaccttct actccaagaa 22620
qqacaqqcct ctctgcaaqa gccatgcctt ctctcatgtg tgagcccctt ctgcccacag 22680
ctgccgcggt ggcccctagc ctgaggggcc tggagtcgtg gccctgcatt tctgggtagg 22740
gctggcaatg gttgccttaa ccctggctcc tggcccgagc ctggggctcc ctgggccctg 22800
ccccaccac cttatectec caccccacte cetecaccac cacageacac egatgetgge 22860
cacaccagec ccettteace tecagtgeca caataaacet gtacccaget gtgtettgtg 22920
tgcccttccc ctgtgcatcc ggaggggcag aatttgaggc acgtggcagg gtggagagta 22980
agatggtttt cttgggctgg ccatctgggt ggtcctcgtg atgcagacat ggcgggctca 23040
tggttagtgg aggaggtaca ggcgagaccc catgtgccag gcccggtgcc cacagacatg 23100
aggggagcca ctggtctggc ctggcttgga ggttagagaa gggtagttag gaagggtagt 23160
tagcatggtg gctcatgcct gtgatcccag cactttggaa ggccaaggtg ggcagatcgc 23220
ttgaggtcag gagttcgaga cctcatggcc aacacggtga aacagcgtct ctagtaaaaa 23280
tacaaaaatt ageegagtgt ggtggggcat geetgtaate ceageeacte aggaggetga 23340
ggcgggaaaa tcacttgaac ctgggaagtg gaggttgcag tgagctgaga tcacaccact 23400
gegegegage etgggtggca gatggeagag egagaeeetg etteaaaaaa aaaaaaaaa 23460
aaaaaaaaa gaagggtagt tgtagttggg ggtggatctg cagagatatg gtgtggaaaa 23520
cagcaatggc cacagcaaag teetggaggg gecagetgee gtecaaacag aagaaggcag 23580
ggctggagag ggtagccctt aggtcctggg aagccacgag tgccaggcag tagagctggg 23640
gctgtctctt gaggttaggg cagggcaagg cacagcagag tttgaaatag gtttgtgttg 23700
tattgcagaa aagaggcccc agaacactga gggagtgcag gagggaggct gggaggagga 23760
```

```
gttgcagcag ggcctagggg cgggggccag gcaagggagg ggcagagagt aatatggcag 23820
agatgggacc cagtggcagg tccgggggat gagggatgga gagaaggaca ggagcgttgc 23880
caggcatctg gcctatacca gacatgctca cgctgtctcc cgcgaacctc ctagcaacct 23940
ttttaagaga caggatetee etaggttgee egggetggtt teaaaeteet gggeteaage 24060
aattetteet eettageece aaagtgetgg cattacaggt gtgagecace atgeetggee 24120
cacttatttt ctagatgagg cacagaaaga ttgggagact tgaccaaggt cacgctgtca 24180
ttgagccatg agccagacta gaatccaggc ctgaagctgg gtgcgctgtc ccaggactgg 24240
ctggcactga gtaccatttg ccagcgagca tetetetggg aagetgaett ctgcccggta 24300
cctggaggac tgtagacctt ggtggtggcg ccgtcactct ggggcttcct gcctcccact 24360
gatgcccgca ccaccctaga gggactgtca teteteetgt eccaageetg gaetggaaag 24420
actgaagaga agcettaagt aggecaggac agetcagtgt gecatggetg ceegteette 24480
agtggtccct ggcatgagga cctgcaacac atctgttagt cttctcaaca ggcccttggc 24540
ceggteceet ttaagagaeg agaagggetg ggeaeggtga eteacacete taateecage 24600
actttggaag getgaggetg gagaaggget ceagettagg agtteaggae eageetggge 24660
aacatggtga gaccctgttt tgttttgttt tttgtttttt tgagatggag tcttgctctg 24720
tegeceagge tggagtgeag
                                                              24740
```

<210> 42

<211> 25

<212> DNA

<213> Rattus norvegicus

<400> 42

gcactacctt gaaggaatcc atggt

25

## INTERNATIONAL SEARCH REPORT

International application No. PCT/US00/11664

A. CLASSIFICATION OF SUBJECT MATTER						
IPC(7) :Please See Extra Sheet. US CL :435/320.1, 325; 530/350, 388.1; 536/23.5; 514/44						
According to International Patent Classification (IPC) or to bot						
B. FIELDS SEARCHED						
Minimum documentation searched (classification system followed by classification symbols)						
U.S.: 435/320.1, 325; 530/350, 388.1; 536/23.5; 514/44						
Documentation searched other than minimum documentation to the	ne extent that such documents are included in the fields searched					
Electronic data base consulted during the international search (	name of data base and, where practicable, search terms used)					
Please See Extra Sheet.	name of data base and, where practicable, scarcif terms used)					
THE SECOND DAILS DIRECT						
C. DOCUMENTS CONSIDERED TO BE RELEVANT						
Category* Citation of document, with indication, where a	appropriate, of the relevant passages Relevant to claim No.					
A BODEN et al. LMP-1, a LIM-dom	ain protein, mediates BMP-6 1-37					
effects on bone formation. Endocrine	- 1					
12, pages 5125-5134, entire documen	t.					
A HOGAN et al. Bone morphogenet	ic proteins: multifunctional 1-37					
regulators of vertebrate development						
1996. Vol. 10, pages 1580-1594, ent	- I					
A KINGSLEY, DM What do BMPs do	· · · · · · · · · · · · · · · · · · ·					
No. 1, pages 16-21, entire document.	* '1					
100. 1, pages 10-21, entire document.	·					
	i					
Further documents are listed in the continuation of Box	C. See patent family annex.					
Special categories of cited documents:	"T" later document published after the international filing date or priority					
"A" document defining the general state of the art which is not considered to be of particular relevance	date and not in conflict with the application but cited to understand the principle or theory underlying the invention					
*E* earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step					
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other	when the document is taken alone					
special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is considered with one or more other such documents such combination.					
means	document referring to an oral disclosure, use, exhibition or other combined with one or more other such documents, such combination being obvious to a person skilled in the art					
*P* document published prior to the international filing date but later than the priority date claimed	*& * document member of the same patent family					
Date of the actual completion of the international search	Date of mailing of the international search report					
16 JULY 2000	29 AUG 2800					
Name and mailing address of the ISA/US	Authorized officer 1 11 / 184					
Commissioner of Patents and Trademarks Box PCT	ANNE MARIE BAKER					
Washington, D.C. 20231 Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196					

Form PCT/ISA/210 (second sheet) (July 1998)\*

## INTERNATIONAL SEARCH REPORT

International application No. PCT/US00/11664

A. CLASSIFICATION OF SUBJECT MATTER: IPC (7):	
A61K 48/00; C07H 21/04; C07K 14/00, 16/00; C12N 15/63, 15/85, 15/86	
B. FIELDS SEARCHED Electronic data bases consulted (Name of data base and where practicable terms used):	
WEST Dialog (file: medicine) search terms: gene(w)transfer, LMP, bone, LIM, mineral?, human, osteogen?	
t de servicio de la servicio del servicio de la servicio della ser	-

Form PCT/ISA/210 (extra sheet) (July 1998)\*

## BROVENCE LISTING

- <120> LIM Mineralization Protein Splice Variants
- <130> splice variant LMP
- <140> 6148.0222-00304
- <141> 2000-04-79
- <150> 60/132,021
- c151> 1999-04-30
- <160> 42
- <170> Patentin Ver. 2.1
- <210× 1
- <211> 457
- <212> PRT
- <213> Rattus norvegious

----

- <400> 1
- Net Asp Ser Pho Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe 1 5 10 15
- Arg Lau Gin Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg 20 25 30
- Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp 35 40 45
- Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile 50 55 50
- Glu Ala Glu Aen Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly 65 70 75 60
- Leu Ser Arg Ala Glt Pro Ala Gln Ser Lys Pro Gln Lys Ala Leu Thr 85 90 95
- Pro Pro Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Ala Ser Leu 100 105 110
- Aen Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Thr Asp Ser Als 115 120 125
- Leu Ser Gln Asn Gly Gln Leu Leu Arg Gln Leu Val Pro Asp Ala Ser 130 135 140
- Lye Gln Arg Leu Met Glu Aen Thr Glu Asp Trp Arg Pro Arg Pro Gly 145 150 155 160

Thr Gly Gln Ser Arg Ser Phe Arg Ila Lau Ala His Leu Thr Gly Thr

- Glu Phe Not Gin Amp Pro Amp Glu Glu Phe Met Lym Lym Ser Ser Glu 180 135 190
- Val Pro Arg Thr Glu Ala Pro Ala Pro Ala Ser Thr Ile Pro Gla Glu 195 200 205
- Ser Trp Pro Gly Pro Thr Thr Pro Ser Pro Thr Ser Arg Pro Pro Trp 210 215 220
- Ala Val Acp Pro Ala Phe Ala Glu Arg Tyr Ala Pro Acp Lys Thr Ger 225 230 235 240
- Thr Val Leu Thr Arg Ris Ser Gln Pro Ala Thr Pro Thr Pro Leu Gln 245 250 255
- ABD Arg Thr Ser Ile Val Glm Ala Ala Gly Gly Gly Thr Gly Gly 265 270
- Gly Ser Asn Asn Gly Lys Thr Pro Val Cys His Gln Cys His Lys Ile 275 280 285
- Ile Arg Gly Arg Tyr Leu Val Ala Leu Gly His Ala Tyr His Pro Glu
  290 295 300
- Glu Phe Val Cys Ser Gln Cys Gly Lys Val Leu Glu Glu Gly Gly Phe 305 310 315 320
- Phe Glu Glu Lys Gly Ala fle Phe Cys Pro Ser Cys Tyr Asp Val Arg 325 330 335
- Tyr Ala Pro Ser Cys Ala Lys Cys Lys Lys Lys Ile Thr Gly Glu Ile 340 345 350
- Met His Ala Leu Lys Met Thr Trp His Val Pro Cys Phs Thr Cys Ala 355 360 365
- Ala Cys Lys Thr Pro Ile Arg Asn Arg Ala Phe Tyr Met Glu Glu Gly 370 375 380
- Als Pro Tyr Cys Glu Arg Asp Tyr Glu Lys Met Phe Gly Thr Lys Cys 385 390 395 400
- Arg Gly Cys Asp Phe Lys Ile Asp Ala Gly Asp Arg Phe Leu Glu Ala 405 410 415
- Leu Gly Phe Ser Try Ris Rep Thr Cys Phe Val Cys Ala Ile Cys Gln
  420 425 430
- Ile Agn Leu Glu Gly Lys Thr Phe Tyr Ser Lys Lys Asp Lys Pro Leu 435 440 445
- Cya Lya Cer His Ala Phe Ser His Val 450 455

```
<210> 3
<211> 1696
<212> DNA
<213> Rattus norvegicus
<400> 2
gcacgaggat cocagogogg ctcctggagg cogccaggea gccgcceage egggcattca 60
ggagcaggta coatggatte etteaaggta gtgetggagg gacetgeece ttgggggette 120
ogtotgcaag ggggcaagga ottesaegtg cocctotoca tototogget cactootgga 180
ggcamggcog cacaggcogg tgtggcogtg ggagactggg tactgagtat cgacggtgag 240
aacgooggaa gootcacaca cattgaagoo cagaacaaga toogtgootg tggggagogo 300
ctcagectgg gtcttageag ageccageet gctcagagea aaccaeagaa ggccetgace 360
cotocogecy accococogay gracactiti gcaccaagoy cotocotosa caagacygoc 430
eggecetteg gggeacecec acetaetgae agegeetgt egeagaatgg acagetgete 480
agacagetgg tecetgatge cageaageag eggetgatgg agastaetga agactggege 540
coccepccag ggacaggeca gtocogttoc tteegcatec ttgeteacet caegggeaca 600
gagttestge aagaeeegga tgaggaatte atgaagaagt caageeaggt geeeaggaea 660
gasgoccag coccagooto aaccataced cagysatect ggootggood caccacecco 720
agecceacca googoccace etgggccgta gatectgcat ttgctgagcg ctatgcccca 780
gacaasacca gcacagtgot gaccogacac agecagecag ccacacetae gcototgoag 840
aaccycacct ccatagttca ggctycagct ggaggyggca caggaggagg cagcaacaat 900
ggcaagacge ctgtatgcca coagtgccae aagatcatce geggeegata cetggtagea 960
ctgggccacg cgtaccatec tgaggaattt gtgtgcagcc agtgtgggaa ggtcctggaa 1030
gagggtggct tettegagga gaagggaget Atettttgee ceteetgeta tgatgtgege 1080
targeaccea gerotgecas argenegaag aagatemetg gagagarent geargegerg 1140
amgatgacet ggcatgttee etgettemee tgtgemgeet gemammeee tateegemme 1200
agggetttet acatggagga gggggetece tactgcgage gagattacga gaagatgttt 1260
ggcacaaagt gtogoggotg tgacttcaag atogatgoog gggaccgttt cotggaagco 1320
ctgggtttca gctggcatga tacgtgtttt gtttgcgcaa tatgtcaaat caacttggaa 1380
ggaaagacet tetaeteeaa gaaggacsag coccetytyca agagecatyc cttttcccae 1440
gtatgagcae etectoacae taetgedaec etactetgec agaagggtga taaaatgaga 1500
gagetetete teeetegaee titetgggtg gggetggeas ceatigteet ageetigget 1550
cetggeraga teetgggget ecetretear agreecettt eccaerette etreaccare 1620
accacegica etcacaggig ciageotopi ageoccagti cacieiggig icacaataaa 1680
                                                                  1696
cetgtatgta getgtg
<210> 3
<211> 260
<212> DNA
<213> Rattus norvegicus
<400> 3
ttotacatgg aggaggggggggge tocotactgo gagogagatt acgagaagat gtttggcaca 60
aagtstegeg getgtgaett caagategat geegggsace gttteetgga ageettggt 120
ttcagetggc stgatscgtg ttttgtttgc gcaatatgtc sastcaactt ggaaggaaag 180
acettetaet coaagaagga caageeeetg tgeaagagee atgeetttte eeaegtatga 240
                                                                  360
gcacctcotc acactactgc
<2105 4
<211> 16
<212> DNA
<213> WNLV
```

<400> 4 aagettttt	tttttg					16
<210> 5 <211> 13 <212> DNA <213> MMLV						
<400> 5 aagetEgget	a <b>t</b> g					13
<210> 6 <211> 223 <212> DWA <213> Homo	sapiens					
c400> 6 atcettget© aaatcaagee ceetggeetg gegtttgeeg	aggtg¢¢¢ag gccctaccg¢	gacagaagee ceeeageeet	accsdccdcc ccsdccccaa	ogccetggge	accccaggag	120
<210> 7 <211> 717 <212> DNA <213> Homo	sapiens					
ggcaaggact caggccggag ctcacacaca ctcagcaggg cctccgcggt gcgcccccgc ccagatgcca acaggccagt	tggecgtggg tcgaagetea eccagetggt acacetttge ecgutgacag gcaageageg egegttectt aggageacet etacacecca	cototocate tgactgggtg gmacaagale tcagagcaaa mcccagcgte cgccccgcaa gotgatgggg ccgcatcott gaagaaatca ggagccctgg	terregetra etgageateg egggertgeg eegeagmagg teceteaaea eagaatggae aacacagagg gecementea agceaggtge eetggeeeta	ccaccccag ccaggacaga actggcgcca agacggccca agacggccca agacggccca agacggccca agacggccca agacggccca agacggccca agacggcaca actggggga	caaageggeg tgegggtage eggettggge egeethtggg accgetggte geggeeggg gtteatgeaa ageweeagee ceetaceage	120 180 240 300 360 420 480 540 600
<210> 8 <211> 1488 <212> DWA <213> Homo	gapiens			•		
Egeggggage aaggeeteeg aacaagaegg	agaatgeggg geeteageet eeceggeege ceeggeeett teegaeeget	gggcctcagc ggaccctccg lggggcgccc	agggcccage cggcccgctg	cggttcagag ttgcacccag acagcgcccc	caaaccgcag cgtctccctc gcaacagaat	120 180 240

PCT/US00/11664 WO 00/66178

```
gaggactggc ggccgcgcc ggggacaggc cagtcgcgtt ccttccgcat ccttgcccac 360
etcacagges cogagitest gessgacceg gaigaggage accigaagaa atcsagcesg 420
gtgcccagga cagaaggcccaggcc tcatctacac cccaggager ctggcctggc 480
ectacogoco coagocotae cageogeecy ecetgagetg tggacoctge gthtgccgag 540
cgctatgccc cggacsaast gagcacagtg ctgacccggc acagccagcc ggccacgccc 600
acgoogotgo agagoogoac otroattgig caggoagotg ooggagggt gocaggaggg 660
ggeageaaca acggesagae teoegtgtgt caccagtgee acaaggtest coggggeege 720
tacctggtgg ogttgggcca cgcgtaccac ccggaggagt ttgtgtgtag ccagtgtggg 780
asysteetgg Asgagggtgg ettetttgag gagaagggcg coatcitetg coesceatge 840
tatgacytgo gotatgcaco cagotytgoc amytgcmagm myaagattac aggcyagato ago
atgraugere tyaagatgae etograagto ractgettta cetotgetee etocaagaag 960
cocatoogga acagggoott otacatggag gagggegtgd ectattgdga gegagactat 1020
gagaagatgt ttggcacgas atoccatggc tgtgacttea agatcgacgc tggggaccoc 1000
ttootggagg cootgggott cagotggcat gacacetget tegtetgtgc gatatgtcag 1140
atcascrigg aaggamagec cttctactoc magaaggaca ggootototg caagagccat 1200
goottototo abgigigaço coctioigos cacagoigos geggiggese ciagootgag 1260
gggcctggag togtggccct gcatttetgg gtagggctgg coatgettgc cttaaccctg 1320
gotoctggcc cgagodtegg otocogggcc cotgcocace cacettatee teccacecca 1380
etecetecae caccacagea caccggtget ggecaeaeea gooccettte acctecagtg 1440
ccacaatana cotgtacora gotgaattoo aaaaaaateca aaaaaaaa
<2105 S
-211> 1644
<233> Homo sapiene
```

<212> DNA

<400> 9

atggatteet teaaggtagt getggaggg ecageacett ggggetteeg getgeaaggg 60 ggcamggact tcaatgtgcc ceteteratt beerggebea etectggggg caaagcggcg 120 caggeoggag tggccdtggg tgactgggtg ctgagcatog atggcgagsa tgcgggtage 180 ctcacacaca togaagetca gascasgato ogggoottgog gygagogoot cagootgggo 240 otoagoaggg cocageeggt teagageaaa eegeagaagg eeteegeeee egeegeggae 300 cetoegoggt acacettige acceagegie teceteaaea agaeggeeeg gecettiggg 360 gegdddddd ddgdtgacag ogdocogoan cagaatggac agcogotoog accgetgghd 420 ccagatgoca gcaagcagog gotgatggag aacacagagg actggcggcc gcggccgggg 480 acaggerage cgrgetecett cogeatectt geocacetea caggeacega geteatgeaa 540 gacceggatg aggageacet gaagaaatea ageeaggtge eeaggacaga ageeecagee 600 enagecteat etacacecca ggagecetgg cetggeceta cogececcag cectaceage 660 egocegooot gagotataga occtaeattt geegageget atgeeeegaa caaaaogage 720 acagtgetga eceggeseay eengemygoo mogoocnogo egotgoagag cogeacetee 780 attgtgcagg cagetgoogg aggggtgcca ggagggggca gcaacsacgg caagactooc 840 gtgtgtcacc agtgccacaa ggtmatmogg ggccgctacc tggtggcgtt gggccacgcg 900 taccaccegg aggagtttgt gtgtagecag tgtgggaagg teetggaaga gggtggette 960 titgaggaga agggegecat ettetgeeca esatgotatg acgtgegeta tgcacceage 1020 tytyccaayt ycaayaayaa gattacayyo gayatcatyo acyccctyaa yatyocctyy 1000 cargigract gettiacrig ignigrater asgangeres tooggaacag geoticiae 1140 atggaggagg gegtgeecta ttgegagega gactatgaga agatgtttgg caegaaatge 1200 catggctgtg acttcaagat egaegetggg gacegettee tggaggeest gggetteage 1260 tggcatgaca cotgettogt otgtgogata tgtcagatca acctggaagg Aasgacette 1320 tactocaaga aggacaggon kobntgoaag agddatgoot tototoatgt gtgagoocot 1380 tetgeccaea getgeegegg tggeccetag cetgagggge etggagtegt gggdetgeat 1440 ttotgggtag ggotggcast ggttgcotta aconhggoto otggcoogag ootgggotoc 1500 egggeeets cocacecate that esteet acceptate et ceateare aragracate 1560

1644

39tyctggcc acaccagece cetticaect reagtgerse nathoacctg tacccagetg 1620

aatteeaaaa aateeaaaaa aasa

<210> 10

c211> 457

<212> PRT

<213> Homo Bapi në

<400> 10

Met Asp Ser Phe Lya Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe 1 5 10 15

Arg Lau Gin Gly Gly Lys Asp Phe Asn Val Pro Lau Ser Ile Ser Arg 20 25 30

Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp 25 40 45

Trp Val Leu Ser Ile Asp Gly Glu Aan Ala Gly Ser Leu Thr His Ile
50 55 60

Glu Ale Gln Aen Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly
65 70 75 80

Leu Ser Arg Ala Gln Pro Val Gln Ser Lys Pro Gln Lys Ala Ser Ala 85 90 95

Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Val Ser Leu 100 105 110

Asn Lye Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala 115 120 125

Pro Glm Glm Asm Gly Glm Pro Leu Arg Pro Leu Val Pro Asp Ala Ser 130 135 140

Lye Gln Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly 145 150 155 160

Thr Gly Gln Ser Arg Ser Phe Arg Ile Leu Ala Hie Leu Thr Gly Thr 165 170 175

Glu Phe Met Gln Asp Pro Asp Glu Glu His Leu Lys Lys Ser Ser Gln 190 185 190

Val Pro Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu 195 200 205

Pro Trp Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro Trp 210 215 220

Ala Val Asp Pro Ala Phe Ala Glu Arg Tyr Ala Pro Asp Lys Thr Ser 225 230 235 240

Thr Val Leu Thr Arg His Ser Gln Pro Ala Thr Pro Thr Pro Leu Gln
245 250 255

Ser Arg Thr Ser Ile Val Gin Ale Ale Ale Gly Gly Val Pro Gly Gly

270

260 265

Gly Ser Asn Asn Gly Lys Thr Pro Val Cys His Gln Cys His Lys Val 275 280 285

Ile Arg Gly Arg Tyr Leu Val Ala Leu Gly His Ala Tyr His Pro Glu 290 295 300

Glu Phe Val Cys Ser Gln Cys Gly Lys Val Leu Glu Glu Gly Gly Phe

Phe Glu Glu Lys Gly Ala Ile Phe Cys Pro Pro Cys Tyr Asp Val Arg 325 330 335

Tyr Ala Pro Ser Cys Ala Lys Cys Lys Lys Lys Ile Thr Gly Glu Ile 340 345 350

Met His Ala Leu Lys Met Thr Trp His Val Ris Cys Phe Thr Cys Ala 355 360 365

Ala Cys Lye Thr Pro Ile Arg Asn Arg Ala Phe Tyr Met Glu Glu Gly 370 375 380

Val Pro Tyr Cys Glu Arg Asp Tyr Glu Lys Met Phe Gly Thr Lys Cys 385 390 395 400

His Gly Cys Asp Phe Lys Ile Asp Ala Gly Asp Arg Phe Leu Glu Ala 405 410 415

Leu Gly Phe Ser Trp His Asp Thr Cys Phe Val Cys Ala Ile Cys Gln 420 425 430

Ile Asn Leu Glu Gly Lys Thr Phe Tyr Ser Lys Lys Asp Arg Pro Leu 435 440 445

Cys Lys Ser His Ala Phe Scr His Val 450 455

<210> 11

<211> 22

<212> DNA

<213> Rattue norvegicus

<400> 11

gecagggttt teccagteac ga

22

<210> 12

<211> 22

<212: DNA

<213> Rattus norvegique

<400> 22

gccagggttt tcccagtcac ga

22

WO 00/66178	PCT/US00/11664
<210> 13 <211> 22 <212> INA	
<213> Homo sapiens	
<400> 13 tettageaga geceageetg et	22
<210> 14 <211> 22 <212> DNA	
<213> Homo sapiens	
<400> 14 gcatgaactc tgtgcccgtg ag	22
<210> 15 <211> 20 <212> DWA	
<213> Rattua norvegicus	+
<400> 15 atcottgete sectemeggg	20
<210> 16 <211> 22 <212> DNA	
<213> Rattus norvegicus	
<400> 16 geactgtget ggttttgtet gg	22
<210> 17 <211> 23 <212> DNA	
<213> Homo sapiens	
<400> 17 catggattee tteaaggtag tge	23
<210> 18 <211> 20 <212> DNA	
<213> Homo sapiens	
<400> 18 gttttgtctg gggcagagcg	20
<210> 19	

<211> 44

```
<212> DNA
<213> Artificial Baquence
<220>
<223> Description of Artificial Sequence: adaptor for
             Marathon RACE reactions
24B05 19
ctaatacgac tcactatagg gctcgagcgg ccgcccgggc aggt
                                                                                                                                                  40
<210> 20
<211> 27
<212> DNA
<213> Artificial Sequence
<220>
<2235 Description of Artificial Sequence: DCR primer
             specific for Marathon RACE adaptor
<400> 20
ccatoctaat acgaeteect ataggge
                                                                                                                                                  27
<210> 21
<211> 765
<212> DMA
<2135 Homo eapiene
c400> 21
costtatta taaaacaca caaacaaca coctageess gecaageass ageesseate 60
atggatteet teaaggtagt getggagggg ceageacett ggggetteeg getgeaaggg 120
ggoaaggact toaatgtgcc ctcctccatt tcccggctca cctctggggg caaggccgtg 180
raggeoggag tggeogtaag tgaetgggtg etgageateg atggegagaa tgcgggtAge 240
ctoacacaca tegaagetca gaacaagate egggeetgeg gggagegeet cageetggge 300
ctcaacaggg cocageoggt toagaacaaa cegeaaaaagg ceteegeeee egeegeggac 360
cottoggeget acaectities accaagosto tocotoaaca agaoggecog geootteggeg 420
gagbacaege abgetgadag ogedecydag dagaatggad agecgeteeg adegetggtd 480
ccagatgcca gcasgcagcg gctgatggag aacacagagg actggcggcc gcggccgggg 540
acaggeragt groutfeett tegratoott geteacetta daggeacega gtteatgeas 600
quocoggatg aggageacct gaagaaatca agccaggtgo coaggacaga agccccagoo 660
ccagocteat ctacacccca ggagocttgg cctggcccta ccgccccag ccctaccage 720
ogoregreet gggetgtgga coetgegttt geogagoget atgee
<210> 22
<211> 1689
<212> DNA
<213> Homo sapiens
<400> 22
cyacycayay caycyccety yergyyccaa ycaygayccy ycatcatyga ticottoaay 60
gragtgetge aggggecage accttggggd ttddggctgc aagggggcaa ggacttcaat 120
stycemeter meatitioning getraction syggestates office office of the style of the st
stgggtgact gggtgctgag catcgatggc sagaatgcsg stagcetrae acacatcgaa 240
```

g¢tcagaaca agatcogggc ctgoggggag cgcctcagcc tgggcctcag cagggcccag 300

```
neggtteaga geaaaccgca gaaggeetee gesceegeeg oggaceatee geggtacacc 360
tttgcacca gegteteest caacaagaog geedggeet ttggggegee coogcoeget 420
gabagogood egeaacagaa tggacagoog otoogacogo tggtoobaga tgccagcaag 490
cageggetga tggagaacac agaggactgg cggccgcggc cggggacagg ccagtcgcgt 540
tectteegea teettgeees coteseage accesittes tgessgacce ggatgaggag 800
caccigasga aatomagoom ggigoccagg acagaagooc cagooccago cicatotaca 660
coccaggage cetggcotgg coctaceges occagoosta coagoogoos goostggget 720
gragacocts cytttgccqs gegetatgcc ceggacsaaa egageacsgr gctgacccgg 700
cacagocage eggecacgee cacgocgety cagagocyca cotocattyt ycaggoayet 640
geoggagggg tgccaggagg gggcagcaac aacggcaaga ctcccgtgtg tcaccagtgc 900
cacaaggtes teeggggeeg ctacetggtg gegttgggee acgegtacea cccgggaggag 960
tttgtgtgta gookgtgtgg gaaggteetg gaagagggtg gettetttga ggagaaggge 1020
gecatettet geccaecatg etatgaegtg egetatgeac ceagetgtge caagtgeaag 1080
aagaagatta caggegagat catgcacgec etgaagatga cetggcaegt geactgettt 1140
acctytycty cotycaagac goodatocyg aacagygoot totacatega ggagggogtg 1200
contattgog agogagacta tgagaagatg titiggosoga satgoostgg cigigactic 1260
aagategaeg etggggaeeg etteetggag geeetggget teagetggea tgaeseetge 1330
ttegtetgtg egatatgtes gstesseetg gssggsssgs eettetsete casqaaggse 1380
aggestatet geaagagsea tgeettetet satgtgtgag sessitistgs seasagetgs 1440
ogeggtggoo cotageetga ggggeetgga gtegtggeee tgeathtetg ggtagggetg 1500
geaatggtig collaacoot ggeboolgge oogagedigg geleddgigge eddigddaa 1960
coaccitate eteccacee acteceica beaccacage acaceggige iggecacace 1620
agececetti cacetecagi gecacaataa accigiacce ageigaatte caasaaatee 1600
<210> 23
<211> 22
<212> DNA
<213> Homo aspisoa
<4005 23
geaetgtget egttttgtec gg
                                                                  22
<310> 24
<311> 21
<213> DNA
<213> Homo gapiens
<400> 24
teettgetea eeteaeggge &
                                                                  21
<210> 25
<311> 30
c212> DNA
<213> Bomo sapiens
<400> 25
                                                                  30
tecteateeg ggtettgeat gaseteggtg
<210> 26
<211> 28
```

c2125 DNA

MO 00/99128	PCT/DE00/11664
<213> Homo sapiens	
<400> 25	
denceodece deparation neceders	2.5
<210> 27	
<211> 24	
c2125 DNA	
<213> Nomo sapiena	
<400> 27	
tecttgetea cetekeggge aceg	24
<210> 28	
<211> 22	
<212> DNA	
<213> Homo sepiens	
<400> 26	
greatechec reactatagy de	22
<210> 29	•
<211> 23	
<213> DNA	
<213> Rattus norvegicus	
<400> 29	
goggotgatg gagaataotg aag	23
<210> 30	
<211> 23	
<212> DNA	
<213> Rattus norvegicus	
<400> 30	23
atohtgtgge actggtggca tac	*~
<210> 31	
<211> 22	
<312> DMA	
c213> Rattus norvegicus	
<40D> 31	**
tgtgtcgggt cagcactgtg ct	22
<210> 32	
<211> 1620	
<213> DNA	
<213> Romo gapiens	

```
<400> 32
atggatteet teaaggtagt getggaggg eeageatett ggggetteeg getgeaaggg 60
ggcamggact teastgtgcc cotetecatt teceggetea etectggggg camageggeg 120
caggooggag tegeccytegg teacteegte cteaecatog ategegagaa teceggetage 180
etcacacaca tegasgetea gaacaagste egggentgeg gggagegeet eageetggge 240
eteageaggg cocagcoggt toagagtaaa cogcagaagg cotocgccc cgccgcggac 300
ecteogeggt acacettige accemagegte tecetemmes agaeggeeeg gecettiggg 360
gegeeecege cegetgacag egecegeaa cagaatggac ageegeteeg acegetggte 420
ccagatocca gcasgcagcg gctgstggsg ascscagagg sctggcggcc gcggccgggg 400
acaggocaqt egeptteett degeateett gedeacctea caggoaccga gitteatgeaa 540
gacceggatg aggageacet gaagaaatea agceaggtge ceaggaeaga agceeeagee 600
coagonteat etacacees ggageeetag cetageeeta eegeeeeaa eeetaceage 660
egecegecet gagetgtgga coetgegttt geogageget atgeceegga caaaaegage 720
acagigotga cooggeadag coageogged acquedage egotgoagag cogcacotoc 780
attotgcagg cagctgccgg aggggtgcca ggagggggca gcaacaacgg caagactccc 840
grgrgreace agrgeeses ggresteegg ggeegersee tggrggegtt gggeesegeg 900
taccaccogg aggagtttgt gigtagccag tgigggaagg tootggaaga gggiggotto 960
tttgaggaga agggddddat ettotgddda ddatgdtatg adgtgdddta tgdaddago 1030
tykycesagt gcasgaagaa gattacaggo gagatcatgo acgocotgaa gatgacotgg 1080
cacgtgcact getttacetg tgetgcetge aagaegeeca teeggaacag ggeettetae 1140
atggaggagg gegtgeeeta ttgegagega gactatgaga agatgtttgg caegaaatge 1200
catggotgtg acttcamgmat cqmcgctggg gaccgcttcc tggmaggccct gggcttcmge 1260
tggcatgaca cotgettegt etgtgegata tgteagatea acetggaagg aaagacette 1330
tactecanga aggacagged tetetgesag agecatgeet tetetestgt gtgagedeet 1380
totgoccaca gotgoogogg tggoocotag cotgagggor otggagtogt ggoocotgcat 1440
tt#tgggtag ggctggcaat ggttgcctta accetggete etggcccgag cetgggctce 1500
egggecoetg coeacccace thatcetese accceactee etceaccace acageacace 1560
gytgetggcc acaccagocc cotttcacct coagtgccac aataaacctg tacccagetg 1620
<210> 33
<211> 1665
<212> DNA
<213> Homo sapiens
<400> 33
egacgowyky chycecoty geogygocaa geaggagoeg geateatyga ticetteaag 60
gtagtgetgg aggggeeage acettgggge tteceggetge aagggggeaa ggaetteaat 120
gtgodddtat ddatttodog gotoactoot gggggodaaag oggogoaggo oggagtggod 180
gtgggtgact gggtgctgag catcgatggc gagaatgcgg gtagcctcac acacatcgaa 240
geteagaaca agateeggge etgeggggag egeeteagee tgggeeteag cagggeecag 300
coggiticaga gcaaacogca gaaggcotec gcccccgccg cggaccctcc gcggtacacc 360
tttgesees gegteteest saasaagasg geesgeset ttggggegee coogcoogst 420
gacagogoco ogoaacagaa tggacagocg etccgaccgo tggtcccaga tgccagcaag 480
eageggetga tggagaacac agaggaetgg eggeegegge eggggaeagg ecagtegegt 540
teetteegea teettgeeda eetescagge seegagttes tgeaagseee ggatgaggag 600
escriganga salcaageca ggigeceagg acagaagees cagececage eleatetaca 660
ecceaggage cetggéetgg centacegen conagenta coagengene genetgaget 720
giggscorig egittgeega gegetatgee eeggacaaaa egagcacagt getgaccegg 780
cacagocago oggecanged cangdegety cagageeyea cetebattigt geaggeaget 840
```

gerggagggg tgecaggagg pggdagdaac aacegeaaga ttercettte teaccagtge 900 cacaaggagg treeggggeg etacotggtg gegttgggee acgogtacca cooggaggag 960 tttgtgtgta gecagtgtgg paaggtettg geatagge acgogtacca cooggaggag 1020 pecatettet gecaccatg ctatgacgtg cgctatgcac coagctgtge caagtgcaag 1080 aagaagatta caggegagat catgeacgee etgaagatga ectggeacgt geactgttt 1140 acctgtgetg cetgcaagae gecoatcegg aacagggeet tetacatgga ggagggegtg 1200

cectatige agogagacta tgagaagatg titiggiacga aatgocatgg cigtgactic 1260 aagategacg cigggaccg citicitggag geeetgget teagetggea tgacacetge 1320 tiegtetgte egatatotea gateaacetg gaaggaaaga cettetaete caagaaggae 1380 aggeetetet geaagageea tgeettetet catgitgiag cecettetge ceaeagetge 1440 egoggtgge cetageetga ggggeetgga gtegtggee tgeatitetg ggtagggetg 1500 geaatggtig cettaaceet ggeteetgge cegageetgg geteeggge ceetgeeae 1560 ceaecttate etecaacee acteetee accacacage acaceggtge tggeeaeae 1620 aggeeeettt caceteeat geoacaataa acetgtacce agetg

<210> 34

<211> 223

<212> PRT

<213> Homo gapiens

<400> 34

Met Asp Ser Phe Lys Val Val Leu Glu Sly Pro Ala Pro Trp Gly Phe 1 5 10 15

Arg Leu Gln Gly Gly Lya Asp Phe Asn Val Pro Leu Ser Ile Ser Arg

Leu-Thr Pro-Riy-Gly-Lye-Ala-Ala-Gln Ala-Gly-Val-Ala-Val-Gly Asp-35 40 45

Trp Val Leu Ser Ilc Asp Gly Glu Asm Alo Gly Ger Leu Thr His Ile 50 60

Glu Ala Gln Asn Lys lle Arg Ala Cys Gly Glu Arg Leu 9er Leu Gly 65 70 75

Lew Ser Arg Ala Gln Pro Val Gln Ser Lys Pro Gln Lys Als Ser Ala 85 90 95

Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Pro Ser Val Ser Leu 100 105 110

Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala 115 120 125

Pro Gln Gln Asn Gly Gln Pro Lev Arg Pro Lev Val Pro Asp Ala Bor 130 135 140

Lye Gln Ary Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly 145 150 155 160

Thr Cly Cln Scr Arg Scr Phe Arg IIe Leo Ala His Leo Thr Gly Thr 165 170 175

Glu Phe Met Gln Asp Pro Asp Glu Clu His Lou Lys Lys Ser Ser Gln 185 190

Val Pro Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu 195 200 205

Pro Trp Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro

210

215

22D

```
<210> 35
<211> 20
<2125 DNA
<213> Homo sapiens
<400> 35
                                                                  20
gagooggoat catggattee
<210> 35
<211> 20
-212> DNA
<213> Romo eapiens
<400> 36
                                                                  30
getgeetgea caatggaggt -
<21.05 37
<211> 1456
<212> DNA
<213> Momo sapiena
<400> 37
cgaegeagag cagegeetts geegggeeaa geaggageeg geateatgga liteetteaag 60
gtagtgetgg aggggccage accttggggc ttdrygctgr aagggggcaa ggaettcaat 120
gtgecectot ocatttoeeg geteseteet gggggesaag eggegnøgge eggsgtggee 180
gtgggtgødt gggtgotgag categøtggm gagaatgegg gtagoeteac acacategam 240
geteagaaca agateeggge etgeggggag egeoteagee tgggeeteag eagggeeeag 300
coggithraga graaacogca gaaggitgrag accortgaca aacagoogct cogarnogrig 360
gtoccagatg deageaagea goggetgatg gagaacadag aggaetggog googeggoog 420
gggacagged agtegogtte ettergeste ettgeocade teacaggeme egagtteatg 480
caagaccogg atgaggagca cotgaagaaa tommyeeagg tgeccaggac agaagcocca 540
georeagost catotadere compgagose tygostygos stadegesse cagoestaes 600
agoogoogu cetgggotgt ggacootgog thtgeegage getatgooco ggacaaaacy 660
agracagigo igacocogos rageragero gecacgeeca egococióna gagerograce 720
tecattgige aggeagetge eggaggggtg egaggaggg geageaacaa eggeaagaet 780
cregtgtgtc accagtgcca caaggtcatc cggggccgct acctggtgge gttgggccac 840
gogtaccacc cggaggagtt tgtgtgtagc cagtgtggga aggtcctgga agagggtggc 900
ttetttgagg agaagggege catettetge coaceatget atgaegtgeg ctatgcacec 960
agotgtgock agtgomagam gamgattach ggcgmgatch tgcmcgccct gamgatgwcc 1020
tggcacgtgc actgetttad ctgtgctgcc tgcaagacgc coatcoggsa cagggccttc 1080
tacatggagg agggegtgee ctattgrgag egagactatg agaagatgtt tggcacgaaa 1140
tgccatggct gtgacttcaa gatcgacgct ggggaccgct tootggaggc cctgggcttc 1200
agotggoatg acacotgott ogtotgtgog atatgtosga toaacotgga aggaaagado 1260
ttotactoca agnaggarag grotetotge aagagecatg collectes tgtgtgagec 1320
cottetgecc acagetgeeg eggtggener tageetgagg ggcotggagt ogtggmentg 1380
cattletggg twgggctggc satggttgee ttaaccetgg mteetggece gageetggge 1440
teccgggccc tgccca
```

<210> 38 <211> 423

<212> PRT

<213> Homo aspiens

<400> 35

Met Aep Ser Phe Lyg Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phs 1 5 10 15

Arg Leu Gln Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg 20 25 30

Leo Thr Pro Gly Gly Lys Ala Ala Sin Ala Gly Val Ala Val Gly Asp 35 40 45

Trp Val Leu Ser Ile Asp Oly Olu Asn Ala Gly Ser Leu Thr His Ile 50 55 60

Glu Ala Glu Aen Lys Ile Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly
65 70 75 90

Let Ser Arg Ale Gln Pro Val Gln Asn Lys Pro Gln Lys Val Gln Thr
85 90 95

Pro Asp Lys Gln Pro Leu Arg Pro Leu Val Pro Asp Ala Ser Lys Gln 100 105 110

Arg Leu Met Glu Asn Thr Glu Asp Trp Arg Pro Arg Pro Gly Thr Gly
115 120 125

Gin Ser Arg Ser Pho Arg Ile Leu Ala His Leu Thr Gly Thr Glu Pho 130 135 140

Met Gln Asp Pro Asp Glu Glu His Leu Lys Lys Ser Ser Gln Val Pro 145 150 155 160

Arg Thr Glu Ala Pro Ala Pro Ala Ser Ser Thr Pro Gln Glu Pro Trp
165 170 175

Pro Gly Pro Thr Ala Pro Ser Pro Thr Ser Arg Pro Pro Trp Ala Val 180 185 190

Amp Pro Ala Phe Ala Glu Arg Tyr Ala Pro Amp Dym Thr Ser Thr Val 195 200 205

Leu Thr Arg His Ser Gln Pro Ala Thr Pro Thr Pro Leu Gln Ser Arg 210 215 220

Thr Ser Ile Val Gln Ala Ala Ala Gly Gly Val Pro Gly Gly Gly Ser 225 230 235 240

Asn Asn Gly Lys Thr Pro Val Cys His Gln Cys His Gln Val IIt Arg 245 250 255

Ala Arg Tyr Leu Val Ala Leu Gly Rie Ala Tyr Hie Pro Glu Glu Phe 260 265 270

Val Cys Ser Gin Cys Gly Lys Val Len Glu Glu Gly Gly Phe Phe Glu

275 290

Glu Lys Gly Als Ile Phe Cys Pro Pro Cys Tyr Asp Val Arg Tyr Als 290 295 300

285

Pro Ser Cys Ala Lys Cys Lys Lys Lys Ile Thr Gly Glu Ile Met His 305 310 315 320

Ala Leu Lys Met Thr Trp His Val Leu Cys Phe Thr Cys Ala Ala Cys 325 330 335

Lys Thr Pro Ile Arg Asn Arg Ala Phe Tyr Met Glu Glu Gly Val Pro 340 345 350

Tyr Cys Glu Arg Asp Tyr Glu Lys Met Phe Gly Thr Lys Cys Gln Trp 355 360 365

Cye Amp Phe Lys Ile Amp Ala Gly Amp Arg Phe Leu Glu Ala Leu Gly 370 375 380

Phe Ger Trp His Asp Thr Cys Phe Val Cys Ala Ile Cys Gln Ile Asn 385 390 395 400

LBU Glu Gly Lys Thr Phe Tyr Ser Lys Lys Asp Arg Pro Leu Cys Lys
405 410 415

Ser His Ala Phe Ser His Val 420

<210> 39

<211> 1575

<212> DMA

<213> Homo mapiens

c400> 39

ogacycagay cagegecety geoggyccaa geaggageeg geateatyga tteetteaag 60 qtaqtqctgg aggggccage accttggggc ttecggetge aagggggcaa ggaettcaat 120 gtgoccotet ccattleceg geteacteet gggggeaaag eggegeagge eggagtggee 180 gtgggtgact gggtgctgag catcgatggg gagaatgcgg gtagcctcac acacatcgaa 240 geteagaaca agateeggge etgeggggag egenteagen tgggeeteag cagngecoan 300 ceggitosga geaaacogea gaaggeetee geereegeeg eggaceetee geggiacase 360 tttgcaccca gegtetecet caacaagaeg gecoggeeet ttggggegee eeegeeget 420 gacagogoco ogoaacagaa toggtograga cocctgacaa acagoogoto ogacogotog 480 toccagatgo cagcaagcag oggotgateg agaacacaga ggactggcgg cogcggccgg 540 ggacaggoca gtogegtter ttregeater ttgcccaret cacaggosee gagtteatge 600 aagaceegga tgaggagcac etgaagaaat caageeaggt geeeaggaea gaageeeeag 660 occoagosto atotacades caggagoset ggeotggese tacegesese agesetasea 720 geogeoegee etgggetgtg gaccetgegt ttgccgageg etatgecceg gacaaaacga 780 gcacagtgot gaccogcar agcosgoogg coacgoodac gcogotgoag agcogcacet 840 ccattgtgca ggcagctgcc ggaggggtgc caggagggg cagcaacaac ggcaagactc 900 cogtytytca ccagtyccac aaggicatee gyggeogeta cetygtyyeg ttyyyeeaeg 960 cgtaccaecc ggaggagttt gbgtgtagec agtgtgggsa ggtcctggss gsgggtgget 1020 tetttgegga gaagggegee stettetgee eaccatgeta tgaegtgege tatgeeecca 1000 getgtgecaa gtgeaagaag aegattaeag gegagateat geaegeeetg aagatgaeet 1140 ggcacgtgcs ctgctctacc tgtgctgcct gcaxgxcgcc catocggaac agggccttct 1200

PCT/US00/11664 WO 00/66178

acatggagga gggcgtgccc tattgcgagc gagactatga gaagatgttt ggcacgaaat 1260 gecatggetg tgacticaag ategacgetg gggacegett cetggaggee etgggettea 1320 getggeatga cacetgette gtetgtgega tatgteagat caacetggaa ggaaagacet 1380 tetachecaa gaaggacagg octotetgca agagecatge etteteteat gtgtgageee 1440 ettetgeeca eagetgeege ggtggeecet ageetgaggg geetggagte gtggeectge 1500 atttetgggt agggetggea atggttgeet taseeetgge teetggeeeg ageetggget 1560 coegggeest geesa

<210> 40

<211> 153

<2125 PRT

<2135 Homo capiens

<400> 40

Met Asp Ser Phe Lys Val Val Leu Glu Gly Pro Ala Pro Trp Gly Phe 1

Arg Leu Gin Gly Gly Lys Asp Phe Asn Val Pro Leu Ser Ile Ser Arg

Leu Thr Pro Gly Gly Lys Ala Ala Gln Ala Gly Val Ala Val Gly Asp

Trp Val Leu Ser Ile Asp Gly Glu Asn Ala Gly Ser Leu Thr His Ile 55

Glu Ala Gin Acn Lyc lie Arg Ala Cys Gly Glu Arg Leu Ser Leu Gly 70

Leu Ser Arg Ala Gln Pro Val Gln Ber Lys Pro Gln Lys Ala Scr Ala 85

Pro Ala Ala Asp Pro Pro Arg Tyr Thr Phe Ala Fro Ser Val Ser Leu

Asn Lys Thr Ala Arg Pro Phe Gly Ala Pro Pro Pro Ala Asp Ser Ala 115 120 125

Pro Gln Gln Asn Gly Cys Arg Pro Len Thr Asn Ser Arg Ser Asp Arg 130 135 140

Trp Ser Gla Met Pro Ala Ser Ser Bly 145 150

<210> 41

<211> 24740

<213> DNA

<213> Homo sapiens

## <400> 41

nnnnnntgta tittatcats tittaasaat caasaacaa aaqqcagitq agqitaqqca 60 tggaggtteg tgwetgtaat wedageackk lgggaageeg aageaegigg atcacetgag 120 gtcaggagtt cgagaccage etgeccaata tggtaaaacc etgtetetac taaaaataca 180 saaaattago caggoatgat ggtgggcaco tgtaatooca gotacttggg agaetgagge 240

PCT/US00/11664

					tgtaateeca	
					catcetgget	
					cctggtggcg	
					aacctgggag	
					cangagtgag	
					gccgagetct	
					aaaggcagtt	
					gttgctatgg	
					tgotgaaggo	
					aggtggetgt	
					cagaagetgg	
					totgomgoto	
tengacacce	tcaagcacag	tgcctcttgc	ccageccage	actctcagtg	aabaacayoa	1020
					aagotggaac	
					daddagaddd	
gecagggcat	agaaatacaa	tgetetesee	tttgacttga	actcattccc	саддадасад	1200
gggagg¢¢¢¢	ctcaggatec	acagatgece	agtetecess	gaggggcctg	gtccccatgg	1260
аддаваястс	catctactcc	teetggcagg	aaggtaagtt	ggaggacgtg	сяядайсяйс	T350
					cctggccscc	
tgccacagtg	ggcgagtttc	ccgtggtggt	gcagaggact	gaggecgeca	cccgctgcca	1440
gatgaagggg	eeggeeetge	tegtgctggg	cccsdecacc	atccagctga	gggaggccaa	1500
					ctccgacaag	
					tgctaatccc	
					depasses	
					gggctgtggc	
					gecetgecee	
¢¢gecacggg	ceaectetet	deactocard	gecacccccg	gagagetteg	<del>ддадасдсса</del>	1860
ecaggacets	agccacccac	gtocaggaaa	atocacctgg	cedadcccdd	асссовдадс	1920
					egetteagtg	
tgcazgcgtg	ccagtgggcc	cccaggcast	dyacscoper	atgagaacct	gtgtgtgctg	2040
deadccadcc	ccacgotgca	caataaaasa	cctgageege	acdsäääccc	oddosáccác	2100
agececacaa	ceagteceat	ctaccacaec	adocaratare.	tgagetggee	cggcccggcc	2160
					99†99% <b>9</b> 99¢	
acaggeogee	¢tgaccctca	ggcaggtttc	яаддс <b>с</b> фадс	tggtgaccct	gctgmgtcgt	2380
gagoggagga	ядайсссяйс	cccttgtgsc	cggccctgaa	cacccaacs	agtggtggcc	2340
					cattgggccc	
atgcagacac	acgcotgtgt	ccaecctggs	ctgcaggaac	я <del>я</del> ддся <b>ддс</b> с	dccratagaa	2460
					taataaagot	
					cattgcggcc	
aacacaagca	ccctgtgccg	gttccagggg	cacaggigae	ctgggcctta	cetgeeacce	2640
gteggeteaa	<b>ecception</b>	gcagacagac	addaradayyy	tcattaggac	tccatgttgc	2700
tetgesegge	ogagtgacac	awaayaa cea	gcggagggag	ctgtgaggct	tacttgtcag	2760
					ctgggaggag	
g¢agtgggg	cggggtgcag	ataassagaa	tttcagagac	accetegice	aaaacacttg	3880
ttecetgetg	assetcease	astttgcaga	tacttctggg	PUCCCCEGGC	gtcagtctcc	2940
teatctgtaa	aggagagaga	accystgacy	tatcaggcat	aatcottgat	gagagtttg¢	1000
					agccttggtc	
					tecatgetge	
					acgcaagcgt	
catgecteca	gggctctgac	tgecoagece	ctcaccgcaa	etecacetea	getgeseses	3240
occttggcac	atcctgascc	teatttteet	gacggacaca	casttttgc	teteteetgt	3300
ccaagcetea	tectetgged	geesecteet	tecageteae	tteettagt	geggeeagta	3380
					gaggectetg	
cccaccatcc	cctctttgtt	cccatagte	cetteceet	grecetete	gtttcatctt	7 6 B O
eccaarctaa	caaagtcccc	ggccttgggc	gagccagacc	LCCCCAGCGC	ctgcacacaq	3540
ctgcccacag	ccadedaaag	ccatttaage	sgactgcctg	catecttett	ascagtgcas	3600

ggcaggcact	<b>ccctgcoscs</b>	agagaccctg	ttccctagta	gggcagettt	tataataaa	366D
agaacotost	gtstatesse	acccaatgtc	toctcacagg	catattgggg	aaacaggtca	3720
ggststscs4	eegtatotgo	aagtgtactg	geatecatet	gtettettee	tacccctacs	3790
gtagsascag	tgtctgtccc	cagetgtget	ctgatccogg	ctcctttcac	ctcagagett	3340
ggaaaattga	getgteccea	ctctctcctg	egeceattea	tectaccage	agottttoca	3 <b>9</b> 00
gccacacgca	aacatgetet	gtastttcac	attttaaacc	ttcccttgac	ctcacattcc	3960
tottoggoca	ectotgttto	tetgtteete	ttcacagcas	asactgttcs	amagagttgt	4020
tgattacttt	catttccact	ttctcscccc	cattetetee	tcaattaact	ctccttcatc	4080
cccatgatgo	cattatgtgg	cttttattag	agteacease	cttattotoc	ввансання	4140
					cccccgttac	
ttgctattcc	tectacetea	taacaatoto	etteccages	totactgctg	ccttctctga	4260
gttcttccca	gggtcctagg	ctcagatgta	gtgtagetea	accetgetae	acaaagaatc	4320
tectosasoc	ctctaaaaat	gtocatgoat	gttetgtgag	tgatctacca	agaaaataaa	4380
aasttttasa	DATCASSESC	ccatgcctgg	gecesaege	aggggctctg	atttcatcag	4440
tetestasst	ggattctggg	catecaddet	cactggattt	coggatgett	gtagtatgca	4500
acctanacta	prosecte	occtorocaa	gecagteatt	etecaggtat	cacegaccc	4560
ctaccta	stoacccoa	apotetotet	teagtgcaca	cctccccctq	agetecagat	4620
ttaggrafee	carteracec	cagacatota	gatotogaan	amacatetee	agatcccatg	4660
					tggacacaga	
daragasaa	9900999999	ctacaperat	309000000	Atteaaacte	acttattcat	ARON
					cgggcagtcg	
gewageacce	cccdwcdddc	anguicutag	eachanasan	9900449499	gcccttggca	4920
					ctagaaccag	
ascecaggac	conggggade	cageatggea		Cattacadac	ctgagetget	5770
trgtttccta	äääettrord	rgaracccca	aggggacogc	danaman	gtccttggaa	5160
#CC#CC#GG#	cgccagaggc	ccáscccan	gccccaggag	terespectate	cagagggggc	2720
tgggtocttg	¢¢¢aggccca	gergeraege	#\$\$\$\$\$ccacc	tgtgetgggs	ggctccccag	5760
gggacacaga	ccagageett	gesceagee	BEGGETCGGG	agccegggt	cotototgat	5250
ggaggactgc	caggaccccc	aggetgeege	Ctattacert	GGEGRITIGG	tgtttcactt	5140
tgtcaatcct	tectttcttc	gtgtgttcat	teacatecae	rataractaa	ccctdddass	5400
atgttegata	agacacatta	gctgtgtgt¢	ttcattgtcc	taacaaagaa	cacaccctgg	5460
asagagcacc	acsadadec	eccattcccc	catetecete	cacacatgga	etctggagat	5520
gcetttteea	catcomgatg	tototggtg¢	tgtgggattc	ttaaataaac	azacatttca	5580
tacagaatgt	<b>andaparta</b>	agstgctatg	gggaaaagta	aagcagaaggg	agggcctagt	5640
gtgtgatgcg	ggtgaggcat	ccagggattg	ctgtttcage	tgtgatcagg	aaaggecetg	5700
ggaggaggcc	acatotgago	agagacctaa	ataaag&tgg	aaacctgttg	ctgagatatc	5760
tggagaagtg	tttcaagggc	caaacscaa	gcatggtggc	tracgrotyt	aatcccagca	5620
ctttgggagg	ccaaggcagg	tggatcgctg	gaggtcagga	gtttgagag¢	agcetgacea	5880
acetggagaa	Accoratcts	tactamment	atesasette	teegggeatg	gtggttcatg	5940
cctgtagtcc	cagetacteg	ggaggttgag	gcaggagaat	cacttgaacg	tgggaggcag	6000
aggttgcagc	aageegagat	сасассвесьс	Chetechgee	tggatgacag	agogagaete	6060
cgtctcasaa	ававававда	ваадаааааа	gazaattees	gasaagtgtt	tcaagcaggg	<b>5120</b>
gaactggcaa	gtggagaggc	сстдаддсяд	aaatatgott	ggcstgctgg	aggaaatgtg	6180
agtgaggagg	teagggtgge	tggagtggag	ggagogagtg	gtaggagtca	gacccagttt	6240
					tgageccact	
					atatcacttt	
ggetgetteg	tgaagactet	ggaaggggca	agggtgaaag	cagggatgcc	cgttaggaga	6420
ccgttacagg	ggogcagqca	caasatgg¢a	gtggctggga	caatggtggc	agcagcggtt	6480
agatgtgaac	atgttgasga	tggsatttgc	agaatetggg	ggaggacaga	ададваадда	6540
tancticate	atttetecte	aaccagttgq	atamatgttg	gtggcacttc	ttgaagtgag	6600
gazogaotta	ggaaggtggg	ABBOTCACAA	gtttgaattq	ggccatgatq	gtctgageta	6660
cctagtarag	tagtteene>	acctititoo	Caquaggac	cactttcata	gaagacaatt	6720
					cagtacattt	
					tcacccacac	
					tcamgtcatt	
etectacete	agectetes	at and cade	ttstaggcat	atacannace	acgcccaget	6960
2202032016	agovococaa	2 ~~ 3 ~ - 2 2 2 4				

41141.14			<b></b>			
					gtetegaact	
					tacaggegtg	
					acctagated	
					etgeegetge	
					rccraacagg	
					ttttggataa	
					gostttgest	
ataggtggtg	ttgaaagcca	caggacagta	cggggtctca	addåätävät	ataaagagag	7440
gcgacatcag	atggccaagg	ccadeddcad	<b>vààsab</b> artilà	gaaggagggg	ccagtggggc	7500
аддрудалас	tgtgaagcca	<del>adagaaaaada</del>	gtgtttcgcg	gaaaaggat¢	aacctggacc	7560
agtgetgecc	ctaggcaggg	CEGGAtGSAS	Cttaaccacc	acggattcca	tggcccatg	7620
geetecagge	cacaggggac	cttgagaaga	gagateteag	gggacgggtg	cccaccadag	7660
ceegactgge	<b>BtggcttcaB</b>	gagataactg	angganagoa	agtggagacg	ogataaacag	7740
acaactecet	ggaggaattt	tactotogag	aggagaatta	aagggtagta	getggagagg	7800
gatgtggggt	caagagaagg	tctttaacga	cgagaactct	cacggcggtt	tgtgcagaac	7860
					gtcctaggag	
					ctcaccccga	
					acgeggggee	
					\$¢\$\$\$ac\$3c	
					ecgetectge	
					catatacaca	
					gggttgcgcc	
					- <b>ttttc</b> ggagt-	
					ggtccgcttt	
					ccctaaggag	
					ggtcggcgag	
					agetgggtgg	
					tggcetggtg	
					teagaacact	
					rgaccagace	
					geeeggggee	
					tecegetoge	
					tgcggcgggg	
					Bccca33c33	
					gaddcaccac	
					ocedocadas	
				_	cocaggggtc	
					ccgctgcsgc	
					coctoccete	
					gtgtcagctt	
•					ccfåååååôcä	
					gatggettgg	
					cctcaagcot	
					ctgtctttcc	
aangtgaggt	ccgccagcca	gcagcccagg	ccagcctgac	aaaatacctg	cctcctatgg	9660
cttgggcgtg	ctcaggggct	gcccgtgcct	gcctggcccc	tgtccaaggc	tggtatectg	9720
agctggcccg	geetgeetge	etgeeegeee	accatgctgg	ceaeteaeet	tetettetet	9780
¢etoteagga	geeggeatea	tggattcctt	caaagtagtg	ctggaggggc	cagcaccttg	9840
gggctteegg	ctgcaagggg	gcaaggactt	caatgtgccc	ctctccattt	cccgggtgag	9900
					ccagagggtg	
					gtctaggtcc	
ggagatgtag	tggtacctgc	ctgccacsag	gactcccsat	gaggtggata	ctgggaggga	10060
					ctcctgtggc	
					tttcaggagg	
					tgtgactata	
					gtttttccag	
					•	

					ggtgaagcag	
					gtetetecee	
					tttgaoocca	
					aagageetge	
cctggtgtgg	gtggagctgg	aggetggete	ctccctgtag	cagggggatt	gccttataag	10520
cccaagaatg	cagodddaeg	otgggatggo	cascagtggc	tgeggtetge	agagotgmaa	10660
agggetggee	taggcctggc	cccctgaacc	ccactggtgg	gcctctcagc	tggtcaccag	10740
					decagaging	
					ggggtgggga	
					tgggcaggca	
					togtggcctg	
					gggtgggceg	
					gcccogccag	
					cctcacacag	
					tgtctgggga	
					gttgggggca	
					ctagetgate	
					totgagttag	
					geasyggetg	
					Eggctgctca	
					teettggtt¢	
					ccacccaccc	
					catectetgt	
					ccctctccag	
					tcccaggage	
					accetgacet	
					ggggttgatt	
					ggetgaeeee	
					agggt cagag	
					ttaacctctc	
					ggttgttgga	
					gtettettat	
					acgcaagcaa	
					etgtgeetgg	
					gcctcagctc	
					ggsccccccc	
					ctacttgatg	
					cagtcaagct	
					agggtcacco	
teteagggee	tgggagatac	ccmatcccag	acattaaaac	tgccagtagc	ccctctacct	12720
					gaactotoac	
					aaagtcattc	
ctgatggtct	cageteatge	tggtggcag¢	catttctccc	agectactgt	ctctactcat	12900
tgccacagga	accagggact	cccagctcaa	gagcctgaag	gattggggtc	aggggaaatt	12960
ggcagtcgag	ggcttgggag	tgacagecat	gtatggccta	cgsagtccca	getgteaact	13020
taggtcccat	tcaggcagtg	ttcacaggga	accgggagat	aacagggcct	gttectggct	13080
ctcaaagggt	cccagcagac	oddtatagat	ggçqqqqqqaq	agggtgctgg	ggggtgagag	13140
					gccagagggc	
					aggcaatggt	
		•			atagcaattt	
					gggaagetea	
				•	tgactgggaa	
					agcagotgoc	
					ttgggtgggg	
					gtcactttgc	
					gccgtgggtg	
<del>-</del>	•					-

netenatant	maddatonat	gacasanata	caaataacet.	cacacacate	gaageteaga	13740
HCC293CGCC	gagearogae	managedetta	acctagacet	cedcecotat.	gegggtggae	13800
atamatamat	32000303033	tanceatana	gatecetoco	accernaces	ctgtcacgct	13860
aradarosaa.	åcåceracae	ageographes	Generalise	CAGRADOTAC	gaggetggee	23920
	accadegacco	astatagaet	topacoocea	oncetacted	ccetcctggc	13980
Salarareca	4944364966	ggtgagggaa	artagapatae	224442442	getaaggegg	14040
acatecers99	ataneattae	accadenact	actorator	retares ces	cgcatgtecg	14100
gceragggee	constatete	tecangggget	acctaccase	accacacta	cgtgctcgcc	14160
					atgggoocct	
cacacacatas	cagoccoogg	desegrated	techantara	tatanaaaat	8888468888	14280
caaaaaaacta	gagergerge	natananana	totateteta	cccanacete	ccaggecccc	14400
apol Laboro	geeegeeege	ettteaana	apacketors	+ Magaabaaa	egecedeged	14450
geggaecete	egegg cacae	Coccedences	agogcocce	post soster	ggeceggeet	34520
rrgggcgccc	cedecedera	acagegeeec	geaguagaau	add recared	geccétgeéé	14586
deseñedeen	angenaticag	Secretain	geoceaegee		tgctgctcag	14560
reraraerae	accecsare.	ggtggaaccg	Lgaggeacge	accordacad	ceggggtggg	14040
\$ctacadaca	cagggccccc	cocdsdacta	tageacorra	cagggcaccg	cctggggagg	14700
ggtetetgaa	pastaccaca	concerdera	acaacraaas	åt tödåt rär	ggtgtcgggc	14000
<b>cagetgagee</b>	ccagacacte	agtgccgcct	tgtccccggc	tgttctgacc	ectecceste	14620
tttcttcctc	tectgtgtet	gtccctttgt	cectttatet	gtetgtetgt	cttatttcct	14880
tcacaggtgc	agacccctga	camptragtg	agecccecte	tgcctgtgcc	tttcttcttc	14940
cttttggcac	tetgggtgge	ggeceeteee	caccctggct	gecetastat	ecactteges	15000
ctcctgtcct	ctcacctacc	cgcccagcag	ggeteetgge	ctcaccctta	cccactocct	15060
cccatcactg	taacccaaac	ccacatgcac	caaatentgg	gaggggctgc	peccacegee	15120
cacececagt	gtggggtt¢t	gagecacace	ctccccacag	acagecçécte	egadegetgg	15180
toocagatge	cagcaagcag	cggctgatgg	agaacacaga	ggactggegg	ccacáaccaa	15240
qqacaggcca	gtogegttee	ttecgcatec	ttgcccacct	Cadaggdadd	gagttcagta	15300
agtgccagcc	cagggcaggg	ggtactttcc	tegeccccag	cccaggcgtg	atecetgaee	15360
ctqtgtcttt	tttggtcaat	geetgeetet	geteteteag	tgcaagaccc	ggatgaggag	15420
CACCEGRAGE	astcaaggta	сарруасруд	caccagocoo	totoccacct	catgactatt	15480
ccattccage	tactgeectg	Egictacted	tgaggeteee	agetggggct	cteaattctc	15540
ecttecttee	tteetteett	cetteettee	tteetteett	cettecttec	tteetteett	15600
ccettectec	tteetteett	ctttcatttc	tteectcect	dettecttee	atcatcacte	15660
ectacctccc	ttepatetet	cetteettee	acttottoct	coctetetet	ctgcccctca	15720
gagaaaaata	tetectogas	ctocapagee	cacqctacac	gragecticeag	gactggcacc	15780
					cageoggetg	
cectetetete	charactatt	ctattentes	tocccaddd	accecttag	tgcctccage	15900
ttetaertae	ctraccecce	ctttcataca	ectaacctaa	geeteetget	<del>ô</del> 墢¢åäccc	15960
torageotac	astagattes	tetapeacta	contractor	htesectata	gtactgctgt	16020
ctaccaante	totastocat	enotenane	atasacatto	teapecetoe	ttgctctgcc	16080
tetettetet	ctttstaase	ctantttaca	tttagagtat	agaggietht.	egtggttegg	16140
antatttaa	ccccgcggab	ottattetea	ntara-para	ataceteaca	asggggccct	16200
accercingg	Cocceptore	ttpstmadet	2022202222	20044453	cctcagaagc	16260
geoungeean	cacagacggc	coccegacac	gaggggagee	cate ant ma	cettactena	16320
999×9cccc3	eetegretee	cagucagaga	coge action	guesactus	cattgctagg	16380
ageagetgee	cccccatcc	ecaeeeeage	gggacecgga	ALCOHACTOS	geagttteca	36040
caccccat	errencede	ggggccagca	ggacceagge	+9999999+99	ggccatgtca	16200
ggaagereag	ccargcaggg	ccttdaardd	CHGALPLIGE	agecaggogo	ccaggacaga	10000
#dccccadcc	ccagcctcat	CERCACCOCK	9929000193	cerdardaga	gggagtgggc	702DC
rcgggcctgg	gcaagggtgg	desdecrees	agggcarggg	986886888	ttotetesge	1554V
rgeetgggge	rccacccccg	cccctgggg	recetagats	eccetteaga	gtcactttcc	TOORA
ccdacadacc	ctaccgcccc	cagecetace	ageogecege	cccgggctát	ggaccetgeg	T0340
tttgccgage	gctatgcccc	ggacaaaacg	ageacagtgc	rgecccggca	cagecagecg	16800
gccacgccca	caccactaca	dadacacaca	tccattgtgc	aggcagctgc	cadadadage	16860
<b>co</b> eggegggg	gcagcaacaa	cggcaagact	cacatarare	accagtgcca	caaggtcatc	16920
cggtgggtgg	cotgttoctg	teegaccetg	gettteecat	cethosácec	agecceacet	16980
gtetgeceae	ctgtcttgcc	teagetgega	ctgggggaa	taaggattca	gtteteaget	17040

				astoccacgo		
ageccaceca	carcaactgc	tagcagcatc	tgecgtggeg	aaatageega	адрдссаасс	17160
				сваадвадда		
cccagcgcgt	etgeesetee	cteetgetet	etgteteeet	ctgctctcag	agcatacagg	17280
				geageateaa		
gcatcagcgt	geeggattet	agagoottoo	teattegoeg	geetggeetg	ctctcatctc	17400
tgtcagotct	tttttttt	tttttgamac	agagteteae	tgtgttgccc	acgttggcgt	17450
				gggttcaaga		
cctcagcctc	ctgagtagct	gggattacag	geaccegeea	ccatgcctgg	ctaattttgt	17580
atttttagta	<b>gagacgggg</b> t	tttaccatgt	tggccaggct	ggtctcsaec	teeteacete	17640
aggtgatoto	aggeotgeet	tggcctccca	aagtgetggg	actacaggtg	tgagccactg	17700
				сводссвосв		
gctccagggt	ttgcccaggg	geggeteage	ttttatgagg	ctccagtcgt	cagocottec	17820
				ccagcagttc		
				aageetgtet		
gtggaaaatg	gggagaatga	actgtccctc	ccaaggcccc	ctggtgggtg	gtcagtcasc	18000
ctgaaggggg	ddaagacccc	cacctctctg	egtgtgetee	ctetgaecge	tetegeetee	18060
ctgcaggggc	cgctacctgg	tggcgctggg	ccacgcgtac	caccoggagg	agtttqtqtq	18120
				gaggagaagg		
				gccaagtgca		
				aagootgtoa		
				gagcctccgg		
				"agtcaggatg		
				246c938938		
				cctgccctgg		
				catactacta		
				tggtgatgtt		
				tggtggctta		
				gacccagget		
				tgcaagatga		
				aggagaagaa		
				cetetgettt		
				ggctgggaaa		
				tcagtcggca		
				tgggaggtcg		
				tggtgmaacc		
				agteceaget		
				gcagtgagcc		
cactgeette	cageetgega	gacagcaaaa	ARRABARA	<b>СВВЕБВЕБББ</b>	аажаходдес	19300
				ggccgagatg		
				accecatete		
				crcdddagdc		
				agatogogoo		
				аваадааада		
toccagoott	gggeteetge	cetaceagee	cacacccctg	gtagagcete	ctctcccace	19740
				cctctasaac		
				atttctgtga		
gtgtgagcag	gggctagttg	ctctctctgg	gcctcactga	agagaagggt	ggcactatgo	19920
				caggtgttgt		
				cttgcttggc		
tcattttaas	atcagtagca	BacBatttaa	aaaatcagaa	gatttgcctg	cstgatgcsg	20100
				tgggaggatt		
				gtttctacea		
				cccagetact		
agggaaagtg	agatotootg	ctttttattt	ctttatgtat	aatgataggg	tettgetetg	20360
ttgaccagge	togagtycag	tggcatgatc	actgctcact	geagecttga	teteetggge	20400
		_	_	<b>-</b>	~~~	

				******		20150
tcagaggate	ctccacctc	agcotoccaa	atagerages	cragaggrac	ccaccagcat	20520
geteageaga	Ettttaaacc	ttttgtaga	gacgaggett	tanananana	decapgetgg	20220
totogaadtd	erggeerega	Bedarecree	caccttggee	сесеннаден	ctgggattac	20500
agacgtgagc	cactátãoco	ageagattte	LCCLLBacac	cragarccca	gcctgagcca	20040
88cs33cec	cotgaatgaa	ccagtagtac	tactcccada	agaagaggcc	ctcctccgtg	20100
tgacacagte	conserrage	corracaggg	accegacceg	âàscccccă	atttaescto	20700
9838649265	tcataacagc	occacaagge	tgggsccage	tececagete	acaagggaag	2002A
asaccaagac	ttgagaaggt	caaggcotgg	ccagacccac	acateergya	ccctcatacc	20080
geet egagge	cocatgotad	ettatagarg	ctccagatgt	gaacactget	ggccctggct	20340
ggeeeegget	\$\$cccaa38	gtcctaggga	rgaacageee	*gcccaggga	gageteagee	21000
cettgtgcct	ctgccccttc	casectecta	cggaggccag	togacteace	cacaaagggc	22120 21000
caggcactgt	ggggatagat	cagotaacaa	aacagttgae	gerteergee	cttctgggcc	21120
<b>ttacatttt</b> g	getggaagaa	dedddagagad	geagaetgea	agcantaagc	gcaataagta	21180
ggttgcctgg	angtastgtt	agateaegtt	peggaaaaca	ggaaagagca	gagogacaag	21340
tgatggggtg	catastacaa	dasadacedc	tggctgctgc	täätätääte	agagtgggcc	21300
ctcatggaga	agactgcatt	cgagcagaaa	crideedadd	acasaaasra	agectagaga	21300
tatctggggc	agag¢agtcc	aggcagaggg	gacageeggt	gtcsagecca	ggacaggagt	31420
ątg¢čtggtg	tgecagttte	aggdaagg	ccagtgtgca	gaggcaaggt	радаводска	21480
<i>2</i> ддадарсар	pagagagac	gggtgggaac	gapgtcagac	etgetggeet	ccagcototg	21540
catgggggctt	ggetettget	gggagcaatg	ggaagcagta	cacagtttca	tgcaggggga	21600
gaaggeetgt	ettgggttgc	aggggcacgc	tgtggeaget	gggatcagag	agaggagett	21660
gtaggccagt	tgttatgtgg	teceacggge	cagatggcca	tggettacet	cacttcaggg	21720
aggetgtgag	aagcactcag	aatotggatg	-tgcattgggg-	gtgggcccca	ctggatttcc	21780
tggtggacct	ggtgtggggk	gtgagaggag	ggtgtgtttg	getgeageag	acaggagaat	21040
ggagttgcca	tecgegtgat	ggggatggct	gtgggaggag	aggtttgggg	tgagggaato	21900
aggaectgeg	tgetggaeat	ggcaagtctg	aaggogoagt	ggtogtdoac	tragagarct	21960
tagagttaga	gatggaggtg	tgggagteet	gaacagbtag	atgtagtgtt	taccgcgaga	22020
addaecaddd	cttgcggcca	geeeteetgt	gttcccgtga	сссадедств	ggcaggaggg	32080
acctanacct	qccqaqtqac	taggaeetee	tt.ccaggaga	teatgeapge	cctgaagatg	22140
Acctograce	tgcactgctt	tacctgtgct	gootgeaaga	concestees	gaacagggcc	22200
ttotacatgg	aggaggggt	gecetattge	gagcgaggta	cccactggcc	agtgagggtg	2226D
angagggatg	gtocatoggg	caggoatgaa	tecaggteet	ctttctctct	geocceatte	22320
teagactatg	agaagatgtt	tggcacgaaa	tgccatgget	gtgaettess	gatugaeget	22300
aggaccact	tectagagge	ectgggette	agetggesty	acacctgctt	cgtctgtgcg	23440
gtgagagggg	execetega	actgageeee	sageceaccy	gecetetett	cattccccag	22500
gagatgcagg	appagetton	aaggggcctc	tectgetgee	cccaacccca	tütilactüää	22560
cotttactat	ccttagatat	gi.cagatcaa	cctqqaagga	aagacottot	acticcaagaa	32620
pgacaggeet	ntictocaaga	qccatgcctt	chelicatgtg	tgagecectt	etgeceaeag	226B0
ataccacaat	gapecetage	ctgaggggcc	togastegte	gecetgeatt	tetgggtagg	22740
octograato	attoccttaa	ccctqqctcc	tggggggggg	ctggggetee	ctgggccctg	22800
CCCCBCCCBC	cttatectcc	caceccaete	cctecaccac	cacageacae	cgatgctggc	22660
CACACCACC	cecttteace	tecartores	ceataaacct	gtacccaget	gtgtcttgtg	22920
taccettece	ctatacatec	deaddadeac	aatttgagge	acgtggcagg	gtggagagta	22980
Agatogtttt	ctteggeteg	centetecat	gatecteata	atgeagaeat	ggcgggctca	23040
touttautou	amradutaca	decaspance	catotoccao	accedatace	cacagacatg	23100
-230003033	ctontetone	ctoart tres	gottagagaa	geotaettae	gaagggtagt	23160
tagggageea	met cat occt	otgatengan	caetttogaa	GGCCBBGGTG	ggcagatcgc	23220
ttweamtrem	gottered	cetestacee	esceccatos	ascadentet	ctagtaassa	2328B
togaggreag	gageecgaga agageegatat	octorings-t	certotaate	reagreacte	aggaggotga	23340
wareasast	agoogagege	stannayota.	centtacea	tnagetgaga	teacaccact	23400
Agenadasas	etabetgaac	ordinagency.	2002ccdogd	Vt t CRasass	teacaccact	23460
gogogogage	Coddd radag	24c2Ana2aA	catacateta	cadapatate	gt,gtggasaa	23520
офинанияня	Aggadacade	tecterates	Paraderend	oringwowed	asdasdacae	23580
caycaatggc	catagosati	recessassas	20000000000	topeanoran	tagaggtagg	23640
ggctggagag	ggtageeece	agg cereggg	wane une use	thtepaster thtepaster	tagagetggg etttetette	23700
gergeetett	daddrreddd.	cadadesaaaa	-acayeagay	PECARGORA	gtttgtgttg	23760
Lattgcagaa	ಇರದಿನಚಿರಿದ್ದಾದ	ядинсястда	Rahvacacua	Pedagadacr	ਫੋਰੋਕੋਬਫੈਰਬਰੋਪੋਤ	22700

```
gttgcagcag ggcctagggg cgggggccag gcaagggagg ggcagagagt aatatggcag 23820
agatgggacc cagtggcagg tocgggggat gagggatgga gagaaggaca qqaqcqttqc 23880
caggeatetg geetataeca gaeatgetea egetgtetee egegaaeete etageaaeet 23940
tgogoogitg telgeaatea ettatteeat titttetiit tiaaettiaa tittittigi 24000
ttttaagaga caggatotoo otaggttgoo ogggotggtt toanactoot gggotcaago 24060
aattottoot oottagoood aaagtgotgg cattacaggt gtgagocace atgootggee 24120
carttatttt otagatgagg cacagaaaga tigggagact igaccaaggi cacgoigtea 24100
ttgagocakg agocagaeta gaatocaggo otgaagotgg gtgogotgto ocaggaetgg 24240
etggcactga gtaccatttg ccagcgagca tetetotggg aagetgactt etgecoggta 24300
cotggaggar tgtagacett ggtggtggeg cegtcactet ggggetteet geeteccaet 24360
gatgeeegea eeacestaga gggastgtsa teteteetgt eesaageetg gaetggaaag 24420
actgaagage agcottaagt aggocagged agotcagtgt gocatggotg coogtootto 24480
agtggtocot ggcatgagga detgdaadad aletgtlagt etteleadea ggcoottgge 24540
coggtopert ttaagagaog agaagggotg ggoacggtga ctcacacete taatcccage 24500
sctttggaag gdtgaggetg gagaaggget ceagettagg agtteaggae eageetggge 24660
aacatggige gaccotgitt tgittigitt titgittitt igagatggag toitgotoig 24720
tegoccagge tegestgcas
                                                                  24740
```

<21D> 42

<211> 25

<212> DNA

<213> Rattus norvegicus

<400> 43

geactacett gaaggaatee atggb

25